

HIGHER SECONDARY FIRST YEAR ONE MARKS**REDUCED SYLLABUS-2021-2022****UNIT - 1. BASIC CONCEPTS OF CHEMISTRY AND CHEMICAL CALCULATIONS**

- 1 40 ml of methane is completely burnt using 80 ml of oxygen at room temperature. The volume of gas left after cooling to room temperature is
 (a) 40 ml CO₂ gas (b) 40 ml CO₂ gas and 80 ml H₂O gas
 (c) 60 ml CO₂ gas and 60 ml H₂O gas (d) 120 ml CO₂ gas
- 2 An element X has the following isotopic composition ²⁰⁰X = 90 %, ¹⁹⁹X = 8 % and ²⁰²X = 2 %. The weighted average atomic mass of the element X is closest to____
 (a) 201 u (b) 202 u (c) 199 u (d) 200 u
- 3 **Assertion :** Two mole of glucose contains 12.044 X10²³ molecules of glucose
Reason : Total number of entities present in one mole of any substance is equal to 6.022 X 10²²
 a) both assertion and reason are true and the reason is the correct explanation of assertion
 b) both assertion and reason are true but reason is not the correct explanation of assertion
 c) assertion is true but reason is false
 d) both assertion and reason are false
- 4 Carbon forms two oxides, namely carbon monoxide and carbon dioxide. The equivalent mass of which element remains constant?
 (a) Carbon (b) oxygen
 (c) both carbon and oxygen (d) neither carbon nor oxygen
- 5 The equivalent mass of a trivalent metal element is 9 g eq⁻¹ the molar mass of its anhydrous oxide is
 (a) 102 g (b) 27 g (c) 270 g (d) 78 g
- 6 The number of water molecules in a drop of water weighing 0.018 g is
 (a) 6.022x10²⁶ (b) 6.022x10²³ (c) 6.022 x 10²⁰ (d) 9.9 x10²²
- 7 1 g of an impure sample of magnesium carbonate (containing no thermally decomposable impurities) on complete thermal decomposition gave 0.44 g of carbon dioxide gas. The percentage of impurity in the sample is
 (a) 0 % (b) 4.4 % (c) 16 % (d) 8.4 %
- 8 When 6.3 g of sodium bicarbonate is added to 30 g of acetic acid solution, the residual solution is found to weigh 33 g. The number of moles of carbon dioxide released in the reaction is
 (a) 3 (b) 0.75 (c) 0.075 (d) 0.3
- 9 When 22.4 litres of H_{2(g)} is mixed with 11.2 litres of Cl_{2(g)} each at 273 K at 1 atm the

- moles of $\text{HCl}_{(g)}$, formed is equal to
- (a) 2 moles of $\text{HCl}_{(g)}$ (b) 0.5 moles of $\text{HCl}_{(g)}$
 (c) 1.5 moles of $\text{HCl}_{(g)}$ (d) 1 moles of $\text{HCl}_{(g)}$
- 10 Hot concentrated sulphuric acid is a moderately strong oxidizing agent. Which of the following reactions does not show oxidizing behaviour?
- (a) $\text{Cu} + 2\text{H}_2\text{SO}_4 \rightarrow \text{CuSO}_4 + \text{SO}_2 + 2\text{H}_2\text{O}$
 (b) $\text{C} + 2\text{H}_2\text{SO}_4 \rightarrow \text{CO}_2 + 2\text{SO}_2 + 2\text{H}_2\text{O}$
 (c) $\text{BaCl}_2 + \text{H}_2\text{SO}_4 \rightarrow \text{BaSO}_4 + 2\text{HCl}$ (d) none of the above
- 12 The equivalent mass of potassium permanganate in alkaline medium is
- $\text{MnO}_4^- + 2\text{H}_2\text{O} + 3e^- \rightarrow \text{MnO}_2 + 4\text{OH}^-$
- (a) 31.6 (b) 52.7 (c) 79 (d) None of these
- 13 Which one of the following represents 180g of water?
- (a) 5 Moles of water (b) 90 moles of water
 (c) $\frac{6.02 \times 10^{23}}{180}$ molecules of water (d) 6.02×10^{24} molecules of water
- 14 7.5 g of a gas occupies a volume of 5.6 litres at 0°C and 1 atm pressure. The gas is
- (a) NO (b) N_2O (c) CO (d) CO_2
- 15 Total number of electrons present in 1.7 g of ammonia is
- (a) 6.022×10^{23} (b) $\frac{6.022 \times 10^{22}}{1.7}$ (c) $\frac{6.022 \times 10^{24}}{1.7}$ (d) $\frac{6.022 \times 10^{23}}{1.7}$
- 16 The correct increasing order of the oxidation state of sulphur in the anions SO_4^{2-} , SO_3^{2-} , $\text{S}_2\text{O}_4^{2-}$, $\text{S}_2\text{O}_6^{2-}$ is
- (a) $\text{SO}_3^{2-} < \text{SO}_4^{2-} < \text{S}_2\text{O}_4^{2-} < \text{S}_2\text{O}_6^{2-}$ (b) $\text{SO}_4^{2-} < \text{S}_2\text{O}_4^{2-} < \text{S}_2\text{O}_6^{2-} < \text{SO}_3^{2-}$
 (c) $\text{S}_2\text{O}_4^{2-} < \text{SO}_3^{2-} < \text{S}_2\text{O}_6^{2-} < \text{SO}_4^{2-}$ (d) $\text{S}_2\text{O}_6^{2-} < \text{S}_2\text{O}_4^{2-} < \text{SO}_4^{2-} < \text{SO}_3^{2-}$
- 17 The mass of a gas that occupies a volume of 612.5 ml at room temperature and pressure (25°C and 1 atm pressure) is 1.1g. The molar mass of the gas is
- (a) 66.25 g mol^{-1} (b) 44 g mol^{-1} (c) 24.5 g mol^{-1} (d) 662.5 g mol^{-1}
- 18 Which of the following contain same number of carbon atoms as in 6 g of carbon-12.
- (a) 7.5 g ethane (b) 8 g methane (c) both (a) and (b) (d) none of these
- 19 Which of the following compound(s) has /have percentage of carbon same as that in ethylene (C_2H_4)
- (a) propene (b) ethyne (c) benzene (d) ethane
- 20 Which of the following is/are true with respect to carbon -12.
- (a) relative atomic mass is 12 u
 (b) oxidation number of carbon is +4 in all its compounds.
 (c) 1 mole of carbon-12 contain 6.022×10^{22} carbon atoms.
 (d) all of these
- 21 Which one of the following is used as a standard for atomic mass.
- (a) $^{12}_6\text{C}$ (b) $^{12}_7\text{C}$ (c) $^{13}_6\text{C}$ (d) $^{14}_6\text{C}$

UNIT – 2 QUANTUM MECHANICAL MODEL OF ATOM

- 1 Electronic configuration of species M^{2+} is $1s^2 2s^2 2p^6 3s^2 3p^6 3d^6$ and its atomic weight is 56. The number of neutrons in the nucleus of species M is
 a) 26 b) 22 c) 30 d) 24
- 2 The energy of light of wavelength 45 nm is _____
 a) $6.67 \times 10^{15} \text{J}$ b) $6.67 \times 10^{11} \text{J}$ c) $4.42 \times 10^{-18} \text{J}$ d) $4.42 \times 10^{-15} \text{J}$
- 3 The energies E_1 and E_2 of two radiations are 25 eV and 50 eV respectively. The relation between their wavelengths ie λ_1 and λ_2 will be
 a) $\frac{\lambda_1}{\lambda_2} = 1$ b) $\lambda_1 = 2\lambda_2$ c) $\lambda_1 = \sqrt{25 \times 50}\lambda_2$ d) $2\lambda_1 = \lambda_2$
- 4 Splitting of spectral lines in an electric field is called
 a) Zeeman effect b) Shielding effect c) Compton effect d) Stark effect
- 5 Based on equation $E = -2.178 \times 10^{-18} \text{J} \left(\frac{z^2}{n^2}\right)$, certain conclusions are written. Which of them is not correct?
 a) Equation can be used to calculate the change in energy when the electron changes orbit
 b) For $n = 1$, the electron has a more negative energy than it does for $n = 6$ which means that the electron is more loosely bound in the smallest allowed orbit
 c) The negative sign in equation simply means that the energy of electron bound to the nucleus is lower than it would be if the electrons were at the infinite distance from the nucleus.
 d) Larger the value of n , the larger is the orbit radius.
- 6 According to the Bohr Theory, which of the following transitions in the hydrogen atom will give rise to the least energetic photon
 a) $n = 6$ to $n = 1$ b) $n = 5$ to $n = 4$ c) $n = 5$ to $n = 3$ d) $n = 6$ to $n = 5$
- 7 Assertion : The spectrum of He^+ is expected to be similar to that of hydrogen
 Reason : He^+ is also one electron system.
 a) If both assertion and reason are true and reason is the correct explanation of assertion.
 b) If both assertion and reason are true but reason is not the correct explanation of assertion.
 c) If assertion is true but reason is false
 d) If both assertion and reason are false
- 8 Which of the following pairs of d-orbitals will have electron density along the axes ?
 a) dz^2, dxz b) dxz, dyz c) $dz^2, dx^2 - y^2$ d) $dxy, dx^2 - y^2$
- 9 Two electrons occupying the same orbital are distinguished by _____ quantum number
 a) azimuthal b) spin c) magnetic d) orbital

- 10 The electronic configuration of Eu (Z=63) Gd (Z= 64) and Tb (Z=65) are
 a) [Xe]4f⁶ 5d¹ 6s², [Xe]4f⁷ 5d¹ 6s² and [Xe]4f⁸ 5d¹ 6s²
 b) [Xe]4f⁷ 5d¹ 6s², [Xe]4f⁷ 5d¹ 6s² and [Xe]4f⁹ 6s²
 c) [Xe]4f⁷, 6s², [Xe]4f⁸ 6s² and [Xe]4f⁸ 5d¹ 6s²
 d) [Xe]4f⁶ 5d¹ 6s², [Xe]4f⁷ 5d¹ 6s² and [Xe]4f⁹ 6s²
- 11 The maximum number of electrons in a sub shell is given by the expression
 a) 2n² b) 2l + 1 c) 4l + 2 d) none of these
- 12 For d-electron, the orbital angular momentum is
 a) $\frac{\sqrt{2} h}{2\pi}$ b) $\frac{\sqrt{2h}}{2\pi}$ c) $\frac{\sqrt{2 \times 4} h}{2\pi}$ d) $\frac{\sqrt{6} h}{2\pi}$
- 13 What is the maximum numbers of electrons that can be associated with the following set of quantum numbers? n = 3, l = 1 and m = -1
 a) 4 b) 6 c) 2 d) 10
- 14 **Assertion :** Number of radial and angular nodes for 3p orbital are 1, 1 respectively.
Reason : Number of radial and angular nodes depends only on principal quantum number.
 a) both assertion and reason are true and reason is the correct explanation of assertion.
 b) both assertion and reason are true but reason is not the correct explanation of assertion.
 c) assertion is true but reason is false
 d) both assertion and reason are false
- 15 The total number of orbitals associated with the principal quantum number n = 3 is
 a) 9 b) 8 c) 5 d) 7
- 16 If n = 6, the correct sequence for filling of electrons will be,
 a) ns → (n - 2) f → (n - 1) d → np b) ns → (n - 1) d → (n - 2) f → np
 c) ns → (n - 2) f → np → (n - 1) d d) none of these are correct
- 17 Consider the following sets of quantum numbers :
 Which of the following sets of quantum number is not possible?
- | | n | L | m | S |
|-----|---|---|----|-------|
| I | 3 | 0 | 0 | + 1/2 |
| Ii | 2 | 2 | 1 | - 1/2 |
| Iii | 4 | 3 | -2 | + 1/2 |
| Iv | 1 | 0 | -1 | + 1/2 |
| V | 3 | 4 | 3 | - 1/2 |
- a) (i), (ii), (iii) and (iv)
 b) (ii), (iv) and (v)
 c) (i) and (iii)
 d) (ii), (iii) and (iv)
- 18 How many electrons in an atom with atomic number 105 can have (n + l)=8
 a) 30 b) 17 c) 15 d) unpredictable

- 19 Electron density in the yz plane of $3 dx^2 - y^2$ orbital is _____
 a) zero b) 0.50 c) 0.75 d) 0.90
- 20 If uncertainty in position and momentum are equal, then minimum uncertainty in velocity is
 a) $\frac{1}{m} \sqrt{\frac{h}{\pi}}$ b) $\sqrt{\frac{h}{\pi}}$ c) $\frac{1}{2m} \sqrt{\frac{h}{\pi}}$ d) $\frac{h}{4\pi}$

A macroscopic particle of mass 100 g and moving at a velocity of 100 cm s^{-1} will have a de Broglie wavelength

- a) $6.6 \times 10^{-29} \text{ cm}$ b) $6.6 \times 10^{-30} \text{ cm}$ c) $6.6 \times 10^{-30} \text{ cm}$ d) $6.6 \times 10^{-32} \text{ cm}$

The ratio of de Broglie wavelengths of a deuterium atom to that of an α -particle, when the velocity of the former is five times greater than that of later, is _____

- a) 4 b) 0.2 c) 2.5 d) 0.4

The energy of an electron in the 3rd orbit of hydrogen atom is $-E$. The energy of an electron in the first orbit will

- a) $-3E$ b) $-\frac{E}{3}$ c) $-\frac{E}{9}$ d) $-9E$

Which of the following does not represent the mathematical expression for the Heisenberg uncertainty principle?

- a) $\Delta x \cdot \Delta p \geq \frac{h}{4\pi}$ b) $\Delta x \cdot \Delta v \geq \frac{h}{4\pi m}$ c) $\Delta E \cdot \Delta t \geq \frac{h}{4\pi}$ d) $\Delta E \cdot \Delta x \geq \frac{h}{4\pi}$

3. PERIODIC CLASSIFICATION OF ELEMENTS

- 1 What would be the IUPAC name for an element with atomic number 222?
 a) bibibium b) bididium c) didibium d) bibibium
- 2 The electronic configuration of the elements A and B are $1s^2 2s^2 2p^6 3s^2$ and $1s^2 2s^2 2p^5$ respectively. The formula of the ionic compound that can be formed between these elements is
 a) AB b) AB_2 c) A_2B d) none of the above
- 3 The group of elements in which the differentiating electron enters the anti-penultimate shell of atoms are called _____ elements
 a) p-block b) d-block c) s-block d) f-block
- 4 In which of the following options the order of arrangement does not agree with the variation of property indicated against it?
 a) $I < Br < Cl < F$ (increasing electron gain enthalpy)
 b) $Li < Na < K < Rb$ (increasing metallic radius)
 c) $Al^{3+} < Mg^{2+} < Na^+ < F^-$ (increasing ionic size)
 d) $B < C < O < N$ (increasing first ionisation enthalpy)
- 5 Which of the following elements will have the highest electronegativity?
 a) Chlorine b) Nitrogen c) Cesium d) Fluorine

6 Various successive ionisation enthalpies (in kJ mol⁻¹) of an element are given below.

IE ₁	IE ₂	IE ₃	IE ₄	IE ₅
577.5	1,810	2,750	11,580	14,820

- a) phosphorus b) Sodium c) Aluminium d) Silicon
- 7 In the third period the first ionization potential is of the order.
- a) Na > Al > Mg > Si > P b) Na < Al < Mg < Si < P
c) Mg > Na > Si > P > Al d) Na < Al < Mg < Si < P
- 8 Identify the wrong statement.
- a) Amongst the isoelectronic species, smaller the positive charge on cation, smaller is the ionic radius
b) Amongst isoelectronic species greater the negative charge on the anion, larger is the ionic radius
c) Atomic radius of the elements increases as one moves down the first group of the periodic table
d) Atomic radius of the elements decreases as one moves across from left to right in the 2nd period of the periodic table.
- 9 Which one of the following arrangements represent the correct order of least negative to most negative electron gain enthalpy
- a) Al < O < C < Ca < F b) Al < Ca < O < C < F
c) C < F < O < Al < Ca d) Ca < Al < C < O < F
- 10 The correct order of electron gain enthalpy with negative sign of F, Cl, Br and I having atomic number 9, 17, 35 and 53 respectively is
- a) I > Br > Cl > F b) F > Cl > Br > I c) Cl > F > Br > I d) Br > I > Cl > F
- 11 Which one of the following is the least electronegative element?
- a) Bromine b) Chlorine c) Iodine d) Hydrogen
- 12 The element with positive electron gain enthalpy is
- a) Hydrogen b) Sodium c) Argon d) Fluorine
- 13 The correct order of decreasing electronegativity values among the elements X, Y, Z and A with atomic numbers 4, 8, 7 and 12 respectively
- a) Y > Z > X > A b) Z > A > Y > X c) X > Y > Z > A d) X > Y > A > Z
- 14 **Assertion:** Helium has the highest value of ionisation energy among all the elements known
Reason: Helium has the highest value of electron affinity among all the elements known
- a) Both assertion and reason are true and reason is correct explanation for the assertion
b) Both assertion and reason are true but the reason is not the correct explanation for the assertion

- c) Assertion is true and the reason is false
d) Both assertion and the reason are false
- 15 The electronic configuration of the atom having maximum difference in first and second ionisation energies is
a) $1s^2 2s^2 2p^6 3s^1$ b) $1s^2 2s^2 2p^6 3s^2$
c) $1s^2 2s^2 2p^6 3s^2 3p^6 3d^6 4s^1$ d) $1s^2 2s^2 2p^6 3s^2 3p^6 3d^1$
- 16 Which of the following is second most electronegative element?
a) Chlorine b) Fluorine c) Oxygen d) Sulphur
- 17 IE_1 and IE_2 of Mg are 179 and 348 kcal mol⁻¹ respectively. The energy required for the reaction $Mg \rightarrow Mg^{2+} + 2e^-$ is
a) +169 kcal mol⁻¹ b) -169 kcal mol⁻¹ c) +527 kcal mol⁻¹ d) -527 kcal mol⁻¹
- 18 In a given shell the order of screening effect is
a) $s > p > d > f$ b) $s > p > f > d$ c) $f > d > p > s$ d) $f > p > s > d$
- 19 Which of the following orders of ionic radii is correct?
a) $H^- > H^+ > H$ b) $Na^+ > F^- > O^{2-}$ c) $F > O^{2-} > Na^+$ d) None of these
- 20 The First ionization potential of Na, Mg and Si are 496, 737 and 786 kJ mol⁻¹ respectively. The ionization potential of Al will be closer to ____
a) 760 kJ mol⁻¹ b) 575 kJ mol⁻¹ c) 801 kJ mol⁻¹ d) 419 kJ mol⁻¹
- 21 Which one of the following is true about metallic character when we move from left to right in a period and top to bottom in a group?
a) Decreases in a period and increases along the group
b) Increases in a period and decreases in a group
c) Increases both in the period and the group
d) Decreases both in the period and in the group
- 22 How does electron affinity change when we move from left to right in a period in the periodic table?
a) Generally increases b) Generally decreases
c) Remains unchanged d) First increases and then decreases
- 23 Which of the following pairs of elements exhibit diagonal relationship?
a) Be and Mg b) Li and Mg c) Be and B d) Be and Al

UNIT – 4 HYDROGEN

- 1 Which of the following statements about hydrogen is incorrect ?
a) Hydrogen ion, H_3O^+ exists freely in solution.
b) Dihydrogen acts as a reducing agent.
c) Hydrogen has three isotopes of which tritium is the most common.
d) Hydrogen never acts as cation in ionic salts.
- 2 Which one of the following statements is incorrect with regard to ortho and para dihydrogen?

- a) They are nuclear spin isomers
 b) Ortho isomer has zero nuclear spin whereas the para isomer has one nuclear spin
 c) The para isomer is favoured at low temperatures
 d) The thermal conductivity of the para isomer is 50% greater than that of the ortho isomer.
- 3 Ionic hydrides are formed by
 a) halogens b) chalogens c) inert gases d) group one elements
- 4 Tritium nucleus contains _____
 a) $1p + 0n$ b) $2p + 1n$ c) $1p + 2n$ d) none of these
- 5 Non-stoichiometric hydrides are formed by
 a) palladium, vanadium b) carbon, nickel
 c) manganese, lithium d) nitrogen, chlorine
- 6 Assertion : Permanent hardness of water is removed by treatment with washing soda.
 Reason : Washing soda reacts with soluble calcium and magnesium chlorides and sulphates in hard water to form insoluble carbonates
 a) Both assertion and reason are true and reason is the correct explanation of assertion.
 b) Both assertion and reason are true but reason is not the correct explanation of assertion.
 c) Assertion is true but reason is false
 d) Both assertion and reason are false
- 7 If a body of a fish contains 1.2 g hydrogen in its total body mass, if all the hydrogen is replaced with deuterium then the increase in body weight of the fish will be
 a) 1.2 g b) 2.4 g c) 3.6 g d) 4.8 g
- 8 The hardness of water can be determined by volumetrically using the reagent
 a) sodium thio sulphate b) potassium permanganate
 c) hydrogen peroxide d) EDTA
- 9 The cause of permanent hardness of water is due to
 a) $\text{Ca}(\text{HCO}_3)_2$ b) $\text{Mg}(\text{HCO}_3)_2$ c) CaCl_2 d) MgCO_3

5. ALKALI AND ALKALINE EARTH METALS

- 1 For alkali metals, which one of the following trends is incorrect?
 a) Hydration energy : $\text{Li} > \text{Na} > \text{K} > \text{Rb}$
 b) Ionisation energy : $\text{Li} > \text{Na} > \text{K} > \text{Rb}$
 c) Density : $\text{Li} < \text{Na} < \text{K} < \text{Rb}$
 d) Atomic size : $\text{Li} < \text{Na} < \text{K} < \text{Rb}$
- 2 Which of the following statements is incorrect?
 a) Li^+ has minimum degree of hydration among alkali metal cations.
 b) The oxidation state of K in KO_2 is +1
 c) Sodium is used to make Na / Pb alloy

- d) MgSO_4 is readily soluble in water
- 3 Which of the following compounds will not evolve H_2 gas on reaction with alkali metals ?
 a) ethanoic acid b) ethanol c) phenol d) none of these
- 4 Which of the following has the highest tendency to give the reaction

$$\text{M}_{(g)}^+ \xrightarrow{\text{Aqueous medium}} \text{M}_{(aq)}^+$$
 a) Na b) Li c) Rb d) K
- 5 sodium is stored in _____
 a) alcohol b) water c) kerosene d) none of these
- 6 RbO_2 is _____
 a) superoxide and paramagnetic b) peroxide and diamagnetic
 c) superoxide and diamagnetic d) peroxide and paramagnetic
- 7 Find the wrong statement
 a) sodium metal is used in organic qualitative analysis
 b) sodium carbonate is soluble in water and it is used in inorganic qualitative analysis
 c) potassium carbonate can be prepared by solvay process
 d) potassium bicarbonate is acidic salt
- 8 Lithium shows diagonal relationship with _____
 a) sodium b) magnesium c) calcium d) aluminium
- 9 In case of alkali metal halides, the ionic character increases in the order
 a) $\text{MF} < \text{MCl} < \text{MBr} < \text{MI}$ b) $\text{MI} < \text{MBr} < \text{MCl} < \text{MF}$
 c) $\text{MI} < \text{MBr} < \text{MF} < \text{MCl}$ d) none of these
- 10 In which process, fused sodium hydroxide is electrolysed for extraction of sodium ?
 a) Castner's process b) Cyanide process c) Down process d) All of these
- 11 The product obtained as a result of a reaction of nitrogen with CaC_2 is
 a) $\text{Ca}(\text{CN})_3$ b) CaN_2 c) $\text{Ca}(\text{CN})_2$ d) Ca_3N_2
- 12 Which of the following has highest hydration energy _____ ?
 a) MgCl_2 b) CaCl_2 c) BaCl_2 d) SrCl_2
- 13 Match the flame colours of the alkali and alkaline earth metal salts in the Bunsen burner

P	Sodium	1	Brick red
Q	Calcium	2	Yellow
R	Barium	3	Violet
S	Strontium	4	Apple green
T	Cesium	5	Crimson red
U	Potassium	6	Blue

- a) p - 2, q - 1, r - 4, s - 5, t - 6, u - 3
 b) p - 1, q - 2, r - 4, s - 5, t - 6, u - 3
 c) p - 4, q - 1, r - 2, s - 3, t - 5, u - 6
 d) p - 6, q - 5, r - 4, s - 3, t - 1, u - 2

- 14 **Assertion :** Generally alkali and alkaline earth metals form superoxides
Reason : There is a single bond between O and O in superoxides.
 a) both assertion and reason are true and reason is the correct explanation of

assertion




- b) both assertion and reason are true but reason is not the correct explanation of assertion
- c) assertion is true but reason is false
- d) both assertion and reason are false
- 15 **Assertion** : BeSO_4 is soluble in water while BaSO_4 is not
Reason : Hydration energy decreases down the group from Be to Ba and lattice energy remains almost constant.
- a) both assertion and reason are true and reason is the correct explanation of assertion
- b) both assertion and reason are true but reason is not the correct explanation of assertion
- c) assertion is true but reason is false
- d) both assertion and reason are false
- 16 Which is the correct sequence of solubility of carbonates of alkaline earth metals?
- a) $\text{BaCO}_3 > \text{SrCO}_3 > \text{CaCO}_3 > \text{MgCO}_3$ b) $\text{MgCO}_3 > \text{CaCO}_3 > \text{SrCO}_3 > \text{BaCO}_3$
- c) $\text{CaCO}_3 > \text{BaCO}_3 > \text{SrCO}_3 > \text{MgCO}_3$ d) $\text{BaCO}_3 > \text{CaCO}_3 > \text{SrCO}_3 > \text{MgCO}_3$
- 17 In context with beryllium, which one of the following statements is incorrect?
- a) It is rendered passive by nitric acid b) It forms Be_2C
- c) Its salts are rarely hydrolysed d) Its hydride is electron deficient and polymeric
- 18 A colourless solid substance (A) on heating evolved CO_2 and also gave a white residue, soluble in water. Residue also gave CO_2 when treated with dilute HCl .
- a) Na_2CO_3 b) NaHCO_3 c) CaCO_3 d) $\text{Ca}(\text{HCO}_3)_2$
- 19 The compound (X) on heating gives a colourless gas and a residue that is dissolved in water to obtain (B). Excess of CO_2 is bubbled through aqueous solution of B, C is formed. Solid (C) on heating gives back X. (B) is
- a) CaCO_3 b) $\text{Ca}(\text{OH})_2$ c) Na_2CO_3 d) NaHCO_3
- 20 Which of the following statement is false ? (NEET - Phase - I)
- a) Ca^{2+} ions are not important in maintaining the regular beating of the heart
- b) Mg^{2+} ions are important in the green parts of the plants
- c) Mg^{2+} ions form a complex with ATP
- d) Ca^{2+} ions are important in blood clotting
- 21 When CaC_2 is heated in atmospheric nitrogen in an electric furnace the compound formed is
- a) $\text{Ca}(\text{CN})_2$ b) CaNCN c) CaC_2N_2 d) CaNC_2
- 22 Among the following the least thermally stable is
- (a) K_2CO_3 b) Na_2CO_3 (c) BaCO_3 d) Li_2CO_3

6.GASEOUS STATE

- 1 Gases deviate from ideal behavior at high pressure. Which of the following statement(s) is correct for non-ideality?

- a) at high pressure the collision between the gas molecule become enormous
 b) at high pressure the gas molecules move only in one direction
 c) at high pressure, the volume of gas become insignificant
 d) at high pressure the intermolecular interactions become significant
- 2 Rate of diffusion of a gas is
 a) directly proportional to its density
 b) directly proportional to its molecular weight
 c) directly proportional to its square root of its molecular weight
 d) inversely proportional to the square root of its molecular weight
- 3 When an ideal gas undergoes unrestrained expansion, no cooling occurs because the molecules
 a) are above inversion temperature b) exert no attractive forces on each other
 c) do work equal to the loss in kinetic energy d) collide without loss of energy
- 4 Equal weights of methane and oxygen are mixed in an empty container at 298 K. The fraction of total pressure exerted by oxygen is
 (a) $\frac{1}{3}$ (b) $\frac{1}{2}$ (c) $\frac{2}{3}$ (d) $\frac{1}{3} \times 273 \times 298$
- 5 In a closed room of 1000 m³ a perfume bottle is opened up. The room develops a smell. This is due to which property of gases?
 a) Viscosity b) Density c) Diffusion d) None
- 6 A bottle of ammonia and a bottle of HCl connected through a long tube are opened simultaneously at both ends. The white ammonium chloride ring first formed will be
 a) At the center of the tube b) Near the hydrogen chloride bottle
 c) Near the ammonia bottle d) Throughout the length of the tube
- 7 The value of universal gas constant depends upon
 a) Temperature of the gas b) Volume of the gas
 c) Number of moles of the gas d) units of Pressure and volume
- 8 The value of the gas constant R is
 a) 0.082 dm³ atm. b) 0.987 cal mol⁻¹ K⁻¹
 c) 8.3 J mol⁻¹ K⁻¹ d) 8 erg mol⁻¹ K⁻¹
- 9 Use of hot air balloon in sports at meteorological observation is an application of
 a) Boyle's law b) Newton's law c) Kelvin's law d) Brown's law
- 10 Consider the following statements
 i) Atmospheric pressure is less at the top of a mountain than at sea level
 ii) Gases are much more compressible than solids or liquids
 iii) When the atmospheric pressure increases the height of the mercury column rises
 Select the correct statement
 a) I and II b) II and III c) I and III d) I, II and III
- 11 If temperature and volume of an ideal gas is increased to twice its values, the initial

pressure P becomes

- a) 4P b) 2P c) P d) 3P
- 12 At identical temperature and pressure, the rate of diffusion of hydrogen gas is $3\sqrt{3}$ times that of a hydrocarbon having molecular formula C_nH_{2n-2} . What is the value of n ?
- a) 8 b) 4 c) 3 d) 1
- 13 Equal moles of hydrogen and oxygen gases are placed in a container, with a pin-hole through which both can escape what fraction of oxygen escapes in the time required for one-half of the hydrogen to escape.
- a) $\frac{3}{8}$ b) $\frac{1}{2}$ c) $\frac{1}{8}$ d) $\frac{1}{4}$
- 14 What is the density of N_2 gas at $227^\circ C$ and 5.00 atm pressure?
(R = 0.082 L atm K^{-1} mol $^{-1}$)
- a) 1.40 g/L b) 2.81 g/L c) 3.41 g/L d) 0.29 g/L
- 15 Which of the following diagrams correctly describes the behaviour of a fixed mass of an ideal gas ? (T is measured in K)
- a)  b)  c)  d) All of these
- 16 25g of each of the following gases are taken at $27^\circ C$ and 600 mm Hg pressure. Which of these will have the least volume?
- a) HBr b) HCl c) HF d) HI

THERMODYNAMICS

- 1 The amount of heat exchanged with the surrounding at constant temperature and pressure is given by the quantity
- a) ΔE b) ΔH c) ΔS d) ΔG
- 2 All the naturally occurring processes proceed spontaneously in a direction which leads to
- a) decrease in entropy b) increase in enthalpy
c) increase in free energy d) decrease in free energy
- 3 In an adiabatic process, which of the following is true ?
- a) $q = w$ b) $q = 0$ c) $\Delta E = q$ d) $P \Delta V = 0$
- 4 In a reversible process, the change in entropy of the universe is
- a) > 0 b) > 0 c) < 0 d) $= 0$
- 5 In an adiabatic expansion of an ideal gas
- a) $w = -\Delta U$ b) $w = \Delta U + \Delta H$ c) $\Delta u = 0$ d) $w = 0$
- 6 The intensive property among the quantities below is

- a) mass b) volume c) enthalpy d) $\frac{\text{mass}}{\text{volume}}$
- 7 An ideal gas expands from the volume of $1 \times 10^{-3} \text{ m}^3$ to $1 \times 10^{-2} \text{ m}^3$ at 300 K against a constant pressure at $1 \times 10^5 \text{ Nm}^{-2}$. The work done is
 a) -900 J b) 900 kJ c) 270 kJ d) -900 KJ
- 8 Heat of combustion is always _____
 a) positive b) negative c) zero d) either positive or negative
- 9 The heat of formation of CO and CO₂ are -26.4 kcal and -94 kcal , respectively. Heat of combustion of carbon monoxide will be
 a) $+26.4 \text{ kcal}$ b) -67.6 kcal c) -120.6 kcal d) $+52.8 \text{ kcal}$
- 10 C(diamond) \rightarrow C(graphite), $\Delta H = -ve$, this indicates that
 a) graphite is more stable than diamond b) graphite has more energy than diamond
 c) both are equally stable d) stability cannot be predicted
- 11 The enthalpies of formation of Al₂O₃ and Cr₂O₃ are -1596 kJ and -1134 kJ , respectively. ΔH for the reaction $2\text{Al} + \text{Cr}_2\text{O}_3 \rightarrow 2\text{Cr} + \text{Al}_2\text{O}_3$ is
 a) -1365 kJ b) 2730 kJ c) -2730 kJ d) -462 KJ
- 12 Which of the following is not a thermodynamic function ?
 a) internal energy b) enthalpy c) entropy d) frictional energy
- 13 If one mole of ammonia and one mole of hydrogen chloride are mixed in a closed container to form ammonium chloride gas, then
 a) $\Delta H > \Delta U$ b) $\Delta H - \Delta U = 0$ c) $\Delta H + \Delta U = 0$ d) $\Delta H < \Delta U$
- 14 Change in internal energy, when 4 kJ of work is done on the system and 1 kJ of heat is given out the system is
 a) $+1 \text{ kJ}$ b) -5 kJ c) $+3 \text{ kJ}$ d) -3 KJ
- 15 The work done by the liberated gas when 55.85 g of iron (molar mass 55.85 g mol^{-1}) reacts with hydrochloric acid in an open beaker at 250 C
 a) -2.48 kJ b) -2.22 kJ c) $+2.22 \text{ kJ}$ d) $+2.48 \text{ kJ}$
- 16 The value of ΔH for cooling 2 moles of an ideal monatomic gas from 125°C to 25°C at constant pressure will be given $C_p = \frac{5}{2}R$
 a) $-250 R$ b) $-500 R$ c) $500 R$ d) $+250 R$
- 17 Given that $\text{C}_{(g)} + \text{O}_{2(g)} \rightarrow \text{CO}_{2(g)} \Delta H^\circ = -a \text{ kJ}$;
 $2 \text{CO}_{(g)} + \text{O}_{2(g)} \rightarrow 2\text{CO}_{2(g)} \Delta H^\circ = -b \text{ kJ}$; Calculate the ΔH° for the reaction $\text{C}_{(g)} + \frac{1}{2}\text{O}_{2(g)} \rightarrow \text{CO}_{(g)}$
 a) $\frac{b+2a}{2}$ b) $2a - b$ c) $\frac{2a-b}{2}$ d) $\frac{b-2a}{2}$
- 18 When 15.68 litres of a gas mixture of methane and propane are fully combusted at 0°C and 1 atmosphere , 32 litres of oxygen at the same temperature and pressure are consumed. The amount of heat of released from this combustion in kJ is

- $(\Delta H_c(\text{CH}_4) = - 890 \text{ kJ mol}^{-1}$ and $\Delta H_c(\text{C}_3\text{H}_8) = - 2220 \text{ kJ mol}^{-1}$)
- a) $- 889 \text{ kJ}$ b) $- 1390 \text{ kJ}$ c) $- 3180 \text{ kJ}$ d) $- 653.66 \text{ KJ}$
- 19 The bond dissociation energy of methane and ethane are 360 kJ mol^{-1} and 620 kJ mol^{-1} respectively. Then, the bond dissociation energy of C-C bond is
a) 170 kJ mol^{-1} b) 50 kJ mol^{-1} c) 80 kJ mol^{-1} d) 220 kJ mol^{-1}
- 20 The correct thermodynamic conditions for the spontaneous reaction at all temperature is
a) $\Delta H < 0, \Delta S > 0$ b) $\Delta H < 0, \Delta S < 0$ c) $\Delta H > 0, \Delta S = 0$ d) $\Delta H > 0, \Delta S > 0$
- 21 The temperature of the system, decreases in an _____
a) Isothermal expansion b) Isothermal Compression
c) adiabatic expansion d) adiabatic compression
- 22 In an isothermal reversible compression of an ideal gas the sign of q, ΔS and w are respectively
a) +, -, - b) -, +, - c) +, -, + d) -, -, +
- 23 Molar heat of vapourisation of a liquid is 4.8 kJ mol^{-1} . If the entropy change is $16 \text{ J mol}^{-1} \text{ K}^{-1}$, the boiling point of the liquid is
a) 323 K b) 270 C c) 164 K d) 0.3 K
- 24 ΔS is expected to be maximum for the reaction
a) $\text{Ca}_{(s)} + \frac{1}{2} \text{O}_{2(g)} \rightarrow \text{CaO}_{(s)}$ b) $\text{C}_{(s)} + \text{O}_{2(g)} \rightarrow \text{CO}_{2(g)}$
c) $\text{N}_{2(g)} + \text{O}_{2(g)} \rightarrow 2\text{NO}_{(g)}$ d) $\text{CaCO}_{3(s)} \rightarrow \text{CaO}_{(s)} + \text{CO}_{2(g)}$
- 25 The values of ΔH and ΔS for a reaction are respectively 30 kJ mol^{-1} and $100 \text{ JK}^{-1} \text{ mol}^{-1}$. Then the temperature above which the reaction will become spontaneous is
a) 300 K b) 30 K c) 100 K d) 200 C

UNIT – 8 PHYSICAL AND CHEMICAL EQUILIBRIUM

- 1 If K_b and K_f for a reversible reactions are 0.8×10^{-5} & 1.6×10^{-4} respectively, the value of the equilibrium constant is,
a) 20 b) 0.2×10^{-1} c) 0.05 d) none of these
- 2 At a given temperature and pressure, the equilibrium constant values for the equilibria $3\text{A}_2 + \text{B}_2 + 2\text{C} \xrightleftharpoons{K_1} 2\text{A}_3\text{BC}$ and $\text{A}_3\text{BC} \xrightleftharpoons{K_2} \frac{3}{2}[\text{A}_2] + \frac{1}{2}\text{B}_2 + \text{C}$ The relation between K_1 and K_2 is
(a) $K_1 = \frac{1}{\sqrt{K_2}}$ (b) $K_2 = K_1^{-1/2}$ (c) $K_1^2 = 2K_2$ (d) $\frac{K_1}{2} = K_2$
- 3 The equilibrium constant for a reaction at room temperature is K_1 and that at 700 K is K_2 . If $K_1 > K_2$, then
a) The forward reaction is exothermic b) The forward reaction is endothermic
c) The reaction does not attain equilibrium d) The reverse reaction is exothermic

- 4 The formation of ammonia from $N_{2(g)}$ and $H_{2(g)}$ is a reversible reaction
 $N_{2(g)} + 3H_{2(g)} \rightleftharpoons 2NH_{3(g)} + \text{Heat}$ What is the effect of increase of temperature on this equilibrium reaction
- a) equilibrium is unaltered b) formation of ammonia is favoured
c) equilibrium is shifted to the left d) reaction rate does not change
- 5 Solubility of carbon dioxide gas in cold water can be increased by
- a) increase in pressure b) decrease in pressure
c) increase in volume d) none of these
- 6 Which one of the following is incorrect statement ?
- a) for a system at equilibrium, Q is always less than the equilibrium constant
b) equilibrium can be attained from either side of the reaction
c) presence of catalyst affects both the forward reaction and reverse reaction to the same extent
d) Equilibrium constant varied with temperature
- 7 K_1 and K_2 are the equilibrium constants for the reactions respectively.
- $$N_{2(g)} + O_{2(g)} \xrightleftharpoons{k_1} 2NO_{(g)} \quad 2NO_{(g)} + O_2 \xrightleftharpoons{k_2} 2NO_{(g)}$$
- What is the equilibrium constant for the reaction
- a) $\frac{1}{\sqrt{k_1 k_2}}$ b) $K_1 = K_2$ c) $\frac{1}{2 K_1 K_2}$ d) $\left(\frac{1}{K_1 K_2}\right)^{\frac{3}{2}}$
- 8 In the equilibrium, $2A_{(g)} \rightleftharpoons 2B_{(g)} + C_{2(g)}$ the equilibrium concentrations of A,B and C_2 at 400 K are $1 \times 10^{-4}M$, $2.0 \times 10^{-3}M$, $1.5 \times 10^{-4}M$ respectively. The value of K_c for the equilibrium at 400 K is
- a) 0.06 b) 0.09 c) 0.62 d) 3×10^{-2}
- 9 An equilibrium constant of 3.2×10^{-6} for a reaction means, the equilibrium is
- a) largely towards forward direction b) largely towards reverse direction
c) never established d) none of these
- 10 $\frac{K_c}{K_p}$ for the reaction, $N_{2(g)} + 3H_{2(g)} \rightleftharpoons 2NH_{3(g)}$ is
- a) $\frac{1}{RT}$ b) \sqrt{RT} c) RT d) $(RT)^2$
- 11 For the reaction $AB(g) \rightleftharpoons A(g) + B(g)$, at equilibrium, AB is 20% dissociated at a total pressure of P, The equilibrium constant K_p is related to the total pressure by the expression
- a) $P = 24 K_p$ b) $P = 8 K_p$ c) $24 P = K_p$ d) none of these
- 12 In which of the following equilibrium, K_p and K_c are not equal?
- a) $2 NO_{(g)} \rightleftharpoons N_{2(g)} + O_{2(g)}$ b) $SO_{2(g)} + NO_2 \rightleftharpoons SO_{3(g)} + NO_{(g)}$
c) $H_{2(g)} + I_{2(g)} \rightleftharpoons 2HI_{(g)}$ d) $PCl_{5(g)} \rightleftharpoons PCl_{3(g)} + Cl_{2(g)}$

- 13 If x is the fraction of PCl_5 dissociated at equilibrium in the reaction $\text{PCl}_5 \rightleftharpoons \text{PCl}_3 + \text{Cl}_2$ then starting with 0.5 mole of PCl_5 , the total number of moles of reactants and products at equilibrium is
 a) $0.5 - x$ b) $x + 0.5$ c) $2x + 0.5$ d) $x + 1$
- 14 The values of for the reactions $\text{X} \rightleftharpoons \text{Y} + \text{Z}$, $\text{A} \rightleftharpoons 2\text{B}$ are in the ratio 9 : 1 if degree of dissociation and initial concentration of X and A be equal then total pressure at equilibrium P_1 , and P_2 are in the ratio
 a) 36 : 1 b) 1 : 1 c) 3 : 1 d) 1 : 9
- 15 In the reaction, $\text{Fe}(\text{OH})_{3(s)} \rightleftharpoons \text{Fe}_{(aq)}^{3+} + 3\text{OH}_{(aq)}^-$, if the concentration of OH^- ions is decreased by $\frac{1}{4}$ times, then the equilibrium concentration of Fe^{3+} will
 a) not changed b) also decreased by $\frac{1}{4}$ times
 c) increase by 4 times d) increase by 64 times
- 16 Consider the reaction where $K_p = 0.5$ at a particular temperature $\text{PCl}_{5(g)} \rightleftharpoons \text{PCl}_{3(g)} + \text{Cl}_{2(g)}$ if the three gases are mixed in a container so that the partial pressure of each gas is initially 1atm, then which one of the following is true
 a) more PCl_3 will be produced b) more Cl_2 will be produced
 c) more PCl_5 will be produced d) none of these
- 17 Equimolar concentrations of H_2 and I_2 are heated to equilibrium in a 1litre flask. What percentage of initial concentration of H_2 has reacted at equilibrium if rate constant for both forward and reverse reactions are equal
 a) 33% b) 66% c) (33)2 % d) 16.5 %
- 18 In a chemical equilibrium, the rate constant for the forward reaction is 2.5×10^2 and the equilibrium constant is 50. The rate constant for the reverse reaction is,
 a) 11.5 b) 5 c) 2×10^2 d) 2×10^{-3}
- 19 Which of the following is not a general characteristic of equilibrium involving physical process
 a) Equilibrium is possible only in a closed system at a given temperature
 b) The opposing processes occur at the same rate and there is a dynamic but stable condition
 c) All the physical processes stop at equilibrium
 d) All measurable properties of the system remains constant
- 20 For the formation of Two moles of $\text{SO}_{3(g)}$ from SO_2 and O_2 , the equilibrium constant is K_1 . The equilibrium constant for the dissociation of one mole of SO_3 into SO_2 and O_2 is
 a) $\frac{1}{K_1}$ b) K_1^2 c) $\left(\frac{1}{K_1}\right)^{1/2}$ d) $\frac{K_2}{2}$

21 Match the equilibria with the corresponding conditions,

- i) Liquid \rightleftharpoons Vapour
 ii) Solid \rightleftharpoons Liquid
 iii) Solid \rightleftharpoons Vapour
 iv) Solute (s) \rightleftharpoons Solute (Solution)

(i)	(ii)	(iii)	(iv)
(a)	1	2	3
(b)	3	1	4
(c)	2	1	3
(d)	3	2	4

- 1) melting point
 2) Saturated solution
 3) Boiling point
 4) Sublimation point
 5) Unsaturated solution


22 Consider the following reversible reaction at equilibrium, $A + B \rightleftharpoons C$, If the concentration of the reactants A and B are doubled, then the equilibrium constant will

- a) be doubled b) become one fourth c) be halved d) remain the same

23 $[\text{Co}(\text{H}_2\text{O})_6]^{2+}(\text{aq})$ (pink) + $4\text{Cl}^-(\text{aq}) \rightleftharpoons [\text{CoCl}_4]^{2-}(\text{aq})$ (blue) + $6\text{H}_2\text{O}(\text{l})$
 In the above reaction at equilibrium, the reaction mixture is blue in colour at room temperature. On cooling this mixture, it becomes pink in colour. On the basis of this information, which one of the following is true?

- a) $\Delta H > 0$ for the forward reaction b) $\Delta H = 0$ for the reverse reaction
 c) $\Delta H < 0$ for the forward reaction
 d) Sign of the ΔH cannot be predicted based on this information.


24 24. The equilibrium constants of the following reactions are :



The equilibrium constant (K) for the reaction ; $2\text{NH}_3 + \frac{5}{2}\text{O}_2 \rightleftharpoons 2\text{NO} + 3\text{H}_2\text{O}$ will be

- (a) $\frac{K_2^3 K_3}{K_1}$ (b) $\frac{K_1 K_3^3}{K_2}$ (c) $\frac{K_2 K_3^3}{K_1}$ (d) $\frac{K_2 K_3}{K_1}$

25 A 20 litre container at 400 K contains $\text{CO}_2(\text{g})$ at pressure 0.4 atm and an excess of SrO (neglect the volume of solid SrO). The volume of the container is now decreased by moving the movable piston fitted in the container. The maximum volume of the container, when pressure of CO_2 attains its maximum value will be :



- a) 2 litre b) 5 litre c) 10 litre d) 4 litre

UNIT – 9 SOLUTIONS

- 1 The molality of a solution containing 1.8g of glucose dissolved in 250g of water is
a) 0.2 M b) 0.01 M c) 0.02 M d) 0.04 M
- 2 Which of the following concentration terms is / are independent of temperature
a) molality b) molarity c) mole fraction d) (a) and (c)
- 3 Stomach acid, a dilute solution of HCl can be neutralised by reaction with Aluminiumhydroxide $\text{Al(OH)}_3 + 3\text{HCl}_{(\text{aq})} \rightarrow \text{AlCl}_3 + 3\text{H}_2\text{O}$
How many millilitres of 0.1 M Al(OH)_3 solution are needed to neutralise 21 mL of 0.1 M HCl ?
a) 14 mL b) 7 mL c) 21 mL d) none of these
- 4 The partial pressure of nitrogen in air is 0.76atm and its Henry's law constant is 7.6×10^4 atm at 300K. What is the molefraction of nitrogen gas in the solution obtained when air is bubbled through water at 300K?
a) 1×10^{-4} b) 1×10^{-6} c) 2×10^{-5} d) 1×10^{-5}
- 5 The Henry's law constant for the solubility of Nitrogen gas in water at 350 K is 8×10^4 atm. The mole fraction of nitrogen in air is 0.5. The number of moles of Nitrogen from air dissolved in 10 moles of water at 350K and 4 atm pressure is
a) 4×10^{-4} b) 4×10^4 c) 2×10^{-2} d) 2.5×10^{-4}
6. Which one of the following is incorrect for ideal solution ?
a) $\Delta H_{\text{mix}} = 0$ b) $\Delta U_{\text{mix}} = 0$
c) $\Delta P = P_{\text{observed}} - P_{\text{Calculated by Raoult's law}} = 0$ d) $\Delta G_{\text{mix}} = 0$
- 6 Which one of the following gases has the lowest value of Henry's law constant ?
a) N_2 b) He c) CO_2 d) H_2
- 7 P_1 and P_2 are the vapour pressures of pure liquid components, 1 and 2 respectively of an ideal binary solution if x_1 represents the mole fraction of component 1, the total pressure of the solution formed by 1 and 2 will be
a) $P_1 + x_1(P_2 - P_1)$ b) $P_2 - x_1(P_2 + P_1)$ c) $P_1 - x_2(P_1 - P_2)$ d) $P_1 + x_2(P_1 - P_2)$
- 8 Osmotic pressure (p) of a solution is given by the relation
a) $p = nRT$ b) $pV = nRT$ c) $pRT = n$ d) none of these
- 9 Which one of the following binary liquid mixtures exhibits positive deviation from Raoult's law ?
a) Acetone + chloroform b) Water + nitric acid
c) HCl + water d) ethanol + water
- 10 The Henry's law constants for two gases A and B are x and y respectively. The ratio of mole fractions of A to B is 0.2. The ratio of mole fraction of B and A dissolved in water will be
a) $\frac{2x}{y}$ b) $\frac{y}{0.2x}$ c) $\frac{0.2x}{y}$ d) $\frac{5x}{y}$

- 11 At 100°C the vapour pressure of a solution containing 6.5g a solute in 100g water is 732mm. If $K_b = 0.52$, the boiling point of this solution will be
 a) 102°C b) 100°C c) 101°C d) 100.52°C
- 12 According to Raoult's law, the relative lowering of vapour pressure for a solution is equal to
 a) mole fraction of solvent b) mole fraction of solute
 c) number of moles of solute d) number of moles of solvent
- 13 At same temperature, which pair of the following solutions are isotonic ?
 a) 0.2 M BaCl_2 and 0.2M urea b) 0.1 M glucose and 0.2 M urea
 c) 0.1 M NaCl and 0.1 M K_2SO_4 d) 0.1 M $\text{Ba}(\text{NO}_3)_2$ and 0.1 M Na_2SO_4
- 14 The empirical formula of a nonelectrolyte(X) is CH_2O . A solution containing six gram of X exerts the same osmotic pressure as that of 0.025M glucose solution at the same temperature. The molecular formula of X is
 a) $\text{C}_2\text{H}_4\text{O}_2$ b) $\text{C}_8\text{H}_{16}\text{O}_8$ c) $\text{C}_4\text{H}_8\text{O}_4$ d) CH_2O
- 15 The K_H for the solution of oxygen dissolved in water is 4×10^4 atm at a given temperature. If the partial pressure of oxygen in air is 0.4 atm, the mole fraction of oxygen in solution is
 a) 4.6×10^3 b) 1.6×10^4 c) 1×10^{-5} d) 1×10^5
- 16 Normality of 1.25M sulphuric acid is
 a) 1.25 N b) 3.75 N c) 2.5 N d) 2.25 N
- 17 Two liquids X and Y on mixing gives a warm solution. The solution is
 a) ideal
 b) non-ideal and shows positive deviation from Raoult's law
 c) ideal and shows negative deviation from Raoult's Law
 d) non-ideal and shows negative deviation from Raoult's Law
- 18 The relative lowering of vapour pressure of a sugar solution in water is 3.5×10^{-3} . The mole fraction of water in that solution is
 a) 0.0035 b) 0.35 c) 0.0035 / 18 d) 0.9965
- 19 The mass of a non-volatile solute (molar mass 80 g mol^{-1}) which should be dissolved in 92g of toluene to reduce its vapour pressure to 90%
 a) 10g b) 20g c) 9.2 g d) 8.89g
- 20 For a solution, the plot of osmotic pressure (p) versus the concentration (c in mol L^{-1}) gives a straight line with slope $310R$ where 'R' is the gas constant. The temperature at which osmotic pressure measured is
 a) $310 \times 0.082 \text{ K}$ b) 310°C c) 37°C d) $\frac{310}{0.082} \text{ K}$
- 21 200ml of an aqueous solution of a protein contains 1.26g of protein. At 300K, the osmotic pressure of this solution is found to be 2.52×10^{-3} bar. The molar mass of

protein will be ($R = 0.083 \text{ L bar mol}^{-1}\text{K}^{-1}$)

- a) $62.22 \text{ Kg mol}^{-1}$ b) 12444g mol^{-1} c) 300g mol^{-1} d) none of these
- 22 What is the molality of a 10% W/W aqueous sodium hydroxide solution ?
a) 2.778 b) 2.5 c) 10 d) 0.4
- 23 Which of the following aqueous solutions has the highest boiling point ?
a) 0.1M KNO_3 b) 0.1 M Na_3PO_4 c) 0.1 M BaCl_2 d) 0.1 M K_2SO_4
- 24 The freezing point depression constant for water is $1.86^\circ \text{K kg mol}^{-1}$. If 5g Na_2SO_4 is dissolved in 45g water, the depression in freezing point is 3.64°C . The Vant Hoff factor for Na_2SO_4 is
a) 2.50 b) 2.63 c) 3.64 d) 5.50
- 25 Equimolal aqueous solutions of NaCl and KCl are prepared. If the freezing point of NaCl is -2°C , the freezing point of KCl solution is expected to be
a) -2°C b) -4°C c) -1°C d) 0°C
- 26 **Assertion :** An ideal solution obeys Raoult's Law
Reason : In an ideal solution, solvent-solvent as well as solute-solute interactions are similar to solute-solvent interactions.
a) both assertion and reason are true and reason is the correct explanation of assertion
b) both assertion and reason are true but reason is not the correct explanation of assertion
c) assertion is true but reason is false
d) both assertion and reason are false

UNIT – 10 CHEMICAL BONDING





- 1 In which of the following Compounds does the central atom obey the octet rule?
a) XeF_4 b) AlCl_3 c) SF_6 d) SCl_2
- 2 In the molecule $\text{O}_A = \text{C} = \text{O}_B$, the formal charge on O_A , C and O_B are respectively.
a) -1, 0, + 1 b) +1, 0, -1 c) -2, 0, +2 d) 0, 0, 0
- 3 Which of the following is electron deficient?
a) PH_3 b) $(\text{CH}_3)_2$ c) BH_3 d) NH_3
- 4 Which of the following molecules contain no π bond?
a) SO_2 b) NO_2 c) CO_2 d) H_2O
- 5 The ratio of number of sigma (σ) and pi (π) bonds in 2-butynal is
a) 8/3 b) 5/3 c) 8/2 d) 9/2
- 6 Which one of the following is the likely bond angles of sulphur tetrafluoride molecule?
a) $120^\circ, 80^\circ$ b) $109^\circ.28'$ c) 90° d) $89^\circ, 117^\circ$

- 7 Assertion: Oxygen molecule is paramagnetic.
Reason :It has two unpaired electron in its bonding molecular orbital
- both assertion and reason are true and reason is the correct explanation of assertion
 - both assertion and reason are true but reason is not the correct explanation of assertion
 - assertion is true but reason is false
 - Both assertion and reason are false
- 8 According to Valence bond theory, a bond between two atoms is formed when
- fully filled atomic orbitals overlap
 - half filled atomic orbitals overlap
 - non- bonding atomic orbitals overlap
 - empty atomic orbitals overlap
- 9 In ClF_3 , NF_3 and BF_3 molecules the chlorine, nitrogen and boron atoms are
- sp^3 hybridised
 - sp^3 , sp^3 and sp^2 respectively
 - sp^2 hybridised
 - sp^3 d, sp^3 and sp^2 hybridised respectively
- 10 When one s and three p orbitals hybridise,
- four equivalent orbitals at 90° to each other will be formed
 - four equivalent orbitals at $109^\circ 28'$ to each other will be formed.
 - four equivalent orbitals, that are lying the same plane will be formed
 - none of these
- 11 Which of these represents the correct order of their increasing bond order.
- $\text{C}_2 < \text{C}_2^{2-} < \text{O}_2^{2-} < \text{O}_2$
 - $\text{C}_2^{2-} < \text{C}_2^+ < \text{O}_2 < \text{O}_2^{2-}$
 - $\text{O}_2^{2-} < \text{O}_2 < \text{C}_2^{2-} < \text{C}_2^+$
 - $\text{O}_2^{2-} < \text{C}_2^+ < \text{O}_2 < \text{C}_2^{2-}$
- 12 Hybridisation of central atom in PCl_5 involves the mixing of orbitals.
- s, p_x , p_y , $d_{x^2-y^2}$
 - s, p_x , p_y , p_{xy} , $d_{x^2-y^2}$
 - s, p_x , p_y , p_z , $d_{x^2-y^2}$
 - s, p_x , p_y , d_{xy} , $d_{x^2-y^2}$
- 13 The correct order of O-O bond length in hydrogen peroxide, ozone and oxygen is
- $\text{H}_2\text{O}_2 > \text{O}_3 > \text{O}_2$
 - $\text{O}_2 > \text{O}_3 > \text{H}_2\text{O}_2$
 - $\text{O}_2 > \text{H}_2\text{O}_2 > \text{O}_3$
 - $\text{O}_3 > \text{O}_2 > \text{H}_2\text{O}_2$
- 14 Which one of the following is diamagnetic.?
- O_2
 - O_2^-
 - O_2^+
 - None of these
- 15 Bond order of a species is 2.5 and the number of electrons in its bonding molecular orbital is found to be 8. The no. of electrons in its antibonding molecular orbital is
- three
 - four
 - Zero
 - can not be calculated from the given information
- 16 Shape and hybridisation of IF_5 are
- Trigonal bipyramidal, Sp^3d^2
 - Trigonal bipyramidal, Sp^3d
 - Square pyramidal, Sp^3d^2
 - Octahedral, Sp^3d^2
- Pick out the incorrect statement from the following
- Sp^3 hybrid orbitals are equivalent and are at an angle of $109^\circ 28'$ with each other
 - dsp^2 hybrid orbitals are equivalent and bond angle between any two of them is 90°

- c) All five sp^3d hybrid orbitals are not equivalent out of these five sp^3d hybrid orbitals, three are at an angle of 120° , remaining two are perpendicular to the plane containing the other three
- d) none of these
- 17 The molecules having same hybridisation, shape and number of lone pairs of electrons are
 a) SeF_4, XeO_2F_2 b) SF_4, XeF_2 c) $XeOF_4, TeF_4$ d) $SeCl_4, XeF_4$
- 18 In which of the following molecules /ions BF_3, NO_2^-, H_2O the central atom is sp^2 hybridised?
 a) NH_2^- and H_2O b) NO_2^- and H_2O c) BF_3 and NO_2^- d) BF_3 and NH_2^-
- 19 Some of the following properties of two species, NO_3^- and H_3O^+ are described below. which one of them is correct?
 a) dissimilar in hybridisation for the central atom with different structure.
 b) isostructural with same hybridisation for the central atom.
 c) different hybridisation for the central atom with same structure
 d) none of these
- 20 The types of hybridisation on the five carbon atom from right to left in the, 2,3 pentadiene.
 a) $sp^3, sp^2, sp, sp^2, sp^3$ b) sp^3, sp, sp, sp, sp^3
 c) $sp^2, sp, sp^2, sp^2, sp^3$ d) $sp^3, sp^3, sp^2, sp^3, sp^3$
- 21 XeF_2 is isostructural with
 a) $SbCl_2$ b) $BaCl_2$ c) TeF_2 d) ICl_2^-
- 22 The percentage of s-character of the hybrid orbitals in methane, ethane, ethene and ethyne are respectively
 a) 25, 25, 33.3, 50 b) 50, 50, 33.3, 25 c) 50, 25, 33.3, 50 d) 50, 25, 25, 50
- 23 Of the following molecules, which have shape similar to carbon dioxide?
 a) $SnCl_2$ b) NO_2 c) C_2H_2 d) All of these.
- 24 According to VSEPR theory, the repulsion between different parts of electrons obey the order.
 a) $l.p - l.p > b.p - b.p > l.p - b.p$ b) $b.p - b.p > b.p - l.p > l.p - b.p$
 c) $l.p - l.p > b.p - l.p > b.p - b.p$ d) $b.p - b.p > l.p - l.p > b.p - l.p$
- 25 Shape of ClF_3 is
 a) Planar triangular b) Pyramidal c) 'T' Shaped d) none of these
- 26 Non- Zero dipole moment is shown by
 a) CO_2 b) p-dichlorobenzene c) carbon tetrachloride d) water
- 27 Which of the following conditions is not correct for resonating structures?
 a) the contributing structure must have the same number of unpaired electrons
 b) the contributing structures should have similar energies

- c) the resonance hybrid should have higher energy than any of the contributing structure.
- d) none of these
- 28 Among the following, the compound that contains, ionic, covalent and Coordinate linkage is
- a) NH_4Cl b) NH_3 c) NaCl d) none of these
- 29 CaO and NaCl have the same crystal structure and approximately the same radii. If U is the lattice energy of NaCl , the approximate lattice energy of CaO is
- a) U b) $2U$ c) $U/2$ d) $4U$

UNIT – 11 FUNDAMENTALS OF ORGANIC CHEMISTRY

- 1 Select the molecule which has only one π bond.
- a) $\text{CH}_3 - \text{CH} = \text{CH} - \text{CH}_3$ b) $\text{CH}_3 - \text{CH} = \text{CH} - \text{CHO}$
- c) $\text{CH}_3 - \text{CH} = \text{CH} - \text{COOH}$ d) All of these
- 2 In the hydrocarbon the state of hybridisation of carbon 1, 2, 3, 4 and 7 are in the following sequence. $\text{CH}_3 - \text{CH}_2 - \text{CH} = \text{CH} - \text{CH}_2 - \text{C} \equiv \text{CH}$
- a) sp, sp, sp^3, sp^2, sp^3 b) $sp^2, sp, sp^3, sp^2, sp^3$
- c) sp, sp, sp^2, sp, sp^3 d) none of these
- 3 The general formula for alkadiene is
- a) C_nH_{2n} b) $\text{C}_n\text{H}_{2n-1}$ c) $\text{C}_n\text{H}_{2n-2}$ d) C_nH_{n-1}
- 4 Structure of the compound whose IUPAC name is 5,6 -dimethylhept -2-ene is
- a)  b)  c)  d) None of these
- 5 The IUPAC name of the Compound is
- 
- a) 2,3 - Dimethylheptane
- b) 3- Methyl -4- ethyloctane
- c) 5-ethyl -6-methyloctane
- d) 4-Ethyl -3 - methyloctane.
- 6 Which one of the following names does not fit a real name?
- a) 3 - Methyl -3-hexanone b) 4-Methyl -3- hexanone
- c) 3- Methyl -3- hexanol d) 2- Methyl cyclohexanone.
- 7 The IUPAC name of the compound $\text{CH}_3 - \text{CH} = \text{CH} - \text{C} \equiv \text{CH}$ is
- a) Pent - 4 - yn-2-ene b) Pent -3-en-1-yne
- c) pent - 2- en - 4 - yne d) Pent - 1 - yn -3 -ene

- 8
- $$\begin{array}{c} \text{H} \quad \text{C}_4\text{H}_9 \\ | \quad | \\ \text{H}_3\text{C}-\text{C}-\text{C}-\text{CH}_3 \\ | \quad | \\ \text{C}_2\text{H}_5 \quad \text{CH}_3 \end{array}$$
- IUPAC name of is
- 9 IUPAC name of is
- $$\begin{array}{c} \text{H} \\ | \\ \text{H}_3\text{C}-\text{C}-\text{CH}=\text{C}(\text{CH}_3)_2 \\ | \\ \text{C}_2\text{H}_5 \end{array}$$
- 10 The IUPAC name of the compound is
- $$\begin{array}{c} \text{H}_3\text{C}-\text{CH}=\text{C}-\text{CH}_2\text{CH}_3 \\ | \\ \text{CH}_2\text{CH}_2\text{CH}_3 \end{array}$$
- 11 The IUPAC name of the compound is
- $$\begin{array}{c} \text{H}_3\text{C}-\text{CH}-\text{COOH} \\ | \\ \text{OH} \end{array}$$
- 12 The IUPAC name of the compound is
- $$\begin{array}{c} \text{H}_3\text{C} \\ \diagdown \\ \text{CH}-\text{CH}-\text{COOH} \\ \diagup \quad | \\ \text{Br} \quad \text{CH}_3 \end{array}$$
- 13 The structure of isobutyl group in an organic compound is
- a) $\text{H}_3\text{C}-\text{CH}_2-\text{CH}_2-\text{CH}_2-$
- b) $\begin{array}{c} \text{CH}_3 \\ | \\ \text{H}_3\text{C}-\text{C}- \\ | \\ \text{CH}_3 \end{array}$
- c) $\begin{array}{c} \text{H}_3\text{C}-\text{CH}-\text{CH}_2- \\ | \\ \text{CH}_3 \end{array}$
- d) $\begin{array}{c} \text{H}_3\text{C}-\text{CH}-\text{CH}_2-\text{CH}_3 \\ | \end{array}$
- 14 The number of stereoisomers of 1, 2 – dihydroxycyclopentane
- a) 1 b) 2 c) 3 d) 4
- 15 Which of the following is optically active?
- a) 3 – Chloropentane b) 2 Chloro propane
- c) Meso – tartaric acid d) Glucose
- 16 The isomer of ethanol is
- a) acetaldehyde b) dimethylether c) acetone d) methyl carbinol
- 17 How many cyclic and acyclic isomers are possible for the molecular formula $\text{C}_3\text{H}_6\text{O}$?
- a) 4 b) 5 c) 9 d) 10
- 18 Which one of the following shows functional isomerism?
- a) ethylene b) Propane c) ethanol d) CH_2Cl_2
- a) 3,4,4 – Trimethylheptane
b) 2 – Ethyl –3, 3– dimethyl heptane
c) 3, 4,4 – Trimethyloctane
d) 2 – Butyl –2 –methyl –3–ethyl-butane.
- a) 2,4,4 – Trimethylpent –2-ene
b) 2,4,4 – Trimethylpent –3-ene
c) 2,2,4 – Trimethylpent –3-ene
d) 2,2,4 – Trimethylpent –2-ene
- a) 3 – Ethyl –2– hexane
b) 3 – Propyl –3– hexene
c) 4 – Ethyl – 4 – hexene
d) 3 – Propyl –2-hexene
- a) 2 – Hydroxypropionic acid
b) 2 – HydroxyPropanoic acid
c) Propan – 2– ol –1 – oic acid
d) 1 – Carboxyethanol.
- a) 2 – Bromo -3 – methyl butanoic acid
b) 2 - methyl - 3- bromobutanoic acid
c) 3 - Bromo - 2 - methylbutanoic acid
d) 3 - Bromo -2,3-dimethyl propanoic acid.

- 19 $\text{H}_2\overset{\text{O}}{\text{C}}-\overset{\text{O}}{\parallel}{\text{C}}-\text{CH}_3$ and $\text{H}_3\text{C}-\overset{\text{O}}{\parallel}{\text{C}}=\text{CH}_2$
 a) resonating structure b) tautomers c) Optical isomers d) Conformers

- 20 $\text{H}_3\text{C}-\overset{\text{O}}{\parallel}{\text{C}}=\text{CH}-\text{COOH}$
 $\quad \quad \quad |$
 $\quad \quad \quad \text{COOC}_2\text{H}_5$

Assertion: 3- carbethoxy -2- butenoic acid.

Reason: The principal functional group gets lowest number followed by double bond (or) triple bond.

- a) both the assertion and reason are true and the reason is the correct explanation of assertion.
 b) both assertion and reason are true and the reason is not the correct explanation of assertion.
 c) assertion is true but reason is false
 d) both the assertion and reason are false.


UNIT – 12 BASIC CONCEPT OF ORGANIC REACTIONS

- 1 What is the hybridisation state of benzyl carbonium ion?
 (a) sp^2 (b) spd^2 (c) sp^3 (d) sp^2d
- 2 Decreasing order of nucleophilicity is
 (a) $\text{OH}^- > -\text{NH}_2 > -\text{OCH}_3 > \text{RNH}_2$ (b) $-\text{NH}_2 > \text{OH}^- > -\text{OCH}_3 > \text{RNH}_2$
 (c) $-\text{NH}_2 > -\text{OCH}_3 > \text{OH}^- > \text{RNH}_2$ (d) $-\text{OCH}_3 > -\text{NH}_2 > \text{OH}^- > \text{RNH}_2$
- 3 Which of the following species is not electrophilic in nature?
 (a) Cl^+ (b) BH_3 (c) H_3O^+ (d) $\overset{+}{\text{N}}\text{O}_2$
- 4 Homolytic fission of covalent bond leads to the formation of
 (a) electrophile (b) nucleophile (c) Carbo cation (d) free radical
- 5 Hyper Conjugation is also known as
 (a) no bond resonance (b) Baker - nathan effect
 (c) both (a)and (b) (d) none of these
- 6 Which of the group has highest +I effect?
 (a) $\text{CH}_3 -$ (b) $\text{CH}_3 - \text{CH}_2 -$ (c) $(\text{CH}_3)_2 - \text{CH} -$ (d) $(\text{CH}_3)_3 - \text{C} -$
- 7 Which of the following species does not exert a resonance effect?
 (a) $\text{C}_6\text{H}_5\text{OH}$ (b) $\text{C}_6\text{H}_5\text{Cl}$ (c) $\text{C}_6\text{H}_5\text{NH}_2$ (d) $\overset{+}{\text{C}}_6\text{H}_5\text{NH}_3$
- 8 -I effect is shown by
 (a) $-\text{Cl}$ (b) $-\text{Br}$ (c) both (a) and (b) (d) $-\text{CH}_3$
- 9 Which of the following carbocation will be most stable?
 (a) Ph_3C^+ - (b) $\text{CH}_3 - \overset{+}{\text{C}}\text{H}_2$ (c) $(\text{CH}_3)_2 - \overset{+}{\text{C}}\text{H}$ (d) $\text{CH}_2 = \text{CH} - \overset{+}{\text{C}}\text{H}_2$

- 10 **Assertion:** Tertiary Carbocations are generally formed more easily than primary Carbocations ions.
Reason: Hyper conjugation as well as inductive effect due to additional alkyl group stabilize tertiary carbonium ions.
- a) both assertion and reason are true and reason is the correct explanation of assertion.
 b) both assertion and reason are true but reason is not the correct explanation of assertion.
 c) Assertion is true but reason is false
 d) Both assertion and reason are false
- 11 Heterolytic fission of C-Br bond results in the formation of
 (a) free radical (b) Carbanion (c) Carbocation (d) Carbanion and Carbocation
- 12 Which of the following represent a set of nucleophiles?
 (a) $\text{BF}_3, \text{H}_2\text{O}, \text{NH}_2^-$ (b) $\text{AlCl}_3, \text{BF}_3, \text{NH}_3$
 (c) $\text{CN}^-, \text{RCH}_2^-, \text{ROH}$ (d) $\text{H}^+, \text{RNH}_3^+, : \text{CCl}_2$
- 13 Which of the following species does not acts as a nucleophile?
 (a) ROH (b) ROR (c) PCl_3 (d) BF_3
- 14 The geometrical shape of carbocation is
 (a) Linear (b) tetrahedral (c) Planar (d) Pyramidal

UNIT – 13 HYDROCARBON

- 1 $\text{C}_2\text{H}_5\text{Br} + 2\text{Na} \xrightarrow{\text{dry ether}} \text{C}_4\text{H}_{10} + 2\text{NaBr}$ The above reaction is an example of which of the following
 a) Reimer Tiemann reaction (b) Wurtz reaction
 c) Aldol condensation (d) Hoffmann reaction
- 2 An alkyl bromide (A) reacts with sodium in ether to form 4, 5– diethyloctane, the compound (A) is
 a) $\text{CH}_3(\text{CH}_2)_3\text{Br}$ (b) $\text{CH}_3(\text{CH}_2)_5\text{Br}$
 c) $\text{CH}_3(\text{CH}_2)_3\text{CH}(\text{Br})\text{CH}_3$ (d) $\text{CH}_3 - (\text{CH}_2)_2 - \text{CH}(\text{Br}) - \text{CH}_2\text{CH}_3$
- 3 The C – H bond and C – C bond in ethane are formed by which of the following types of overlap
 a) $\text{sp}^3 - \text{s}$ and $\text{sp}^3 - \text{sp}^3$ (b) $\text{sp}^2 - \text{s}$ and $\text{sp}^2 - \text{sp}^2$
 c) $\text{sp} - \text{sp}$ and $\text{sp} - \text{sp}$ (d) $\text{p} - \text{s}$ and $\text{p} - \text{p}$



5 Which of the following is optically active

- a) 2 – methyl pentane b) citric acid c) Glycerol d) none of these

6 The compounds formed at anode in the electrolysis of an aqueous solution of potassium acetate are

- a) CH₄ and H₂ b) CH₄ and CO₂ c) C₂H₆ and CO₂ d) C₂H₄ and Cl₂

7 The general formula for cyclo alkanes

- a) C_nH_n b) C_nH_{2n} c) C_nH_{2n-2} d) C_nH_{2n+2}

8 The compound that will react most readily with gaseous bromine has the formula

- a) C₃H₆ b) C₂H₂ c) C₄H₁₀ d) C₂H₄

9 Which of the following compounds shall not produce propene by reaction with HBr followed by elimination (or) only direct elimination reaction

- a) b) CH₃-CH₂-CH₂-OH c) H₂C = C = O d) CH₃-CH₂-CH₂Br

10 Which among the following alkenes on reductive ozonolysis produces only propanone ?

- a) 2 – Methyl propene b) 2 – Methyl but – 2 - ene
c) 2, 3 – Dimethyl but – 1 – ene d) 2, 3 – Dimethyl but – 2 – ene

11 The major product formed when 2 – bromo – 2 – methyl butane is refluxed with ethanolic KOH is


- a) 2 – methylbut – 2 – ene b) 2 – methyl butan – 1 – ol
c) 2 – methyl but – 1 – ene d) 2 – methyl butan – 2 – ol

12 Major product of the below mentioned reaction is,



- a) 2-chloro – 1 – iodo – 2 – methyl propane b) 1-chloro – 2 – iodo – 2 – methylpropane
c) 1,2 – dichloro – 2 – methyl propane d) 1, 2 – diiodo – 2 – methyl propane

13 The IUPAC name of the following compound is



- a) trans-2-chloro-3-iodo – 2 – pentane
b) cis-3 – iodo – 4 – chloro – 3 – pentane
c) trans-3-iodo-4-chloro – 3 – pentene
d) cis-2 – chloro – 3 – iodo – 2 –






pentene

14 Cis – 2 – butene and trans – 2 – butene are

- a) conformational isomers
c) configurational isomers




- b) structural isomers
d) optical isomers

15 Identify

16 Consider the nitration of benzene using mixed con H_2SO_4 and HNO_3 if a large quantity of KHSO_4 is added to the mixture, the rate of nitration will be


- a) unchanged b) doubled c) faster d) slower

17 In which of the following molecules, all atoms are co-planar




d) both (a) and (b)





18



c) both (a) and (b)



19 Which one of the following is nonaromatic ?




20 Which of the following compounds will not undergo Friedal – crafts reaction easily?

- a) Nitro benzene b) Toluene c) Cumene d) Xylene
- 21 Some meta-directing substituents in aromatic substitution are given. Which one is most deactivating ?
 a) - COOH b) - NO₂ c) - C ≡ N d) - SO₃H
- 22 Which of the following can be used as the halide component for Friedel-Crafts reaction ?
 a) Chloro benzene b) Bromo benzene
 c) chloroethene d) isopropyl chloride
- 23 An alkane is obtained by decarboxylation of sodium propionate. Same alkane can be prepared by
 a) Catalytic hydrogenation of propene b) action of sodium metal on iodomethane
 c) reduction of 1 - chloro propane d) reduction of bromomethane
- 24 Which of the following is aliphatic saturated hydrocarbon
 a) C₈H₁₈ b) C₉H₁₈ c) C₈H₁₄ d) All of these
- 25 Identify the compound 'Z' in the following reaction


$$\text{C}_6\text{H}_5\text{O} \xrightarrow[623\text{ K}]{\text{Al}_2\text{O}_3} \text{X} \xrightarrow{\text{O}_3} \text{Y} \xrightarrow[\Delta]{\text{Zn}/\text{H}_2\text{O}} \text{Z}$$
 a) Formaldehyde b) Acetaldehyde c) Formic acid d) none of these
- 26 Peroxide effect (Kharasch effect) can be studied in case of
 a) Oct - 4 - ene b) hex - 3 - ene c) pent - 1 - ene d) but - 2 - ene
- 27 2 - butyne on chlorination gives
 a) 1 - chloro butane b) 1, 2 - dichloro butane
 c) 1, 1, 2, 2 - tetrachlorobutane d) 2, 2, 3, 3 - tetra chloro butane

UNIT - 14 HALOALKANES AND HALOARENES


- 1 The IUPAC name of
- 
- a) 2-Bromo pent - 3 - ene
 b) 4-Bromo pent - 2 - ene
 c) 2-Bromo pent - 4 - ene
 d) 4-Bromo pent - 1 - ene
- 2 Of the following compounds, which has the highest boiling point?
 a) n-Butyl chloride b) Isobutyl chloride
 c) t-Butyl chloride d) n-propyl chloride
- 3 Arrange the following compounds in increasing order of their density
 A) CCl₄ B) CHCl₃ C) CH₂Cl₂ D) CH₃Cl
 a) D < C < B < A b) C > B > A > D c) A < B < C < D d) C > A > B > D

- 4 With respect to the position of -Cl in the compound $\text{CH}_3-\text{CH}=\text{CH}-\text{CH}_2-\text{Cl}$, it is classified as
 a) Vinyl b) Allyl c) Secondary d) Aralkyl
- 5 What should be the correct IUPAC name of diethyl chloromethane?
 a) 3 - Chloro pentane b) 1-Chloropentane
 c) 1-Chloro-1, 1, diethyl methane d) 1 -Chloro-1-ethyl propane
- 6 C -X bond is strongest in
 a) Chloromethane b) Iodomethane c) Bromomethane d) Fluoromethane


7 In the reaction X is



8 Which of the following compounds will give racemic mixture on nucleophilic substitution by OH^- ion?



9 The treatment of ethyl formate with excess of RMgX gives



10 Benzene reacts with Cl_2 in the presence of FeCl_3 and in absence of sunlight to form





- a) Chlorobenzene b) Benzyl chloride
 c) Benzal chloride d) Benzene hexachloride

11 **Assertion:** In mono haloarenes, electrophilic substitution occurs at ortho and para positions.

Reason: Halogen atom is a ring deactivator

- a) If both assertion and reason are true and reason is the correct explanation of assertion.

- b) If both assertion and reason are true but reason is not the correct explanation of assertion.
 c) If assertion is true but reason is false.
 d) If both assertion and reason are false.
- 12 Consider the reaction,

$$\text{CH}_3\text{CH}_2\text{CH}_2\text{Br} + \text{NaCN} \rightarrow \text{CH}_3\text{CH}_2\text{CH}_2\text{CN} + \text{NaBr}$$
 This reaction will be the fastest in
 a) ethanol b) methanol c) DMF (N,N'-dimethyl formamide) d) water
- 13 The most easily hydrolysed molecule under $\text{S}_{\text{N}}1$ condition is
 a) allyl chloride b) ethyl chloride c) isopropylchloride d) benzyl chloride
- 14 The carbocation formed in $\text{S}_{\text{N}}1$ reaction of alkyl halide in the slow step is
 a) sp^3 hybridised b) sp^2 hybridised c) sp hybridised d) none of these
- 15 The major products obtained when chlorobenzene is nitrated with HNO_3 and H_2SO_4
 a) 1-chloro-4-nitrobenzene b) 1-chloro-2-nitrobenzene
 c) 1-chloro-3-nitrobenzene d) 1-chloro-1-nitrobenzene
- 16 Which one of the following is most reactive towards nucleophilic substitution reaction ?
- a)  b)  c)  d) 
- 17 The raw material for Rasching process
 a) chloro benzene b) phenol c) benzene d) anisole
- 18 acetone $\xrightarrow[\text{ii) H}_2\text{O} / \text{H}^+]{\text{i) CH}_3\text{MgI}}$ X, X is
 a) 2-propanol b) 2-methyl-2-propanol c) 1-propanol d) acetone
- 19 Silverpropionate when refluxed with Bromine in carbon tetrachloride gives
 a) propionic acid b) chloro ethane c) bromo ethane d) chloro propane

NGK CHEMISTRY