

Sun Tuition Center

poon thotta pathai hindu mission hospital opposite
villupuram cell - 9629216361

Life is a good circle, you choose the best radius..

10th Standard – Maths Formulas

SETS AND FUNCTIONS

1. Power set $n[P(A)] = 2^{n(A)}$

2. Symmetric difference

i) $X \Delta Y = (X \setminus Y) \cup (Y \setminus X)$
ii) $X \Delta Y = (X \cup Y) \setminus (X \cap Y)$

3. Commutative property

a) $A \cup B = B \cup A$
b) $A \cap B = B \cap A$

4. Associative property

a) $A \cup (B \cup C) = (A \cup B) \cup C$
b) $A \cap (B \cap C) = (A \cap B) \cap C$

5. Distributive property

a) $A \cap (B \cup C) = (A \cap B) \cup (A \cap C)$
b) $A \cup (B \cap C) = (A \cup B) \cap (A \cup C)$

6. De – Morgan's laws of set difference

a) $A \setminus (B \cup C) = (A \setminus B) \cap (A \setminus C)$
b) $A \setminus (B \cap C) = (A \setminus B) \cup (A \setminus C)$

7. De – Morgan's laws

a) $(A \cup B)' = A' \cap B'$
b) $(A \cap B)' = A' \cup B'$

8. Cardinality of sets

a) $n(A \cup B) = n(A) + n(B) - n(A \cap B)$
b) $n(A \cup B \cup C) = n(A) + n(B) + n(C) - n(A \cap B) - n(B \cap C) - n(A \cap C) + n(A \cap B \cap C)$

9. One – one function

Every different element of A has a different image in B .

10. Onto function

Every element in B has a pre – image in A .

SEQUENCES AND SERIES

I) Arithmetic progression

1. General form $a, a+d, a+2d, a+3d, \dots$
2. General term (or) $n^{\text{th}} \text{ term}$ $T_n = a + (n-1)d$
3. Common difference $d = t_2 - t_1 = \dots = t_n - t_{n-1}$
4. Number of terms in an A.P $n = \frac{l-a}{d} + 1$

5. Sum of n terms of an A.P

a) $S_n = \frac{n}{2}[2a + (n-1)d]$
b) $S_n = \frac{n}{2}[a + l]$

6. If 3 terms in A.P are $a-d, a, a+d$

II) Geometric progression

7. General form $a, ar, ar^2, ar^3, \dots, ar^n$.
8. General term (or) $n^{\text{th}} \text{ term}$ $T_n = ar^{n-1}$.
9. Common ratio $r = \frac{t_2}{t_1} = \frac{t_3}{t_2} = \dots = \frac{t_n}{t_{n-1}}$.

10. Sum to n terms of a G.P

a) If $r > 1$ $S_n = \frac{a(r^n - 1)}{r-1}$
b) If $r < 1$ $S_n = \frac{a(1 - r^n)}{1-r}$
c) If $r = 1$ $S_n = na$
d) Sum of infinite series $S_\infty = \frac{a}{1-r}$

11. If 3 terms in G.P are

$$\frac{a}{r}, a, ar$$

III) Special series

a) $\sum_1^n n = 1 + 2 + 3 + \dots + n = \frac{n(n+1)}{2}$
b) $\sum_1^n n^2 = 1^2 + 2^2 + 3^2 + \dots + n^2 = \frac{n(n+1)(2n+1)}{6}$
c) $\sum_1^n n^3 = 1^3 + 2^3 + 3^3 + \dots + n^3 = \left(\frac{n(n+1)}{2}\right)^2$

9th free coaching classes going on...

Sun Tuition Center- 9629216361

d) $\sum_1^n (2n - 1) = 1 + 3 + 5 + \dots + (2n - 1) = \left(\frac{l+1}{2}\right)$

12. Fibonacci sequence

$$\begin{aligned} F_1 = F_2 = 1 & \quad F_n = F_{n-1} + F_{n+1} & n = 3, 4, 5 \dots \\ 1, 1, 2, 3, 5, 8, 13, 21, 34, 55, 89 \dots \end{aligned}$$

ALGEBRA

1. $(a + b)^2 = a^2 + 2ab + b^2$

2. $(a - b)^2 = a^2 - 2ab + b^2$

3. $a^2 - b^2 = (a+b)(a-b)$

4. i) $(a+b)^3 = a^3 + 3a^2b + 3ab^2 + b^3$
ii) $(a+b)^3 = a^3 + b^3 + 3ab(a+b)$

5. $a^3 + b^3 = (a+b)^3 - 3ab(a+b)$

6. $a^3 + b^3 = (a+b)(a^2 - ab + b^2)$

7. i) $(a-b)^3 = a^3 + 3a^2b - 3ab^2 - b^3$
ii) $(a-b)^3 = a^3 - b^3 - 3ab(a+b)$

8. $a^3 - b^3 = (a-b)^3 + 3ab(a-b)$

9. $a^3 - b^3 = (a-b)(a^2 + ab + b^2)$

10. $(a + b + c)^2 = a^2 + b^2 + c^2 + 2ab + 2bc + 2ac$

11. $(x + a)(x + b) = x^2 + (a + b)x + ab$

12. $(x + a)(x + b)(x + c) = x^3 + (ab + bc + ac)x^2 + (a + b + c)x + abc$

13. General form of a quadratic equation

i) $ax^2 + bx + c = 0$

ii) $x^2 - (\text{sum of roots})x + \text{product of roots} = 0$

14. Let $ax^2 + bx + c = 0$ be a quadratic equation then the value of x is

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

15. Product rule

$$a^m \times a^n = a^{m+n}$$

16. Division rule

$$\frac{a^m}{a^n} = a^{m-n} \quad m < n; a \neq 0.$$

17. Power rule

$$(a^m)^n = a^{mn}$$

18. $(ax \times b)^m = a^m \times b^m$

19. $\left(\frac{a}{b}\right)^m = \frac{a^m}{b^m}$

20. Let α and β are the roots of the equation $ax^2 + bx + c = 0$ then

21. $\alpha^2 + \beta^2 = (\alpha + \beta)^2 - 2\alpha\beta$

22. $|\alpha - \beta| = \sqrt{(\alpha + \beta)^2 - 4\alpha\beta}$

23. $\alpha^3 + \beta^3 = (\alpha + \beta)^3 - 3\alpha\beta(\alpha + \beta)$

24. $\alpha^3 - \beta^3 = (\alpha - \beta)^3 + 3\alpha\beta(\alpha - \beta)$

25. $\frac{1}{\alpha} + \frac{1}{\beta} = \frac{\alpha + \beta}{\alpha\beta}$

26. $\frac{\alpha}{\beta} + \frac{\beta}{\alpha} = \frac{\alpha^2 + \beta^2}{\alpha\beta} = \frac{(\alpha + \beta)^2 - 2\alpha\beta}{\alpha\beta}$

CO – ORDINATE GEOMETRY

1. Distance between any two points

$$D = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2} \text{ units}$$

2. Area of a triangle $A = \frac{1}{2} \begin{vmatrix} x_1 & x_2 & x_3 & x_1 \\ y_1 & y_2 & y_3 & y_1 \end{vmatrix}$ sq. units

3. Area of a quadrilateral $A = \frac{1}{2} \begin{vmatrix} x_1 & x_2 & x_3 & x_4 & x_1 \\ y_1 & y_2 & y_3 & y_4 & y_1 \end{vmatrix}$ sq. units

4. Section formula internally is $P = \left(\frac{lx_2 + mx_1}{l+m}, \frac{ly_2 + my_1}{l+m} \right)$

5. Section formula externally is $P = \left(\frac{lx_2 - mx_1}{l-m}, \frac{ly_2 - my_1}{l-m} \right)$

6. Midpoint formula $M(x, y) = \left(\frac{x_1 + x_2}{2}, \frac{y_1 + y_2}{2} \right)$

7. Centroid of a triangle $G(x, y) = \left(\frac{x_1 + x_2 + x_3}{3}, \frac{y_1 + y_2 + y_3}{3} \right)$

8. Slope of a straight line

a) $m = \tan \theta \quad \theta \neq 90^\circ$ when θ is given

9. $m = \frac{y_2 - y_1}{x_2 - x_1} = \frac{y_1 - y_2}{x_1 - x_2} \quad x_1 \neq x_2$ when two points given

10. $m = \frac{-\text{coefficient of } x}{\text{coefficient of } y} \quad \text{when } ax + by + c = 0 \text{ is given}$

11. Equation of a straight line

a) General format $ax + by + c = 0$

b) Intercept form $y = mx + c$

9th free coaching classes going on...

Sun Tuition Center - 9629216361

- c) One point is given $y - y_1 = m(x - x_1)$
d) Two points are given $\frac{y-y_1}{y_2-x_1} = \frac{x-y_1}{x_2-x_1}$
e) x intercept(a), y intercept (b) is given $\frac{x}{a} + \frac{y}{b} = 1$
12. Equation of x -axis $y = 0$
13. Equation of y -axis $x = 0$
14. Parallel to x -axis $y = k$
15. Parallel to y -axis $x = k$
16. Parallel to $ax + by + c = 0$ is $ax + by + k = 0$.
17. Perpendicular to $ax + by + c = 0$ is $bx - ay + k = 0$.

TRIGONOMETRY

1. $\sin\theta = \frac{\text{Opposite side}}{\text{Hypotenuse side}}$
2. $\cos\theta = \frac{\text{Adjacent side}}{\text{Hypotenuse side}}$
3. $\tan\theta = \frac{\text{Opposite side}}{\text{Adjacent side}}$
4. $\cosec\theta = \text{Reciprocal of } \sin\theta = \frac{\text{Hypotenuse side}}{\text{Opposite side}}$
5. $\sec\theta = \text{Reciprocal of } \cos\theta = \frac{\text{Hypotenuse side}}{\text{Adjacent side}}$
6. $\cot\theta = \text{Reciprocal of } \tan\theta = \frac{\text{Adjacent side}}{\text{Opposite side}}$
7. $\tan\theta = \frac{\sin\theta}{\cos\theta}$
8. $\cot\theta = \frac{\cos\theta}{\sin\theta}$
9. $\cosec\theta = \frac{1}{\sin\theta}$
10. $\sec\theta = \frac{1}{\cos\theta}$

11. Pythagorean identities

i) $\sin^2\theta + \cos^2\theta = 1 \Rightarrow \cos^2\theta = 1 - \sin^2\theta \Rightarrow \sin^2\theta = 1 - \cos^2\theta$

ii) $\cosec^2\theta - \cot^2\theta = 1 \Rightarrow \cosec^2\theta = 1 + \cot^2\theta \Rightarrow \cot^2\theta = \cosec^2\theta - 1$

iii) $\sec^2\theta - \tan^2\theta = 1 \Rightarrow \sec^2\theta = 1 + \tan^2\theta \Rightarrow \tan^2\theta = \sec^2\theta - 1$

12. Trigonometry θ value table

Θ	0°	30°	45°	60°	90°
$\sin\theta$	0	$\frac{1}{2}$	$\frac{1}{\sqrt{2}}$	$\frac{\sqrt{3}}{2}$	1
$\cos\theta$	1	$\frac{\sqrt{3}}{2}$	$\frac{1}{\sqrt{2}}$	$\frac{1}{2}$	0
$\tan\theta$	0	$\frac{1}{\sqrt{3}}$	1	$\sqrt{3}$	∞
$\cosec\theta$	∞	2	$\sqrt{2}$	$\frac{2}{\sqrt{3}}$	1
$\sec\theta$	1	$\frac{2}{\sqrt{3}}$	$\sqrt{2}$	2	∞
$\cot\theta$	∞	$\sqrt{3}$	1	$\frac{1}{\sqrt{3}}$	0

MENSURATION

I) Solid right circular cylinder

- 1) Curved surface Area = $2\pi rh$ sq. units
- 2) Total surface Area = $2\pi r(h + r)$ sq. units
- 3) Volume = $\pi r^2 h$ cu. units

II) Right circular hollow cylinder

- 1) Curved surface Area = $2\pi h(R + r)$ sq. units
- 2) Total surface Area = $2\pi(R + r)(R - r + h)$ sq. units
- 3) Volume = $\pi h(R + r)(R - r)h$ cu. units

III) Solid right circular cylinder

- 1) Curved surface Area = πrl sq. units
- 2) Total surface Area = $\pi r(l + r)$ sq. units
- 3) Volume = $\frac{1}{3}\pi r^2 h$ cu. units

IV) Sphere

- 1) Curved surface Area = $4\pi r^2$ sq. units

9th free coaching classes going on...

Sun Tuition Center - 9629216361

2) Volume = $\frac{1}{3} \pi r^3$ cu. units

V) Hollow sphere Volume = $\frac{4}{3} \pi (R^3 - r^3)$ cu. Units

VI) Solid hemisphere

1) Curved surface Area = $2\pi r^2$ sq. units

2) Total surface Area = $3\pi r^2$ sq. units

3) Volume = $\frac{2}{3} \pi r^3$ cu. Units

VII) Hollow hemisphere

1) Curved surface Area = $2\pi(R^2 - r^2)$ sq. units

2) Total surface Area = $\pi(3R^2 + r^2)$ sq. units

3) Volume = $\frac{2}{3} \pi(R^3 - r^3)$ cu. Units

VIII) A sector of a circle converted into cone

1) CSA of a cone = Area of the sector

$$\pi r l = \frac{\theta}{360^\circ} \times \pi r^2 \text{ cu. Units}$$

IX) Frustum Volume = $\frac{1}{3} \pi h(R^2 + Rr + r^2)$

X) Volume of water flows out through a pipe = (cross section area \times Speed \times Time)

STATISTICS

1. Range = highest value – lowest value

2. The coefficient of range = $\frac{L-S}{L+S}$

3. Standard deviation

i) Direct method $\sigma = \sqrt{\frac{\sum x^2}{n} - (\frac{\sum x}{n})^2}$

ii) Actual mean method $\sigma = \sqrt{\frac{\sum(x-x)^2}{n}}$

iii) Assumed mean method $\sigma = \sqrt{\frac{\sum d^2}{n} - (\frac{\sum d}{n})^2}$ $d = x - A$

iv) Step deviation method $\sigma = \sqrt{\frac{\sum d^2}{n} - (\frac{\sum d}{n})^2} \times C$ $d = \frac{x-A}{C}$

v) Standard deviation for first n Natural numbers $\sigma = \sqrt{\frac{n^2-1}{n}}$

vi) Standard deviation = $\sqrt{\text{variance}}$

vii) Coefficient of variance (C.V) = $\frac{\sigma}{x} \times 100$

PROBABILITY

1. A random experiment is one which is the exact outcome cannot be predicted before conducting the experiment.

2. The set of all possible outcomes of a random experiment is called its sample space. It is denoted by S .

3. $P(A) = \frac{\text{number of outcomes favourable to } A}{\text{total number of outcomes}}$

4. $P(A) = \frac{n(A)}{n(S)}$

5. If A and B are two mutually exclusive events, then $A \cap B = \emptyset$

6. The probability of an event A lies between 0 and 1.

$$0 \leq P(A) \leq 1$$

7. $P(A) + P(\bar{A}) = 1$.

8. Addition theorem on probability

$$\frac{n(A \cup B)}{n(S)} = \frac{n(A)}{n(S)} + \frac{n(B)}{n(S)} - \frac{n(A \cap B)}{n(S)}$$

9. Probability of sure event is 1.

10. Probability of impossible event is 0.

*Life is a good circle,
you choose the best radius...*

sun tuition center

poon thotta pathai hindu mission hospital opposite

villupuram

9th free coaching classes going on...