

A Valuable material from SS PRITHVI's

Class 12



PRITEDUCATION

PRACTISE! PERFORM! PERFECT!

Think Question Papers, Think Priteducation!

JOIN OUR TELEGRAM CHANNEL FOR MORE MATERIALS: [CLICK HERE](#)

A COLLECTION OF

COMPULSORY QUESTIONS

SUBJECT:

M A T H

MR. SS PRITHVI

UPDATED ON: 1/12/2023, 4:25 PM

JOIN OUR TELEGRAM CHANNEL FOR MORE MATERIALS: [CLICK THIS](#)



FIRST MID TERM

1	Write in polar form of the complex number $3 - i\sqrt{3}$.
2	Find the rank of the matrix $\begin{bmatrix} 2 & -2 & 4 & 3 \\ -3 & 4 & -2 & -1 \\ 6 & 2 & -1 & 7 \end{bmatrix}$ by reducing it to an echelon form
3	If $\text{adj}A = \begin{bmatrix} -1 & 2 & 2 \\ 1 & 1 & 2 \\ 2 & 2 & 1 \end{bmatrix}$, find A^{-1} .
4	State and prove triangle inequality of complex number.
5	Construct a cubic equation whose roots are 1, 1, -2
6	Show that the equation $z^3 + 2z$ has five roots
7	$1950x^{20} + 15x^4 + 26x^4 + 2020 = 0$, discuss the nature of roots for this equation.
8	Solve: $x^4 - 10x^3 + 26x^2 - 10x + 1 = 0$



9	Solve : $\sin^2 x - 5 \sin x + 4 = 0$.
10	Find the square root of $-7 + 24i$.
11	Let $A = \{a + \sqrt{5}b : a, b \in \mathbb{Z}\}$. Check whether the usual multiplication is a binary operation on A.
12	Find the angle between the straight line $\frac{x+3}{2} = \frac{y-1}{2} = -z$ with coordinate axes.
13	Find a polynomial equation of minimum degree with rational coefficients, having $2 - \sqrt{3}i$ as a root.
14	Form a polynomial equation with integer coefficients $\sqrt{\frac{\sqrt{2}}{\sqrt{3}}}$ as a roots.
15	Show that the equation $x^9 - 5x^5 + 4x^4 + 2x^2 + 1 = 0$ has atleast 6 imaginary solutions.
16	Find the sum of squares of roots of the equation $2x^4 - 8x^3 + 6x^2 - 3 = 0$.



17	Solve the following system of linear equations using matrix inversion method: $2x - y = 3, 5x + y = 4$
18	Find the square root of $5 - 12i$.
19	If A is a non-singular matrix of odd order, prove that $ \text{adj } A $ is positive.
20	If k is a real, discuss the nature of the roots of the polynomial $2x^2 + kx + k = 0$, in terms of k .
21	Determine the number of positive and negative roots of the equation $x^9 - 5x^8 - 14x^7 = 0$.
22	Solve the equation $3x^3 - 16x^2 + 23x - 6 = 0$ if the product of two roots is 1.
23	Solve: $2x + 3y = 10, x + 6y = 4$ using Cramer's rule.
24	For any vector \vec{a} , prove that $\hat{i} \times (\vec{a} \times \hat{i}) + \hat{j} \times (\vec{a} \times \hat{j}) + \hat{k} \times (\vec{a} \times \hat{k}) = 2\vec{a}$.



25

Find the point of intersection of the lines $\frac{x-1}{2} = \frac{y-2}{3} = \frac{z-3}{4}$ and $\frac{x-4}{5} = \frac{y-1}{2} = z$.



QUARTERLY

1	If the system of linear equation $x+2ay+az = 0$, $x+3by+bz = 0$, $x+4cy+cz = 0$ has a non-trivial solution then show that a, b, c are in H.P.
2	Prove that $\tan\left(\frac{\pi}{4} + \frac{1}{2}\cos^{-1}\left(\frac{a}{b}\right)\right) + \tan\left(\frac{\pi}{4} - \frac{1}{2}\cos^{-1}\left(\frac{a}{b}\right)\right) = \frac{2b}{a}$
3	Find the volume of the parallelopiped whose coterminus edges are given by the vectors $2\vec{i} - 3\vec{j} - 4\vec{k}$, $\vec{i} + 2\vec{j} - \vec{k}$ and $3\vec{i} - \vec{j} + 2\vec{k}$
4	A particle acted on by constant forces $8\vec{i} + 2\vec{j} - 6\vec{k}$ and $6\vec{i} + 2\vec{j} - 2\vec{k}$ is displaced from the point $(1, 2, 3)$ to the point $(5, 4, 1)$. Find the total work done by the forces.
5	Find the angle between the planes $\vec{r} \cdot (\hat{i} + \hat{j} - 2\hat{k}) = 3$ and $2x - 2y + z = 2$
6	Find the equation of the ellipse with foci $(\pm 2, 0)$ and vertices $(\pm 3, 0)$
7	Show that the points $(2, 3, 4)$, $(-1, 4, 5)$ and $(8, 1, 2)$ are collinear.



8	Show that $\vec{r} = (6\hat{i} + \hat{j} + 2\hat{k}) + s(\hat{i} + 2\hat{j} - 3\hat{k})$ and $\vec{r} = (3\hat{i} + 2\hat{j} - 2\hat{k}) + t(2\hat{i} + 4\hat{j} - 5\hat{k})$ are skew lines.
9	If $2\hat{i} - \hat{j} + 3\hat{k}$, $3\hat{i} + 2\hat{j} + \hat{k}$, $\hat{i} + m\hat{j} + 4\hat{k}$ are coplanar, find the value of m.
10	If $ z = 2$ show that $3 \leq z + 3 + 4i \leq 7$.
11	Find the value of $\tan^{-1}(\sqrt{3}) - \sec^{-1}(-2)$.
12	Prove that the point of intersection of the tangents at t_1 and t_2 on the parabola $y^2 = 4ax$ is $(at_1 t_2, a(t_1 + t_2))$
13	If $\omega \neq 1$ is complex cubic root of unity prove that $(2 - \omega)(2 - \omega^2)(2 - \omega^{10})(2 - \omega^{11}) = 49$
14	If e_1 and e_2 are the eccentricities of the ellipse $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$, ($a^2 > b^2$) and the hyperbola $\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$ then prove that $e_1^2 + e_2^2 = 2$.
15	Show that the points $(2, 3, 4)$, $(-1, 4, 5)$ and $(8, 1, 2)$ are collinear



16	Show that $ 3z - 5 + i = 4$ represents a circle and find its centre and radius.
17	If $z = (2 + 3i)(1 - i)$ find z^{-1} .
18	For any vector \vec{a} , prove that $\hat{i} \times (\vec{a} \times \hat{i}) + \hat{j} \times (\vec{a} \times \hat{j}) + \hat{k} \times (\vec{a} \times \hat{k}) = 2\vec{a}$.
19	Prove by vector method that the parallelograms on the same base and between the same parallels are equal in area.
20	Find the angle between the line $\vec{r} = (2\hat{i} - \hat{j} + \hat{k}) + t(\hat{i} + 2\hat{j} - 2\hat{k})$ and the plane $\vec{r} \cdot (6\hat{i} + 3\hat{j} + 2\hat{k}) = 8$
21	Determine whether the three vectors $2\vec{i} + 3\vec{j} + \vec{k}$, and $\vec{i} - 2\vec{j} + 2\vec{k}$ and $3\vec{i} + \vec{j} + 3\vec{k}$ are coplanar.
22	The volume of the parallelepiped whose co terminous edges are $7\vec{i} + \lambda\vec{j} - 3\vec{k}$, $\vec{i} + 2\vec{j} - \vec{k}$, $-3\vec{i} + 7\vec{j} + 5\vec{k}$ is 90 cu.units. Find the value of λ



23	Find the volume of the parallelopiped with its edges represented by the vectors $\hat{i} + \hat{j}$, $\hat{i} + 2\hat{j}$, $\hat{i} + \hat{j} + \pi\hat{k}$
24	If α and β are the roots of $x^2 + x + 1 = 0$ then find the value of $\alpha^{2020} + \beta^{2020}$.
25	If z is a complex number of unit modulus and argument θ . Find the value of $\arg\left(\frac{1+z}{1+\bar{z}}\right)$.
26	If $\vec{a} = 2\hat{i} + 3\hat{j} - 5\hat{k}$, $\vec{b} = -\hat{i} + \hat{j} + 2\hat{k}$ and $\vec{c} = 4\hat{i} - 2\hat{j} + 3\hat{k}$, show that $(\vec{a} \times \vec{b}) \times \vec{c} \neq \vec{a} \times (\vec{b} \times \vec{c})$
27	Find the square root of $7 - 24i$.
28	A ball is thrown vertically upwards, moves according to the law $S = 13.8t - 4.9t^2$ where S is in metres and t is in seconds. (i) Find the velocity at $t = 1$ (ii) Find the acceleration at $t = 1$ (iii) Find the maximum height reached by the ball?
29	If $\vec{a} = \hat{i} - \hat{k}$, $\vec{b} = \hat{j} - \hat{k}$, $\vec{c} = \hat{i} + \hat{j} + \hat{k}$ find $\vec{a} \cdot (\vec{b} \times \vec{c})$ {blurred content is $[\vec{b} \times \vec{c}]$ }



30	Find the inverse of the non - singular matrix $A = \begin{bmatrix} 0 & 5 \\ -1 & 6 \end{bmatrix}$ by Gaus - Jordan method.
31	obtain the polar form of $1 + i \tan \alpha$. where α is an acute angle
32	Solve the equation $x^3 - 9x^2 + 26x - 24 = 0$.
33	Find the acute angle between the planes $\vec{r} \cdot (2\hat{i} + 2\hat{j} + 2\hat{k}) = 11$ and $4x - 2y + 2z = 15$
34	Prove that $\tan^{-1}\left(\frac{1}{2}\right) + \tan^{-1}\left(\frac{1}{3}\right) = \frac{\pi}{4}$
35	Find the length of latus rectum of parabola $y^2 = 4ax$
36	Prove that $ z_1 + z_2 \leq z_1 + z_2 $.
37	If z is a complex number of unit modulus and argument θ . Find the value of $\arg\left(\frac{1+z}{1+\bar{z}}\right)$.



38

If $P = \begin{pmatrix} 1 & \alpha & 3 \\ 1 & 3 & 3 \\ 2 & 4 & 4 \end{pmatrix}$ is the adjoint of a 3×3 matrix A and $|A| = 4$. Find the value of α .



SECOND MID TERM

1	If $U(x, y, z) = \log(x^3 + y^3 + z^3)$, find $\frac{\partial U}{\partial x} + \frac{\partial U}{\partial y} + \frac{\partial U}{\partial z}$.
2	Determine whether * is a binary operation on \mathbb{R} defined by $a * b = a\sqrt{b}$.
3	Assuming $\log_{10} e = 0.4343$ find an approximate value of $\log_{10} 1003$.
4	Prove that in an algebraic structure the Identity element (If exists) must be unique.
5	Let $A = \begin{bmatrix} 1 & 0 & 1 & 0 \\ 0 & 1 & 0 & 1 \\ 1 & 0 & 0 & 1 \end{bmatrix}$, $B = \begin{bmatrix} 0 & 1 & 0 & 1 \\ 1 & 0 & 1 & 0 \\ 1 & 0 & 0 & 1 \end{bmatrix}$, $C = \begin{bmatrix} 1 & 1 & 0 & 1 \\ 0 & 1 & 1 & 0 \\ 1 & 1 & 1 & 1 \end{bmatrix}$ be any three boolean matrices of the same type. Find (i) $A \wedge B$ ii) $(A \wedge B) \vee C$
6	Let $A = \begin{bmatrix} 0 & 1 \\ 1 & 1 \end{bmatrix}$, $B = \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix}$ be any two boolean matrices of the same type Find $A \vee B$ and $A \wedge B$.
7	State and prove commutative laws of conjunction and disjunction by using Truth table
8	Prove that the function $f(x) = x^3$ is strictly increasing on $(-\infty, \infty)$.



9	Evaluate : $\lim_{x \rightarrow 0} x ^{\sin x}$
10	Evaluate: $\int_0^1 x^5 (1-x^2)^5 dx$
11	If $g(x, y) = 3x^2 - 5y + 2y^2$, $x(t) = e^t$ and $y(t) = \cos t$, then $\frac{dg}{dt}$ is equal to
12	Prove that the function $f(x) = x^2 - 2x - 3$ is strictly increasing in $(2, \infty)$
13	Show that the percentage error in the n^{th} root of a number is approximately $\frac{1}{n}$ times the percentage error in the number.
14	Check whether the statement $p \rightarrow (q \rightarrow p)$ is a tautology or a contradiction.
15	Let $U(x, y, z) = x^2 - xy + 3 \sin x$, $x, y, z \in \mathbb{R}$. Find the linear approximation for u at $(2, -1, 0)$.
16	If X is the random variable with distribution function $F(x)$ given by $F(x) = \begin{cases} 0 & x < 0 \\ x & 0 \leq x < 1 \\ 1 & 1 \leq x \end{cases}$ then find i) the probability density function (ii) $P(0.2 \leq x \leq 0.7)$



17

Evaluate $\int_0^8 |x - 5| dx$



Half - Yearly

1	Write the Maclaurin series expansion of e^{-x} .
2	Draw the Geometrical diagram for the sum of two complex numbers Z_1 and Z_2 and verify the result.
3	In the set Q define $a \odot b = a + b + ab$. For what value of y , $3 \odot (y \odot 5) = 7$?
4	Evaluate: $\int_{\pi/6}^{\pi/3} \frac{dx}{1 + \sqrt{\cot x}}$
5	Solve $2x^3 - 9x^2 + 10x - 3 = 0$
6	If $A = \begin{bmatrix} 5 & 3 \\ -1 & -2 \end{bmatrix}$ show that $A^2 - 3A - 7I_2 = 0$ then find A^{-1} .
7	Find the differential equation of the curve represented by $xy = ae^x + be^{-x} + x^2$
8	If $A = \begin{bmatrix} -3 & -2 \\ \lambda & -2 \end{bmatrix}$, find the value of λ so that $A^2 = \lambda A - 2I$



9	If $\omega \neq 1$ is complex cubic root of unity form a quadratic equation with roots 2ω and $2\omega^2$.
10	Evaluate $\int_0^1 \log\left(\frac{1-x}{x}\right) dx$
11	Prove that the function $f(x) = x^2 - 2x - 3$ is strictly increasing in $(2, \infty)$.
12	Evaluate $\int_{\frac{\pi}{6}}^{\frac{\pi}{3}} \frac{dx}{1 + \sqrt{\tan x}}$
13	Evaluate: $\sin(\sin^{-1}(16))$
14	Evaluate: $\int_0^{\frac{\pi}{2}} \log(\tan x) dx$
15	If $ z - 2 + i \leq 2$, then find the greatest value of $ z $



16	The line PP' is a focal chord of the parabola $y^2 = 8x$ and if the coordinates of P are $(18,12)$ then find the coordinates of P'														
17	Find polynomial equation of minimum degree with rational coefficients, having $2 + \sqrt{3}i$ as a root.														
18	Evaluate : $\int_2^3 \frac{\sqrt{x}}{\sqrt{5-x} + \sqrt{x}} dx$														
19	Suppose that $f(x)$ given below represents a probability mass function. <table><tr><td>x</td><td>1</td><td>2</td><td>3</td><td>4</td><td>5</td><td>6</td></tr><tr><td>$f(x)$</td><td>k^2</td><td>$2k^2$</td><td>$3k^2$</td><td>$4k^2$</td><td>k</td><td>$2k$</td></tr></table> Find the value of K .	x	1	2	3	4	5	6	$f(x)$	k^2	$2k^2$	$3k^2$	$4k^2$	k	$2k$
x	1	2	3	4	5	6									
$f(x)$	k^2	$2k^2$	$3k^2$	$4k^2$	k	$2k$									
20	Establish the equivalence property connecting the bi-conditional with conditional: $p \leftrightarrow q \equiv (p \rightarrow q) \wedge (q \rightarrow p)$														
21	If $ z - 2 + i \leq 2$, then find the greatest value of $ z $														
22	The line PP' is a focal chord of the parabola $y^2 = 8x$ and if the coordinates of P are $(18,12)$ then find the coordinates of P'														



23	Evaluate: $\int_0^{2\pi} \sin^7\left(\frac{x}{4}\right) dx$
24	Find the distance between the planes $x+2y+3z+7=0$ and $2x+4y+6z+7=0$
25	Find the modulus and amplitude of $\frac{1+2i}{1-(1-i)^2}$.
26	If $A = \begin{pmatrix} 1 & 3 & 4 \\ 1 & 4 & 5 \\ 2 & 5 & 7 \end{pmatrix}$, find A^{-1} .
27	If $u(x,y,z) = \log(e^x + e^y + e^z)$, find $\frac{\partial u}{\partial x} + \frac{\partial u}{\partial y} + \frac{\partial u}{\partial z}$
28	Verify $(AB)^{-1} = B^{-1}A^{-1}$ with $A = \begin{bmatrix} 4 & 0 \\ -1 & 2 \end{bmatrix}$, $B = \begin{bmatrix} -4 & -2 \\ 1 & -1 \end{bmatrix}$
29	Find df for $f(x) = x^2 + 3x$ for $x = 3$ and $dx = 0.002$



30	Evaluate : $\sin\left[\sin^{-1}\left(\frac{3}{5}\right) + \sec^{-1}\left(\frac{5}{4}\right)\right]$
31	Find polynomial equation of minimum degree with rational coefficients, having $2 + \sqrt{3}i$ as a root.
32	Evaluate : $\int_2^3 \frac{\sqrt{x}}{\sqrt{5-x} + \sqrt{x}} dx$
<div>SS PRITHVI</div>	



Revision - I & 2

1	If the radius of a sphere, with radius 10cm, has to decrease by 0.1 cm, approximately how much will its volume decrease.
2	If $x + y \geq 0$ prove $\cos^{-1}x + \cos^{-1}y = \cos^{-1} [xy - \sqrt{1-x^2} \sqrt{1-y^2}]$
3	Prove that $q \rightarrow p \equiv \neg p \rightarrow \neg q$
4	Establish the equivalence property connecting the bi-conditional $p \leftrightarrow q \equiv (p \rightarrow q) \wedge (q \rightarrow p)$
5	If $w = x + 2y + z^2$ and $x = \cos t$, $y = \sin t$, $z = t$, find $\frac{dw}{dt}$
6	Two balls are chosen randomly from an urn containing 6 red and 8 black balls. Suppose that we win ₹ 15 for each red ball selected and we lose ₹ 10 for each black ball selected. X denotes the winning amount, then find the value of x and number of points in its reverse images.
7	State Rolle's Theorem.



8	Find the distance between the parallel planes $x + 2y - 2z + 1 = 0$ and $2x + 4y - 4z + 5 = 0$
9	The probability density function of X is given by $f(x) = \begin{cases} k x e^{-2x} & \text{for } x > 0 \\ 0 & \text{for } x \leq 0 \end{cases}$ Find the value of k
10	If μ and σ^2 are the mean and variance of the discrete random variable X , and $E(X + 3) = 10$ and $E(X + 3)^2 = 116$, find μ and σ^2 .
11	Find the length of the perpendicular from the point $(1, -2, 3)$ to the plane $x - y + z = 5$
12	Solve $(1 + x^2) \frac{dy}{dx} = 1 + y^2$.
13	Construct truth table for $(p \vee q) \vee \neg q$
14	If $A = \begin{bmatrix} 8 & -4 \\ -5 & 3 \end{bmatrix}$ verify that $A (\text{adj}A) = (\text{adj}A) A = A I_2$



15	Find the volume of the solid formed by revolving the the region bounded by the parabola $y = x^2$, x -axis, ordinates $x = 0$ and $x = 1$ about the x -axis.
16	Find the values in the interval $\left(\frac{1}{2}, 2\right)$ satisfied by the Rolle's theorem for the function $f(x) = x + \frac{1}{x}$, $x \in \left[\frac{1}{2}, 2\right]$.
17	Write the Properties of cumulative distribution function
18	$G = \{1, -1, i, -i\}$ Verify (i) Closure Property (ii) Identity property (iii) Inverse property with respect to complex number Multiplication on G
19	Write the statements in words corresponding to $\sim p$, $q \vee \sim p$, where p is 'it is cold' and q is it is raining.
20	If $A = \begin{bmatrix} 8 & -4 \\ -5 & 3 \end{bmatrix}$ verify that $A(\text{adj } A) = (\text{adj } A) A = A I_2$
21	Express $(\bar{a} + \bar{b} + \bar{c}, \bar{a} - \bar{b}, \bar{c})$ in terms of $(\bar{a} \ \bar{b} \ \bar{c})$



22	Construct the truth table for $(\sim p \vee q) \rightarrow (q \wedge p)$
23	Find the equation of tangent to the curve $y = x^2 - x^4$ at $(1,0)$.
24	Obtain the equation of circle for which $(3,4)$ and $(2,-7)$ are the end of a diameter.
25	If $z_1 = 3$, $z_2 = -7i$, $z_3 = 5 + 4i$ show that $z_1(z_2 z_3) = (z_1 z_2)z_3$.
26	If $A = \begin{pmatrix} 5 & 2 \\ 7 & 3 \end{pmatrix}$ and $B = \begin{pmatrix} 2 & -1 \\ -1 & 1 \end{pmatrix}$ then verify $(AB)^{-1} = B^{-1}A^{-1}$.
27	On \mathbb{Z} , define \otimes by $(m \otimes n) = m^n + n^m : \forall m, n \in \mathbb{Z}$. Is \otimes binary on \mathbb{Z} ?
28	If $\vec{a}, \vec{b}, \vec{c}$ are three vectors prove that $[\vec{a} + \vec{b}, \vec{b} + \vec{c}, \vec{c} + \vec{a}] = 2[\vec{a}, \vec{b}, \vec{c}]$
29	Find a polynomial equation of minimum degree with rational coefficients having $2 - \sqrt{3}$ as a root.



30	Determine whether the pair of straight lines $\vec{r} = (2\hat{i} + 6\hat{j} + 3\hat{k}) + t(2\hat{i} + 3\hat{j} + 4\hat{k})$, $\vec{r} = (2\hat{j} - 3\hat{k}) + s(\hat{i} + 2\hat{j} + 3\hat{k})$ are parallel. Find the shortest distance between them.
31	The probability density function of X is given by $f(x) = \begin{cases} kxe^{-2x} & \text{for } x > 0 \\ 0 & \text{for } x \leq 0 \end{cases}$. Find the value of k
32	Prove De Morgan's law by using Truth table.
33	Find the 'local extrema of the function $f(x) = x^4 + 32x$.
34	Find the length of the perpendicular from the point $(1, -2, 3)$ to the plane $x - y + z = 5$
35	Construct the truth table for the following statements, $\neg(p \wedge \neg q)$
36	If $f(x, y) = \frac{1}{\sqrt{x^2 + y^2}}$ then, prove that $x \frac{\partial f}{\partial x} + y \frac{\partial f}{\partial y} = -f$.
37	If $A = \begin{vmatrix} \cos \theta & \sin \theta \\ -\sin \theta & \cos \theta \end{vmatrix}$ find A^{-1}



38	The time T , taken for a complete oscillation of a single pendulum with length l , is given by the equation $T = 2\pi \sqrt{\frac{l}{g}}$, where g is a constant. Find the approximate percentage error in the calculated value of T corresponding to an error of 2 percent in the value of l .
39	Form the differential equation obtained by eliminating a and b from $y = ae^{3x} + be^{-3x}$ is
40	Evaluate: $\int_0^3 \frac{\sqrt{x} dx}{\sqrt{x} + \sqrt{3-x}}$
41	Show that $\left(\frac{1+i}{1-i}\right)^3 - \left(\frac{1-i}{1+i}\right) = -2i$.
42	Which one of the points $10 - 8i$, $11 + 6i$ is closest to $1 + i$.
43	Let $A = \begin{bmatrix} 1 & 0 & 1 & 0 \\ 0 & 1 & 0 & 1 \\ 1 & 0 & 0 & 1 \end{bmatrix}$, $B = \begin{bmatrix} 0 & 1 & 0 & 1 \\ 1 & 0 & 1 & 0 \\ 1 & 0 & 0 & 1 \end{bmatrix}$ are the boolean matrices. Find i) $A \cup B$ ii) $A \cap B$
44	Compute $P(X = k)$ for the binomial distribution, $B(n, p)$ when $n = 9$, $p = \frac{1}{2}$, $k = 7$



45	<p>Show that the points $1, -\frac{1}{2} + i\frac{\sqrt{3}}{2}$ and $-\frac{1}{2} - i\frac{\sqrt{3}}{2}$ are the vertices of an equilateral triangle.</p>
46	<p>Determine whether the three vectors $2\hat{i} + 3\hat{j} + \hat{k}$, $\hat{i} - 2\hat{j} + 2\hat{k}$ and $3\hat{i} + \hat{j} + 3\hat{k}$ are coplanar.</p>
47	<p>A binary operation $*$ is defined on Q by $a * b = \frac{a+b}{2}$, $\forall a, b \in Q$ verify whether $*$ satisfies closure property, commutative property and associative property.</p>
48	<p>Find the inverse of the non-singular matrix $A = \begin{bmatrix} 0 & -5 \\ -1 & -6 \end{bmatrix}$, by Gauss-Jordan method.</p>
49	<p>Verify $(AB)^T = B^T A^T$ with $A = \begin{bmatrix} -4 & 1 \\ 1 & 3 \end{bmatrix}$, $B = \begin{bmatrix} 4 & -3 \\ 3 & 0 \end{bmatrix}$.</p>
50	<p>Find differential dy for the function $y = (3 + \sin 2x)^2$</p>
51	<p>If the straight lines $\frac{x-5}{5m+2} = \frac{2-y}{5} = \frac{1-z}{-1}$ and $x = \frac{2y+1}{4m} = \frac{1-z}{-3}$ are perpendicular to each other. Find the value of m.</p>



52	Determine whether * is a binary operation on \mathbb{R} , defined by $a * b = a\sqrt{b}$
53	Find the value of $\cos^{-1}\left(\cos \frac{\pi}{7} \cos \frac{\pi}{17} - \sin \frac{\pi}{7} \sin \frac{\pi}{17}\right)$
54	Show that $\sim(p \wedge q) \equiv \sim p \vee \sim q$
55	Prove that $[\vec{a} \times \vec{b}, \vec{b} \times \vec{c}, \vec{c} \times \vec{a}] = [\vec{a}, \vec{b}, \vec{c}]$



Public , Common and PTA

1	Let * be a binary operation on set Q of rational numbers defined as $a * b = \frac{ab}{8}$. Write the identity for *, if any.
2	A fair coin is tossed a fixed number of times. If the probability of getting seven heads is equal to that of getting nine heads, find the probability of getting exactly two heads.
3	If the system of linear equation $x+2ay+az = 0$, $x+3by+bz = 0$, $x+4cy+cz = 0$ has a non-trivial solution then show that a, b, c are in H.P.
4	Prove that $\tan\left(\frac{\pi}{4} + \frac{1}{2}\cos^{-1}\left(\frac{a}{b}\right)\right) + \tan\left(\frac{\pi}{4} - \frac{1}{2}\cos^{-1}\left(\frac{a}{b}\right)\right) = \frac{2b}{a}$.
5	Write the Maclaurin series expansion of e^{-x} .
6	Draw the Geometrical diagram for the sum of two complex numbers Z_1 and Z_2 and verify the result.



7	<p>முனை (2, 1) மற்றும் (1, 3) என்ற புள்ளி வழியாக செல்வதும், இடப்பக்கம் திறப்பு உடையதுமான பரவளையத்தின் சமன்பாடு காண்க.</p> <p>Find the equation of the parabola if the curve is open leftward, vertex is (2, 1) and passing through the point (1, 3).</p>
8	<p>If the lines $\frac{x-x_1}{l_1} = \frac{y-y_1}{m_1} = \frac{z-z_1}{n_1}$ and $\frac{x-x_2}{l_2} = \frac{y-y_2}{m_2} = \frac{z-z_2}{n_2}$ lie on the same plane, then write the number of ways to find the Cartesian equation of the above plane and explain in detail.</p>
9	<p>Show that, if $x = r \cos \theta$, $y = r \sin \theta$, then $\frac{\partial r}{\partial x}$ is equal to $\cos \theta$.</p>
10	<p>Show that $((\neg q) \wedge p) \wedge q$ is a contradiction.</p>
11	<p>Show that the differential equation corresponding to $y = A \sin x$, where A is an arbitrary constant, is $y = y' \tan x$.</p>
12	<p>Show that $\int_0^1 \frac{\sqrt{x}}{\sqrt{1-x} + \sqrt{x}} dx = \frac{1}{2}$.</p>



13	Find the equation of the tangent to the parabola $y^2 = 4ax$ at $(at^2, 2at)$.
14	If $\vec{a}, \vec{b}, \vec{c}$ are three vectors then prove that $[\vec{a} + \vec{b}, \vec{b} + \vec{c}, \vec{c} + \vec{a}] = 2[\vec{a} \vec{b} \vec{c}]$.
15	Show that the distance from the origin to the plane $3x + 6y + 2z + 7 = 0$ is 1.
16	Prove that the general equation of the circle whose diameter is the line segment joining the points $(-4, -2)$ and $(-1, -1)$, is $x^2 + y^2 + 5x + 3y + 6 = 0$.
17	Form the differential equation of the curve $y = ax^2 + bx + c$ where a, b and c are arbitrary constants.
18	Prove that $\int_0^1 x e^x dx = 1$.
19	Express $e^{\cos\theta + i \sin\theta}$ in $a + ib$ form.
20	If $a + b + c = 0$ and a, b, c are rational numbers then, prove that the roots of the equation $(b + c - a)x^2 + (c + a - b)x + (a + b - c) = 0$ are rational numbers.



21	Let $A = \begin{bmatrix} 0 & 1 \\ 1 & 1 \end{bmatrix}$, $B = \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix}$ be any two boolean matrices of the same type. Find $A \vee B$ and $A \wedge B$.
22	Show that the polynomial equation $9x^9 + 2x^5 - x^4 - 7x^2 + 2 = 0$ has at least six imaginary roots.
23	Evaluate $\lim_{x \rightarrow 0} \frac{xe^x - \sin x}{x}$
24	If $\vec{a}, \vec{b}, \vec{c}$ are three vectors prove that $[\vec{a} + \vec{b}, \vec{b} + \vec{c}, \vec{c} + \vec{a}] = 2[\vec{a}, \vec{b}, \vec{c}]$
25	Evaluate : $\int_0^{\frac{\pi}{2}} \frac{\sin x}{1 + \cos^2 x} dx$
26	If $A = \begin{bmatrix} 3 & -2 \\ \lambda & -2 \end{bmatrix}$, find the value of λ so that $A^2 = \lambda A - 2I$.
27	If $A = \begin{bmatrix} 2 & 3 \\ 1 & 2 \end{bmatrix}$, $B = \begin{bmatrix} 4 & 0 \\ 2 & 5 \end{bmatrix}$, find $\text{adj}(AB)$.



28	Find the magnitude and direction cosines of the moment about the point $(0, -2, 3)$ of a force $\hat{i} + \hat{j} + \hat{k}$ whose line of action passes through the origin.
29	If $\omega \neq 1$ is a cubic root of unity and $(1 + \omega)^7 = A + B\omega$, then find A and B .
30	The population of a city grows at the rate of 5 % per year. Calculate the time taken for the population doubles. [Given $\log 2 = 0.6912$]
31	Find the vector equation of the plane passing through the point $(2, 2, 3)$ having 3, 4, 3 as direction ratios of the normal to the plane
32	If $x + iy = \sqrt{\frac{a+ib}{c+id}}$, prove that $x^2 + y^2 = \sqrt{\frac{a^2 + b^2}{c^2 + d^2}}$
33	Find the value of $\cos^{-1}\left(\frac{1}{2}\right) - 2\sin^{-1}\left(-\frac{1}{2}\right)$
34	Find the equation of the tangent to the curve $x^2y - x = y^3 - 8$ at $x = 0$

WITH REGARDS,

SS PRITHVI,

PRIT-EDUCATION.