www.Padasalai.Net

www.TrbTnpsc.com

12

Time : 3.00 Hrs

Register No. 12 A08

# Half-Yearly Examination - 2023 MATHEMATICS

20 x 1 = 20

 $7 \times 2 = 14$ 

Marks : 90

### PART - 1

#### Answer all the questions. Choose the correct answer from the given four alternatives.

1. If  $\rho(A) = \rho((A|B)) < \text{number of unknowns then the system AX = B of linear equations}$ a) consistent and has unique solution b) consistent c) consistent and has infinitely many solutions d) In consistent

2. If 
$$(AB)^{-1} = \begin{pmatrix} 12 & -17 \\ -19 & 27 \end{pmatrix}$$
 and  $A^{-1} = \begin{pmatrix} 1 & -1 \\ -2 & 3 \end{pmatrix}$  then  $B^{-1} = a$   $\begin{pmatrix} 2 & -5 \\ -3 & 8 \end{pmatrix}$  b)  $\begin{pmatrix} 8 & 5 \\ 3 & 2 \end{pmatrix}$  c)  $\begin{pmatrix} 3 & 1 \\ 2 & 1 \end{pmatrix}$  d)  $\begin{pmatrix} 8 & -5 \\ -3 & 2 \end{pmatrix}$ 

- З. How many roots does the quation  $z^2 = \frac{1}{z}$  have? a) 1 b) 2 c) 3 d) 4
- 4 If  $\alpha$  and  $\beta$  are the roots of  $x^2 + x + 1 = 0$  then  $\alpha^{2020} + \beta^{2020}$  is a) -1 b) -2 c) 1 d) 2
- The polynomial  $x^3 kx^2 + 9x$  has three real zeros if and only if k satisfies a)  $|k| \le 6$  b) k = 0 c)  $|k| \ge 6$  d)  $|k| \ge 6$ 5.
- For a 5th degree polynomial equations with real coefficients, which of the following is possible?
- a) it can have 1 real root and 4 purely maginary roots b) it can have 2 real roots and 3 purely imaginary roots
- c) it can have 4 real roots and 1 purely imaginary root d) it can have 5 purely imaginary roots

7. 
$$\sin^{-1}\left(\tan\frac{\pi}{4}\right) - \sin^{-1}\left(\sqrt{\frac{3}{x}}\right) = \frac{\pi}{6}$$
 Then x is a root of the equationa)  $x^2 - x - 6 = 0$  b)  $x^2 - x - 12 = 0$  c)  $x^2 + x - 12 = 0$  d)  $x^2 + x - 6 = 0$ 

8. If 
$$\sin^{-1}x + \sin^{-1}y + \sin^{-1}z = \frac{3\pi}{2}$$
, the value of  $x^{2017} + y^{2016} + z^{2019} - \frac{1}{x^{101} + y^{101} + z^{101}}$  is a) 0 b) 1 c) 2 d) 3

The locus of a point whose distance from (-2, 0) is  $\frac{2}{3}$  times its distance from the line x =  $-\frac{9}{2}$  is

- a) a circle b) a parabola c) an ellipse d) a hyperbola 10. If P(x, y) be any point on  $16x^2 + 25y^2 = 400$  with foci F<sub>1</sub>(3, 0) and F<sub>2</sub>(-3, 0) then PF<sub>1</sub> + PF<sub>2</sub> is a) 6 b) 8 c) 10 d) 12 11. The distance between the plane x + 2y + 3z + 7 = 0 and 2x + 4y + 6z + 7 = 0 is a)  $\frac{\sqrt{7}}{2}$  b)  $\frac{7}{2}$  c)  $\frac{\sqrt{7}}{2\sqrt{2}}$  d)  $\frac{7}{2\sqrt{2}}$ 12. The volume of the parallel piped with its edges represented by the vectors i + j, i + 2j,  $i + j + \pi k$  a)  $\frac{\pi}{2}$  b)  $\frac{\pi}{3}$  c)  $\frac{\pi}{4}$  d)  $\pi$
- The point of inflection of the curve  $y = (x 1)^3$  is a) (0, 1) b) (1, 0) c) (0, 0) d) (1, 1) 14. Angle between  $y^2 = x$  and  $x^2 = y$  at the origin is a)  $\tan^{-1} \left(\frac{3}{4}\right)$  b)  $\tan^{-1} \left(\frac{4}{3}\right)$  c)  $\frac{\pi}{4}$  d)  $\frac{\pi}{2}$
- 15. If  $g(x, y) = 3x^2 5y + 2y^2$ ,  $x(t) = e^t$  and y(t) = cost then  $\frac{dg}{dt}$  is equal to a)  $6e^{2t} + 5sint 4cost sint$  b)  $6e^{2t} 5sint + 4cost sint$  c)  $3e^{2t} 5sint + 4cost sint$  d)  $3e^{2t} + 5sint + 4cost sint$
- 16. If  $f(x, y) = e^{xy}$ , then  $\frac{\partial^2 f}{\partial x \partial y}$  is equal to a)  $xye^{xy}$  b)  $(1 + x)e^{xy}$  c)  $(1 + y)e^{xy}$  d)  $(1 + xy)e^{xy}$
- 17. The value of  $\int \sin^4 x dx$  is a)  $\frac{3\pi}{2}$  b)  $\frac{3\pi}{4}$  c)  $\frac{3\pi}{6}$  d)  $\frac{3\pi}{8}$ 18. If  $\int_{0}^{\pi} \frac{1}{4+x^2} dx = \frac{\pi}{8}$  then a - is a) 1 b) 2 c) 3 d) 4
- 19. The slope at any point of a curve y = f(x) is given by  $\frac{dy}{dx} = 3x^2$  and it passes through (-1, 1). Then the equation of the curve is a)  $y = x^3 + 2$  b)  $y = x^3 + 5$  c)  $y = 3x^2 + 4$  d)  $y = 3x^3 + 4$
- 20. If cosx is the integrating factor of the linear differential equation  $\frac{dy}{dx}$  + Py = Q then P is a) tanx b) cotx c) -cotx d) -tanx PART 2

- 21. If  $\frac{z+3}{z-5i} = \frac{1+4i}{2}$ , find the complex number z in rectangular form.

- 22. Determine the number of positive and negative roots of the equation  $x^9 5x^6 14x^7 = 0$ 23. For what value of x, the inequality  $\frac{\pi}{2} < \cos^{-1}(3x 1) < \pi$  holds? 24. Identify the type of conic section for each of the following equations i)  $3x^2 + 3y^2 4x + 3y + 10 = 0$  ii)  $3x^2 + 2y^2 = 14$
- Find the magnitude and the direction cosines of the torque about the point (2, 0, -1) of a force 2i + j k whose line of action passes 25. through the origin

26. Evaluate :  $x \rightarrow 0 \left(\frac{\sin x}{x^2}\right)$ 

Use the linear approximation to find approximate value of  $\sqrt[3]{26}$ 27

12 Mathematics - 1

#### Kindly send me your answer keys to us - padasalai.net@gmail.com

#### www.Padasalai.Net

### www.TrbTnpsc.com

∫e<sup>i×t</sup>dx valuate the limit

Determine the Order, degree (if exists) of the following differential equation 29

$$\left(\frac{d^2y}{dx^2}\right)^2 + \left(\frac{dy}{dx}\right)^2 = x \sin\!\left(\frac{d^2y}{dx^2}\right)$$

7 x 5 = 35

1 and the decryption matrix as its inverse. Decrypt the received encoded message (10 1) (6 1) with encryption matrix 30

where the system of codes is described by the numbers 1-26 to the letters A-Z respectively, and the number 0 to a blank space PART - 3 7 x 3 = 21

Solve the following system of equations by Cramer's rule.  $\frac{3}{x} + 2y = 12$ ,  $\frac{2}{x} + 3y = 13$ 31

32. Find the value of 
$$\frac{1+\sin\frac{\pi}{10}+i\cos\frac{\pi}{10}}{1+\sin\frac{\pi}{10}-i\cos\frac{\pi}{10}}$$

Obtain the condition that the roots of  $x^3 + px^2 + qx + \gamma = 0$  are in A.P. 33

- Find the domain of cos ' 34 Find the point of intersection of the lines  $\frac{x-1}{2} = \frac{y-2}{3} = \frac{z-3}{4}$  and  $\frac{x-4}{5} = \frac{y-1}{2} = z$ . Write the Maclaurin series expansion for the function  $f(x) = \tan^{-1} x, -1 \le x \le 1$ 35
- 38
- 37
- If U (x, y, z) = log (x<sup>3</sup> + y<sup>3</sup> + z<sup>3</sup>), find  $\frac{\partial u}{\partial x} + \frac{\partial u}{\partial x} + \frac{\partial u}{\partial z}$ Find the area of the region bounded by the y-axis and the parabola x = 5 4y y<sup>2</sup> 38
- Solve [y[1 xtanx] + x<sup>2</sup> cosx] dx xdy = 0 39.
- 40. The Earth is revolving around the Sun in elliptical orbit when Sun is located at one of the focus. If the distance between Sun and the other

focus is 575 x 10° km and eccentricity is  $\frac{1}{2}$  then find the maximum and minimum distance between the earth and sun in earth's orbit. PART - 4

#### Answer all the questions.

ы

Ιź

D

41 a) Test the consistency of the following system of linear equations and solve it if it is consistent. x - y + z = -9, 2x -

$$3x - y + z = 6$$
,  $4x - y + 2z = 7$  (OR) b) if  $z = x + iy$  and arg  $\left(\frac{z-1}{z+2}\right) = \frac{\pi}{4}$ , show that  $x^2 + y^2 + 3x - 3y + 2 = 0$ 

42. a) Identify the type of conic and find centre, foci, vertices and directrices of the following : 9x<sup>2</sup> - y<sup>2</sup> - 36x - 6y + 18 = 0

If 
$$u = \sin^{-1}\left(\frac{x+y}{\sqrt{x}+\sqrt{y}}\right)$$
, show that  $x\frac{\partial u}{\partial y} + y\frac{\partial u}{\partial y} = \frac{1}{2} \tan u$ 

a) Find the local extrema of the function  $f(x) = 4x^6 - 6x^4$  (OR) b) Solve :  $x^4 + 3x^3 - 3x - 1 = 0$ 43

44. a) If 
$$a = 2i+3j-k$$
,  $b = 3i+5j+2k \cdot c = 1-2j+3k$  then find the following  
i)  $a \times (b \times c)$  ii) [a b c] iii) (a c) b (OR) b) Evaluate :  $\int \frac{x}{1+\sin x} dx$ 

45. a) A pot of boiling water at 100°C is removed from a stove at time t = 0 and left to cool in the kitchen. After 5 minutes, the water temperature has decreased to 80°C, and another 5 minutes later it has dropped to 65°C. Determine the temperature of the kitchen. (OR) b) If a., a., a., an is an arithmetic progression with common difference d, prove that

$$\inf_{n} \tan \left[ \frac{d}{1 + a_1 a_2} \right] + \tan \left[ \frac{d}{1 + a_2 a_3} \right] + \dots \tan^{-1} \left[ \frac{d}{1 + a_n a_{n-1}} \right] = \frac{a_n - a_1}{1 + a_1 a_n}$$

a) Find the dimensions of the rectangle with maximum area that can be inscribed in a circle of radius 10cm. (OR) 46

) Solve 
$$(1 + 2e^{y}) dx + 2e^{y} \left(1 - \frac{x}{y}\right) dy = 0$$

a) Parabolic cable of a 60m portion of the roadbed of a suspension bridge are positioned as shown below. Vertical cables are to be 47 spaced every fim along this portion of the roadbed. Calculate the lengths of first two of these vertical cables from the vertex



points (-1, 2, 0), (2, 2, -1) and parallel to the straight

(OR) b) Find the vector parametric vector non-parametric and Cartesian form of the equation of the plane passing through the

t line 
$$\frac{x-1}{2} = \frac{2y+1}{2} = \frac{z+1}{-1}$$

## 12 Mathematics - 2

## Kindly send me your answer keys to us - padasalai.net@gmail.com