

**SRIMAAN COACHING CENTRE-TRICHY-UG-TRB-ZOOLOGY**

**2023-24**

**GRADUATE TEACHERS / BLOCK RESOURCE TEACHER EDUCATOR (BRTE)**

**SRIMAAN**

**UNIT-3- STUDY MATERIAL-TO CONTACT:8072230063.**

# SRIMAAN

## UG-TRB

## ZOOLOGY

**GRADUATE TEACHERS / BLOCK RESOURCE TEACHER EDUCATORS (BRTE)**

**UNIT-3- CELL AND MOLECULAR BIOLOGY (NEW SYLLABUS 2023-2024)**

### UG-TRB MATERIALS

**TAMIL / ENGLISH / MATHS / CHEMISTRY / PHYSICS /  
BOTANY / ZOOLOGY / HISTORY / GEOGRAPHY AVAILABLE.**

**PG-TRB STUDY MATERIALS:-TAMIL/ENGLISH/ MATHEMATICS/PHYSICS  
CHEMISTRY/COMMERCE (T/M & E/M)/BOTANY (T/M & E/M)/ ZOOLOGY  
HISTORY (T/E)/ECONOMICS (T/E)/ GEOGRAPHY /BIO-CHEMISTRY**

**PGTRB-COMPUTER INSTRUCTOR GRADE-I -TO CONTACT -8072230063.**

**SRIMAAN COACHING CENTRE-TRICHY-UG-TRB-ZOOLOGY**

**2023-24**

**GRADUATE TEACHERS / BLOCK RESOURCE TEACHER EDUCATOR (BRTE)**

**SRIMAAN**

**UNIT-3- STUDY MATERIAL-TO CONTACT:8072230063.**

## **TRB-POLYTECHNIC LECTURER MATERIALS:**

**MATHEMATICS / ENGLISH / PHYSICS / CHEMISTRY/COMPUTER SCIENCE/ IT /  
EEE / ECE / EIE/ICE/MECHANICAL/CIVIL/MOP AVAILABLE.**

**UG-TRB: ALL SUBJECT STUDY MATERIALS AVAILABLE.**

**SCERT/DIET/GTTI STUDY MATERIAL AVAILABLE.**

**DEO & BEO (T/M & E/M) STUDY MATERIALS AVAILABLE.**

**TNPSC (CESSE)-(T/M & E/M) STUDY MATERIAL AVAILABLE.**

**PG-TRB: COMPUTER INSTRUCTOR GRADE-1 -FULL  
STUDY MATERIAL WITH QUESTION BANK AVAILABLE**

**TNPSC-ASSISTANT DIRECTOR OF CO-OPERATIVE AUDIT STUDY  
MATERIAL AVAILABLE.**

**TNEB-(ASSESSOR/AE/JA) MATERIALS WITH QUESTION BANK AVAILABLE**

**UG-TRB / PG-TRB / TRB-POLYTECHNIC / DEO & BEO  
MATERIALS ARE SENDING THROUGH COURIER.**

**TO CONTACT**

**8072230063**

**PG-TRB STUDY MATERIALS:-TAMIL/ENGLISH/ MATHEMATICS/PHYSICS  
CHEMISTRY/COMMERCE (T/M & E/M)/BOTANY (T/M & E/M)/ ZOOLOGY  
HISTORY (T/E)/ECONOMICS (T/E)/ GEOGRAPHY /BIO-CHEMISTRY**

**PGTRB-COMPUTER INSTRUCTOR GRADE-I -TO CONTACT -8072230063.**

**SRIMAAN COACHING CENTRE-TRICHY.****TO CONTACT:8072230063.****UG-TRB : ZOOLOGY****UNIT-III-CELL AND MOLECULAR BIOLOGY****Compound microscope****MICROSCOPY-PRINCIPLES AND TYPES**

Microorganisms are too small to be seen by our unaided eyes and the microscopes are of crucial importance as they help to view the microbes. A microscope is an optical instrument consisting of one or more lenses in order to magnify images of minute objects. Thus it is important to gain a preliminary knowledge about the principles of microscope and its types. This chapter gives a brief introduction to microscopy.

**PROPERTIES OF LIGHT**

To understand how a light microscope operates, one must know something about the way in which lenses bend and focus light to form images.

When a ray of light passes from one medium to another, **refraction** occurs, i.e., the ray is bent at the interface. The **refractive index** is a measure of how greatly a substance slows the velocity of light, and the direction and magnitude of bending is determined by the refractive indexes of the two media forming the interface.

When light passes from air into glass, a medium with a greater refractive index, it is slowed and bent toward the normal, a line perpendicular to the surface. As light leaves glass and returns to air, a medium with a lower refractive index, it accelerates and is bent away from the normal. Thus a prism bends light because glass has a different refractive index from air, and the light strikes its surface at an angle. Lenses act like a collection of prisms operating as a unit. When the light source is distant so that parallel rays of light strike the lens, a convex lens will focus these rays at a specific point, the **focal point**. The distance between the center of the lens and the focal point is called the **focal length**. Our eyes cannot focus on objects nearer than about 25 cm or 10 inches. This limitation may be overcome by using a convex lens as a simple magnifier (or microscope) and holding it close to an object. A magnifying glass provides a clear image at much closer range, and the object appears larger. Lens strength is related to focal length; a lens with a short focal length will magnify an object more than a weaker lens having a longer focal length.

**PRINCIPLES OF LIGHT MICROSCOPY**

The light is the primary source on which magnification is based in light microscopes. The magnification is obtained by a system of optical lenses using light waves. Magnification refers the number of times a specimen is appeared to be larger than its original size.

**BASIC UNITS FOR MICROSCOPE**

1 meter = 1000 millimeter

1 millimeter = 1000 micrometer ( $\mu\text{m}$ ) =  $10^{-6}$  meter1 micrometer = 1000 nanometer (nm) =  $10^{-9}$  meter1 Angstrom (1 A) =  $10^{-10}$  meter

1 nanometer = 10 Angstrom

Relative size of the microorganisms and their visibility. Man can see about 0.5 mm sized object whereas the light microscopes can be used to visualize upto 1  $\mu\text{m}$  and EM (electron microscopes) can be used to view 1 nm objects.

## BASIC QUALITY PARAMETERS OF MICROSCOPIC IMAGES

The microscopic images should have four basic quality parameters, through which the microscopes can be graded.

1. **Focus:** It refers whether the image is well defined or blurry (out of focus). The focus can be adjusted through course and fine adjustment knobs of the microscope which will adjust the focal length to get clear image. The thickness of specimen, slide and coverslip also decide the focus of the image. (Thin specimens will have good focus).
2. **Brightness:** It refers how light or the dark the image is. Brightness of the image is depends on the illumination system and can be adjusted by changing the voltage of the lamp and by condenser diaphragm.
3. **Contrast:** It refers how best the specimen is differentiated from the background or the adjacent area of microscopic field. More the contrast will give good images. It depends on the brightness of illumination and colour of the specimen. The contrast can be achieved by adjusting illumination and diaphragm and by adding colour to the specimen. The phase contrast microscopes are designed in such a way that the contrast can be achieved with out colouring the specimen.
4. **Resolution:** It refers the ability to distinguish two objects close to each other. The resolution depends on the resolving power, which refers minimum distance between the two objects which can be distinguishable.

## MAGNIFICATION AND RESOLUTION

The total **magnification** of compound microscope is the product of the magnifications of objective lens and eyepiece. Magnification of about 1500x is the upper limit of compound microscopes. This limit is set because of the resolution.

**Resolution** refers the ability of microscopes to distinguish two objects close to each other, it depends on resolving power, which refers the minimum distance. Ex : Man has the resolving power of 0.2 mm (meaning that he can distinguish two objects with a distance of 0.2 mm close to each other) If he want to see beyond the limit of his resolving power, further magnification is necessary.

$$\text{Resolving power} = \frac{\mu}{n(\sin \theta)}$$

where,  $\mu$  is the wave length of light source and  $n(\sin \theta)$  is the numerical aperture (NA).

For compound microscopes, resolving power is  $\mu/2NA$ . The resolving power of an microscope can be improved either by reducing the wave length of light or by increasing the  $n(\sin \theta)$  value.

**Numerical aperture ( $n \sin^{\theta}$ )** measures how much light cone spreads out between condenser & specimen. More spread of light gives less resolving power means better resolution. The numerical aperture depends on the objective lens of the microscope. There are two types of objective lenses are available in any compound microscope.

### THE LIMIT OF RESOLUTION

The limit of resolution refers the smallest distance by which two objects can be separated and still be distinguishable or visible as two separate objects.

Optical Instrument	Resolving Power	RP in Angstroms
Human eye	0.2 millimeters (mm)	2,000,000 $\text{A}^{\circ}$
Light microscope	0.20 micrometers ( $\mu\text{m}$ )	2000 $\text{A}^{\circ}$
Scanning electron microscope (SEM)	5-10 nanometers (nm)	50-100 $\text{A}^{\circ}$
Transmission electron microscope (TEM)	0.5 nanometers (nm)	5 $\text{A}^{\circ}$

### TYPES OF MICROSCOPE

Microbiologists use a variety of microscopes, each with specific advantages and limitations. Microscopes are of two categories.

**a. Light Microscope:** Magnification is obtained by a system of optical lenses using light waves. It includes (i) Bright field (ii) Dark field (iii) Fluorescence (iv) Phase contrast and (v) UV Microscope.

**b. Electron Microscope:** A system of electromagnetic lenses and a short beam of electrons are used to obtain magnification. It is of two types: (I) Transmission electron microscope (TEM) (ii) Scanning electron microscope (SEM).

### LIGHT MICROSCOPE

Light microscopy is the corner stone of microbiology for it is through the microscope that most scientists first become acquainted with microorganisms. Light microscopes can be broadly grouped into two categories.

**(a) Simple microscope:** It consists of only one bi-convex lens along with a stage to keep the specimen.

**(b) Compound microscope:** It employs two separate lens systems namely, (i) objective and (ii) ocular (eye piece).

### BRIGHT FIELD MICROSCOPE

The compound student microscope is a bright field microscope. It consists of mechanical and optical parts.

#### 1. Mechanical parts

These are secondary but are necessary for working of a microscope. A 'Base', which is horsehoe, shaped supports the entire framework for all parts. From the base, a 'Pillar' arises. At the top of the pillar through an 'Inclination Joint' arm or limb is attached. At the top of the pillar, a stage with a central circular opening called 'Stage aperture' is fixed, with a stage clip to fix the microscopic slide. Beneath the stage, there is one stage called 'sub stage' which carries the condenser. At the top of the arm, a hollow cylindrical tube of standard diameter is attached in-line with the stage aperture, called 'body tube'. The body tube moves up and down by two separate arrangements called 'coarse adjustment' worked with pinion head and 'fine adjustment' worked with micrometer head. At the bottom of the body tube an arrangement called 'revolving nose-piece' is present for screwing different objectives. At the top of the body tube eye-piece is fixed.

## 2. Optical parts

It includes mirror, condenser, objective and ocular lenses. All the optical parts should be kept in perfect optical axis.

**a. Objectives :** Usually 3 types of magnifying lenses (i) Low power objective (10x) (ii) High dry objective (45x) and (iii) Oil immersion objective (100x)

**b. Eye-piece :** Mostly have standard dimensions and made with different power lenses. (5x, 10x, 15x, 20x). A compound microscope with a single eyepiece is said to be monocular, and one with two eyepieces is said to be binocular.

**c. Condenser :** Condenses the light waves into a pencil shaped cone thereby preventing the escape of light waves. Also raising or lowering the condenser can control light intensity. To the condenser, iris diaphragm is attached which helps in regulating the light.

**d. Mirror :** It is mounted on a frame and attached to the pillar in a manner that it can be focused in three different directions. The mirror is made of a lens with one plane surface and another concave surface. Plane surface is used, when the microscope is with a condenser.

**In case of microscopes with oil immersion,** when light passes from a material of one refractive index to material of another, as from glass to air or from air to glass, it bends. The refractive index of air is 1.0, which is less than that glass slide (1.56). So, when light passes from glass (dense medium) to air (lighter medium), the rays get refracted, which led to loss of resolution of image. Light of different wavelengths bends at different angles, so that as objects are magnified the images become less and less distinct. This loss of resolution becomes very apparent at magnifications of above 400x or so. Even at 400x the images of very small objects are badly distorted. Placing a drop of oil (Cedar wood oil) with the same refractive index (1.51) as glass between the cover slip and objective lens eliminates two refractive surfaces and considerably enhances resolution, so that magnifications of 1000x or greater can be achieved. Oil immersion is essential for viewing individual bacterial cell. A disadvantage of oil immersion viewing is that the oil must stay in contact, and oil should be viscous.

## DARK-FIELD MICROSCOPE

In dark-field microscopy, specimen is brightly illuminated against a dark background. This type of microscope possesses a special type of condenser, which prevents the parallel and the

oblique rays entering in to the objective and thus making the microscopic field dark. In the absence of specimen the entire field will appear as dark. In the presence of specimen, which differs in refractive index, the oblique rays are scattered by reflection and refraction and the scattered rays enter the objective making the specimen brightly illuminated.

#### *Dark-Field microscope*

Maximum magnification of 1500x and resolution of 0.1 – 0.2  $\mu\text{m}$  can be obtained. It is useful in studying the morphology and motility of microorganisms. Dark field is especially useful for finding cells in suspension. Dark field makes it easy to obtain the correct focal plane at low magnification for small, low contrast specimens. The following are its uses:

- a. Initial examination of suspensions of cells such as yeast, bacteria, small protists, or cell and tissue fractions including cheek epithelial cells, chloroplasts, mitochondria, even blood cells (small diameter of pigmented cells makes it tricky to find them sometimes despite the color).
- b. Initial survey and observation at low powers of pond water samples, hay or soil infusions, purchased protist or metazoan cultures.
- c. Examination of lightly stained prepared slides. Initial location of any specimen of very small size for later viewing at higher power.
- d. Determination of motility in cultures

#### **PHASE CONTRAST MICROSCOPE**

In the 1930's Frederick Zernike devised a method of converting phase changes into differences in light intensity. This invention leads to the development of phase contrast microscope.

It has a special type of condenser, objective and a special magnifier. Light passing from one material into another of slightly different refractive index will undergo a change in the phase. This change in the phase of the light will in-turn increase the contrast. A system of rings in the condenser and objective separate the diffracted rays from the specimen. These diffracted rays from the specimen and the undiffracted rays combine and the phase difference is converted into difference in light intensity. In phase-contrast microscopy, unstained/living organisms can be examined. As a result, the internal contrast of various parts of a specimen against its surroundings is increased. It has a resolving power of 0.1 – 0.2  $\mu\text{m}$ .

**To the left:** Arrangement of the ring-shaped mask below the objective and of the phase-ring within the objective

**To the right:** The path of light rays within a phase-contrast microscope. 1. ring-shaped mask, 2. Condenser, 3. Specimen, 4. Objective, 5. Phase plate, 6. Focal plane of the objective. The wave character of the light is indicated by the change of light and dark areas.

#### **UV MICROSCOPE**

Resolution of a microscope depends upon the wavelength of light used. If, longer the wavelength of light used, lower will be the resolving power while shorter the wavelength,

more will be the resolution. With this principle, UV rays of shorter wavelength are used as light source. Since UV rays can't penetrate the glass, quartz lenses are used. Since the UV rays are invisible, photographic plates should be used to record the image or special type of filters should be used to eliminate the UV rays from reaching the eyepiece. This is used in conjunction with fluorescent microscopy. Upon illumination with UV light certain fluorescent dyes emit light in visible range, which can be directly viewed.

## FLUORESCENCE MICROSCOPE

Certain chemical compounds absorb light and reemit part of the radiant energy as light of longer wavelength. Such substances are called fluorescent and the phenomenon is termed as fluorescence. In fluorescence microscopy, a high intensity mercury lamp is used as the light source, which emits white light. The exciter filter transmits only blue light to the specimen and blocks out all the colours. The blue light is reflected downward to the specimen by the dichroic mirror. The specimen is stained with fluorescent dye (acrydine orange). Only certain portions of the specimen retain the dye, others do not. The stained portion of the specimen absorb blue light and emit green light, which passes upwards, penetrate the dichroic mirror and reaches the barrier filter. This filter allows only green light to pass through and the eye receives only green light emitted from the specimen against the black background whereas unstained portions are invisible. Also, ultraviolet light is used to excite molecules so that they release light of a different wavelength.

This technique is especially important in **immunology** in which the reactions of antigens and antibodies are studied in great detail. Fluorescent antibody staining is now widely used in diagnostic procedures to determine whether an antigen is present. Not all bacteria get stained with fluorescent chemicals. *Excitation and fluorescence with chromatic beam splitters.* Similar to the interference filters these are specially coated mirrors used under  $45^\circ$  to the illuminating beam. They reflect certain spectral ranges, while others are completely transmitted. The separating line between reflection and transmission may be set at any point of the spectrum. 1. Exciting radiation, 2. Fluorescence emission

## Confocal Laser Scanning Microscope (CLSM)

A schematic diagram of a confocal laser scanning microscope is shown in figure 16.2A. Let us see how exactly a CLSM works:

- i. Light source and illumination: Light sources used in confocal microscopes are lasers. The microscope works in epi-illumination mode. The laser beam is spread by a diverging lens so as to fill the back aperture of the objective lens which functions as condenser as well. The expanded laser light is reflected by the dichroic mirror on the objective that focuses the light as an intense diffraction-limited spot on the sample. The fluorescence from the illuminated spot is collected by the objective and sent to the eyepiece/camera/detector through a pinhole aperture.
- ii. Pinhole aperture: The fluorescence light emitted by the illuminated sample is focused as the confocal point at the pinhole. Any light coming from below or above the focal plane is blocked by the pinhole plate.
- iii. Raster scanning: As the fluorescence is detected from a diffraction limited spot, the focused laser spot is scanned over the sample in a raster fashion collecting light from the entire focal plane (Figure 16.3A). The laser spot is scanned over the sample by changing the direction of the incident radiation as shown in Figure 16.3B. As the position of the illuminating spot changes, the pinhole moves so as to be confocal with the illuminated spot of the same focal plane.
- iv. Emission filter: The light that passes through the pinhole is filtered by the emission filter before it reaches the detector.



## ELECTRON MICROSCOPE

In electron microscope, short beam of electrons and magnetic condenser lenses are employed to produce the image. The electrons have short wavelength, which helps in better resolution. It is possible to resolve objects as small as  $10^{\circ}\text{A}$ , which is 100 times more than that of light microscope. It can magnify object up to 200,000X.

In electron microscope, a hot tungsten filament forms the source of electrons. The object is placed in the path of moving electrons. Since electrons move only in the vacuum, the entire path of electrons should be kept under vacuum. The magnetic condenser lens causes the primary magnification. A second magnetic lens amplifies the primary image and this image is viewed on a fluorescent screen or captured on photographic plates. There are two types of electron microscope.

- a. Transmission electron microscope (TEM)
- b. Scanning electron microscope (SEM)

## TRANSMISSION ELECTRON MICROSCOPE

A high voltage established between the filament and the anode accelerates the electrons from the hot tungsten filament. The electron beam is focused on the specimen with an electromagnetic condenser. Ultra thin sections of the specimen must be prepared since electrons can penetrate matter only a short distance. This is done by embedding or freezing the specimen and sectioning it with a diamond or a glass Knife. The sections are floated in water and collected in a copper grid. They are stained with heavy metals such as gold or palladium and kept within the evacuated column of the electron microscope.

TEM has a projector lens that project the image onto a fluorescent viewing screen or film plate, because the beam cannot be viewed directly. With TEM greater resolution and higher magnifications than light and Scanning Electron Microscope can be obtained. In TEM, the differential scattering of electrons by the specimen makes the contrast. Since most of the atoms of the biological material are of low mass, the contrast of the specimen is low. Staining with heavy metals such as platinum, uranium or tungsten can increase the contrast.

They are of various types such as,

- a. **Positive staining:** The heavy metals are fixed on the specimen.
- b. **Negative staining:** It is used to increase the electron opacity of the surrounding area.

Two techniques commonly employed for the observation of biological specimen are:

- a. **Metal shadowing:** The dried specimen is exposed at an acute angle to a stream of heavy metals like platinum, palladium or gold and thereby producing an image, that reveals the three-dimensional structure of the object.
- b. **Freeze fracturing:** The frozen specimen is fractured with a knife, and the exposed surface is coated with a heavy metal (Gold) at an acute angle. A supporting layer of carbon is evaporated on the metal surface. Then the specimen is destroyed and the replica is examined. This method is used for studying cell wall and cell membrane. The penetrating power of electrons is low; hence the ultra thin sections of the specimens should be used.

## SCANNING ELECTRON MICROSCOPE

The specimen is coated with a thin layer of heavy metal and the specimen is subjected to a narrow beam of electrons, which rapidly moves and scans the surface of the specimen. The irradiated specimen depending upon its physical and chemical composition will release secondary electrons. These secondary electrons are then collected by anode detector, which generates an electronic signal. Then the electronic signal is scanned in TV system to produce an image on a cathode ray tube. Magnification on SEM is about 75,000 to 1,00,000 times.

### Limitations

- Specimen is kept under high vacuum on the path of electron beam. So, living cells can't be examined.
- Electrons have low penetration capacity, hence ultra thin section and staining should be done which is time consuming and also sometimes alter or distort the structures of microorganisms.
- High cost and specialized techniques prevent its use in all microbiology labs in spite of greater magnification and resolution.

## .ATOMIC FORCE MICROSCOPE

Atomic force microscope is a type of scanning probe microscope that records the force between the probe and the specimen. The working principle of an AFM can be understood like this: Consider yourself to be in a dark room in front of a table. The table has a book, a pen, a wristwatch, a spoon, a fork, and a screw driver. Will you be able to selectively lift the spoon if asked to do so? The answer for most people is yes. You can distinguish two distinct objects by touching them with your fingers. In this example, your fingers act as the probes, your arm acts as the positioner of your fingers, and your brain works as the processing unit. An AFM works exactly the same way; it has three basic components: a probe, a positioner, and a processing unit.

### Modes of operation

An AFM experiment can be recorded in both attractive and repulsive regimes of the Lennard-Jones potential. There are three basic modes of AFM imaging. Another mode, called force spectroscopy is not used for imaging but for characterizing physico-chemical properties of the

*Contact mode AFM:* In contact mode AFM, the tip is brought in close contact with the specimen (in the repulsive regime) and scanned over the surface. As the tip is in contact with the sample throughout the scan, the frictional forces are very high. This mode of operation therefore may not be suitable for soft samples including biological samples.

*Non-contact mode AFM:* In non-contact mode AFM, a cantilever with very high spring constant is oscillated very close to the sample (in the attractive regime). The quantities that are measured are changes in the oscillation amplitude and the phase. The forces between the tip and the sample are very small, of the order of piconewtons. This mode is therefore well-suited for very soft samples but resolution is compromised. *Intermittent mode or tapping mode AFM:* A stiff cantilever is oscillated so close to the specimen that a small part of oscillation lies in the repulsive regime of the Lennard-Jones potential. The tip therefore intermittently touches the sample while scanning. This mode of imaging allows imaging with very high resolution and has become the method of choice for scanning the soft biological samples.

*Force mode AFM/Force spectroscopy:* Force mode of AFM is not an imaging mode. Briefly, the sample is brought close to the cantilever, pushed against it causing deflections in it, and then

withdrawn. A plot of force (depends on the spring constant of the cantilever) against the distance is called a force spectrum. Force spectroscopy mode is often used to study the interactions of the tip with the sample and to determine the mechanical properties of the specimen.

### **Resolution**

Atomic force microscopes can provide resolutions comparable to that obtained with electron microscopes. As neither light nor particles are used to generate the images, resolution of atomic force microscopes does not depend on any wavelength. The resolution of an AFM is determined by the shape and the diameter of the tip. It is also evident that the resolution in the X-Y plane is poorer as compared to that in the Z-direction. A Z-resolution of ~0.2 nm or better is often achieved using AFM

### **Advantages of AFM**

Both AFM and EM provide very high resolution images but AFM has few distinct advantages over EM:

- i. Easy sample preparation: AFM does not involve a tedious sample preparation. A sample to be analyzed can simply be placed on a smooth surface and scanned.
- ii. Imaging in solution: Unlike EM; it is possible, in fact routine; to record AFM images in solution. No other microscopic method, except the scanning probe microscopes, provides a subnanometer resolution in solution.
- iii. Manipulation: An AFM tip can be used to mechanically manipulate the specimen at very high spatial resolution.

## **CELL FIXATION**

In the preceding section you have learnt about the basic principle of microscopy and the working of a compound microscope. However, before we start observing specimens, it is necessary to fix cells or tissues so that we can preserve them in a near lifelike condition. Cell fixation prevents autolysis and putrefaction of the specimen and also protects the tissue / cells during any further processing steps.

Cell fixation can be done either by **physical or chemical** means. Physical fixation is carried out by increasing (heating) or decreasing (cryopreservation) the temperature whereas chemical fixation depends on the use of either organic solvents or cross linking agents. Generally chemical fixation methods are used to preserve most specimens.

The **organic solvents** include alcohol and acetone that cause removal of lipids; dehydration of tissues; denaturation and precipitation of proteins; changes in protein conformation and their solubility. They are also referred to as **coagulants** based on the effect produced. The dehydrating effect of ethanol can be reduced by the swelling effect of acetic acid and they are together used in some cytological fixatives.

The **cross linking methods** uses aldehydes (such as paraformaldehyde, glutaraldehyde) or osmium tetroxide that reacts with proteins and other biomolecules by forming intra- and inter molecular cross links. The covalent bonds result in a better retention of cellular architecture than organic solvents.

The rigidity imparted to cells by cross linking is an added asset in sectioning fragile tissues. Fixing kills the cells immediately but allows us to visualize its morphology and internal structures that were present just before fixing. This is helpful because once the cells are fixed then the investigator can observe them at ease. Fixing also permeabilises cells.

## Fixation

Fixation of tissue is the first and crucial step for specimen preparation. Biological specimen can be fixed either with chemical fixatives or freezing method. The advantages of tissue fixation are many. We have discussed this step in the preceding unit also as it is utilised in some forms of light microscopic observations.

It serves to:

- \* preserve cells and tissue components as close to their living state as possible with minimum alterations,
- \* protect cell from autolysis and decomposition, and
- \* protect and stabilize cellular structure from subsequent treatments such as dehydration, staining and from irradiation by electron beam.

## Staining

Staining is the last step in which the sectioned tissue is stained with a solution of heavy metal salts to increase the light scattering power of the section. A number of stains are available to enhance contrast as many biological materials are transparent to electrons. These stains fall into two major categories - positive and negative stains. A drop of heavy metal salts is deposited onto to sectioned tissue and then the stained specimen is placed on a small circular metal grid for viewing the image under the electron microscope. The sample can also be stored and observed later. In positive staining, the image appears as dark area against light background as these stains deposit electron dense material on the area of interest. Lead citrate and uranyl acetate are common stains.

The positive stains are used to view specific molecules within cells, for instance proteins, lipids and DNA. The Uranyl ions react strongly with phosphate and amino groups, staining DNA and some proteins whereas lead citrate is taken up more strongly by lipid containing membranes and it produces a general increase in contrast.

The negative stains are used to view intact biological structure such as sub cellular organelles, bacteria, viruses. These stains penetrate and darken the interstices between areas of interest and produce an image in which the specimen appears light against a stained dark background. Negative staining technique developed by Brenner and Horne 1959.

## Centrifugation

**Centrifugation** is a technique that uses centrifugal force to separate and purify biological components such as cells and subcellular organelles, suspended in liquid medium. The particles are separated at different sedimentation rate according to their size, shape or density in an applied centrifugal force.

This is the last unit of Block 1 deals with the principle, types and application of centrifugation. This unit begins with principle of centrifugation. Here you will learn about sedimentation coefficient, Svedberg unit and relative centrifugal force. This will be followed by a description of the type of centrifuges and rotors. You will also learn about the working principle and applications of differential centrifugation and density gradient centrifugation (Rate Zonal and Isopycnic centrifugations) which are widely used in Cell and Molecular biology. The unit ends with safety measures and centrifuge maintenance.

### PRINCIPLE OF CENTRIFUGATION

The basic principle of centrifugation is based on sedimentation. Sedimentation is the tendency of particles in liquid suspension to settle down gradually to the bottom of the tube under the influence of the earth's gravitational force. When a particle moves in a circular motion, it experiences two forces *i.e.* centrifugal force which is outward force and centripetal force which is towards the central axis.

Larger particles tend to move faster than smaller ones;

| A denser particle moves faster rate than less dense ones;

| The denser the solution, more slowly the particle will move, and | Greater the frictional coefficient, the slower the particle will move.

**The sedimentation coefficient (s)** is the ratio of sedimentation velocity ( $v$ ) and centrifugal force

( $G$ ). It is a constant for a given particle or solvent. More often, the notation provided by **Theodor**

**Svedberg** is used to describe sedimentation coefficients. One Svedberg is taken as  $1 \times 10^{-13}$  seconds.

Many biological components are described by their sedimentation coefficients.

## Type of Centrifuges

All centrifuges are based on the same principle but they differ in the maximum speed of rotation, rotor designs, temperature control and presence or absence of vacuum. We will describe here briefly four major type of centrifuges.

**Benchtop centrifuge:** Modern benchtop centrifuge (Fig 4.6a) is a low speed (3000- 7000 x g) centrifuge that can be with or without a cooling system. It is routinely used in clinical laboratories for the separation the blood cells, plasma and serum as well as in cell culture applications.

**Micro centrifuge (microfuge):** Microcentrifuge is widely used for routine laboratory work. The sample volumes centrifuged are small (0.5 – 1.5ml) and it can accommodate 24 samples. It generates up to 15,000 x g, with or without refrigeration. It is primarily used to spin small centrifuge tubes (Eppendorf) at high speed. It is useful formany biological applications such as pelleting of nucleic acids and for the concentration of proteins samples..

**Refrigerated centrifuges:** They are refrigerated and can generate high centrifugal force up to 60,000 x g. Depending on the rotor sizes.and type, variable number of tubes and volumes can be centrifuged. It is used for differential separation of organelles, separating fractions during partial purification using ammonium sulphate or in obtaining tissue extracts and other temperature sensitive separations

## CELL TYPE

### Introduction

A structure containing a mass of cytoplasm surrounded by semi-permeable membrane called plasma membrane is called a cell. It encloses cytoplasm, many cell organelles along with nucleus or nuclear material. On the basis of organization of membranes, variety and structure of cytoplasmic organelles and complexity of nuclear region, the cells are classified into two types: Prokaryotic cell and Eukaryotic cell. These terms were suggested by **Hans Ris** in **1960s**.

### History and Origin

A cell was defined as “unit of biological activity delimited by a semi permeable membrane and capable of self-reproduction in a medium free of other living systems” by **Loewy and Siekevitz (1963)**.

The study of cell has been made possible with the help of light microscope. **Robert Hooke (1665)** with the help of light microscope discovered that a section of cork is made up of small cavities surrounded by firm walls. He used the term “**cell**” for the first time to describe his investigations on the “texture of a piece of cork”.

Later on **A. Van Leeuwenhoek (1632-1723)** observed various unicellular organisms and cells like bacteria, protozoan’s, red blood cells and sperm etc. He observed nucleus in some erythrocytes and all this was made possible with the improved microscopes. In **1809, Mirble M.** stated that all plant tissues are composed of cells. In the same year, importance of cells in living organisms was described by **J.B. Lamarck**. **Robert Brown** in **1831** observed nucleus in certain plant cells. *Mimosa* cells were boiled in nitric acid by **Dutrochet (1837)** to separate the cells to conclude that all organic tissues are composed of globular cells, united by simple adhesive forces. “All living organism are composed of cells” was stated by **Schwann, T. (1839)** after examining a variety of animals and plant tissues.

## BASIC COMPONENTS OF PROKARYOTIC AND EUKARYOTIC CELL

### Prokaryotic Cells

Prokaryotic cells are the most primitive cells and have simple structural organization. It has a single membrane system. They include bacteria, viruses, blue-green algae, mycoplasmas, rickettsias, spirochetes etc. Cyanobacteria or blue green algae are the largest and most complex prokaryote, in which photosynthesis of higher plants type have evolved. **Prokaryotes** are included in the kingdom **Monera** and the super kingdom **Prokaryota**. The Prokaryotes have the following characters:

1. The size of prokaryotic cells ranges between 1 to 10  $\mu\text{m}$ . They occur in a variety of forms.
2. Prokaryotic cell consists of three main components:
  - (I) **Outer covering:** It is composed of inner cell or plasma membrane, middle cell wall and outer slimy capsule.
    - a. **Cell membrane:** Cell membrane made up of lipids and proteins, is thin and flexible and controls the movement of molecules across the cell. Respiratory enzymes are carried by it for energy releasing reactions. **Mesosomes**, the in-folds of plasma membrane

**STUDY MATERIALS AVAILABLE**

**IF YOU NEED : UG-TRB: ZOOLOGY FULL STUDY MATERIAL AVAILABLE (NEW SYLLABUS 2023-2024).**

- \* **UGTRB (ALL SUBJECT) STUDY MATERIALS AVAILABLE.**
- \* **TET-PAPER-1,PAPER-2 STUDY MATERIALS WITH Q.BANK AVAILABLE**
- \* **TRB-POLYTECHNIC LECTURER (ALL SUBJECT) STUDY MATERIAL WITH QUESTION BANK.**

**SRIMAAN COACHING CENTRE-TRICHY.**

**TO CONTACT:8072230063.**

**SRIMAAN COACHING CENTRE-TRICHY- TET/PG-TRB / UG-TRB  
BEO/ DEO/TRB-POLYTECHNIC/COM.INS/TNEB/ SCERT / JSO  
STUDY MATERIALS AVAILABLE- CONTACT:8072230063.**

**2023-24  
SRIMAAN**

# **UG-TRB MATERIALS**

**GRADUATE TEACHERS / BLOCK RESOURCE TEACHER EDUCATORS (BRTE)**

- **UG TRB: TAMIL MATERIAL WITH QUESTION BANK.**
- **UG TRB: ENGLISH STUDY MATERIAL +Q. BANK.**
- **UG-TRB: MATHEMATICS MATERIAL WITH Q. BANK (E/M)**
- **UG TRB: PHYSICS MATERIAL WITH QUESTION BANK (E/M)**
- **UG TRB: CHEMISTRY MATERIAL + QUESTION BANK (E/M)**
- **UG TRB: HISTORY MATERIAL + Q.BANK (E/M)**
- **UG TRB: ZOOLOGY MATERIAL + QUESTION BANK (E/M)**
- **UG TRB: BOTANY MATERIAL +QUESTION BANK (T/M& E/M)**
- **UG TRB: GEOGRAPHY STUDY MATERIAL (E/M)**

**SCERT/DIET/GTTI (LECTURER) STUDY MATERIAL AVAILABLE.**

**TRB-POLYTECHNIC LECTURER-(NEW SYLLABUS)  
STUDY MATERIALS AVAILABLE**

- **MATHEMATICS STUDY MATERIAL with Question Bank.**
- **ENGLISH STUDY MATERIAL with Question Bank.**
- **PHYSICS STUDY MATERIAL with Question Bank.**



**SRIMAAN COACHING CENTRE-TRICHY- TET/PG-TRB / UG-TRB  
BEO/ DEO/TRB-POLYTECHNIC/COM.INS/TNEB/ SCERT / JSO  
STUDY MATERIALS AVAILABLE- CONTACT:8072230063.**

**2023-24  
SRIMAAN**

- **CHEMISTRY STUDY MATERIAL with Question Bank.**
- **MODERN OFFICE PRACTICE STUDY MATERIAL with Q.B.**
- **COMPUTER SCIENCE STUDY MATERIAL with Question Bank.**
- **INFORMATION TECHNOLOGY STUDY MATERIAL with Q.Bank.**
- **ECE STUDY MATERIAL with Question Bank.**
- **EEE STUDY MATERIAL With Question Bank.**
- **MECHANICAL STUDY MATERIAL With Question Bank.**
- **CIVIL STUDY MATERIAL With Question Bank.**
- **EIE STUDY MATERIAL with Question Bank.**
- **ICE STUDY MATERIAL with Question Bank.**

## **PG-TRB MATERIALS**

- **PG TRB: TAMIL STUDY MATERIAL +QUESTION BANK (T/M)**
- **PG TRB: ENGLISH MATERIAL WITH QUESTION BANK.**
- **PG-TRB: MATHEMATICS MATERIAL WITH Q.BANK (E/M)**
- **PG TRB: PHYSICS MATERIAL WITH QUESTION BANK (E/M)**
- **PG TRB: CHEMISTRY MATERIAL + QUESTION BANK (E/M)**
- **PG TRB: COMMERCE MATERIAL WITH Q.BANK (T/M)&(E/M)**
- **PG TRB:ECONOMICS MATERIAL+Q. BANK (T/M & E/M)**

**SRIMAAN COACHING CENTRE-TRICHY- TET/PG-TRB / UG-TRB  
BEO/ DEO/TRB-POLYTECHNIC/COM.INS/TNEB/ SCERT / JSO  
STUDY MATERIALS AVAILABLE- CONTACT:8072230063.**

**2023-24  
SRIMAAN**

➤ **PG TRB: HISTORY MATERIAL + Q. BANK (T/M & E/M)**

➤ **PG TRB: ZOOLOGY MATERIAL + QUESTION BANK (E/M)**

➤ **PG TRB: BOTANY MATERIAL +QUESTION BANK (T/M& E/M)**

➤ **PG TRB: GEOGRAPHY STUDY MATERIAL (E/M)**

**TNPSC-(CESE / JSO) STUDY MATERIAL AVAILABLE.**

**PG-TRB: COMPUTER INSTRUCTOR-GRADE-I (NEW  
SYLLABUS)-2023-2024 STUDY MATERIAL AVAILABLE  
TNPSC-GROUP-I C**

**TNPSC-DEO (District Educational Officer(Group – I C Services)  
(TAMIL & ENGLISH MEDIUM) STUDY MATERIAL AVAILABLE.**

**TRB-BEO (Block Educational Officer)  
(TAMIL & ENGLISH MEDIUM) STUDY MATERIAL AVAILABLE.**

**10% Discount for all materials. Materials are sending through**

**COURIER.**

**TO CONTACT**

**8072230063**

**SRIMAAN**