

திரைக் குறியீடு தேர்வு: 2024

பாடம்: கணிதம் - மூலக் கருவிகள்

①

பு. பு. மூலக் கருவிகள் பற்றி - தகுந்தவர்

PART - I TYPE - A			PART - II		
1	a	$\frac{8}{3}$	21	$\sum_{n=1}^{12} i^n = i^1 + i^2 + \dots + i^{12}$ $= (i^1 + i^2 + i^3 + i^4) + (i^5 + i^6 + i^7 + i^8) + (i^9 + i^{10} + i^{11} + i^{12})$ $= 0 + 0 + 0 = 0$	
2	d	$\frac{3\pi a^4}{16}$	22	$\alpha + \beta = \frac{7}{2}, \alpha\beta = \frac{13}{2}$ $\alpha^2 + \beta^2 = \frac{-3}{4}, \alpha^2\beta^2 = \frac{169}{4}$ $x^2 + \frac{3}{4}x + \frac{169}{4} = 0$ $4x^2 + 3x + 169 = 0$	
3	a	10	23	$df = (2x+3)dx$ $x = 3 \text{ மீட்டர் } dx = 0.02$ $df = (6+3)(0.02)$ $= 9(0.02)$ $= 0.18$	
4	d	2	24	$y = mx \quad \text{--- (1)}$ $\frac{dy}{dx} = m$ $(1) \Rightarrow y = x \frac{dy}{dx}$ $\Rightarrow \frac{dy}{dx} = \frac{y}{x}$	
5	d	8	25	$E(x) = \int_1^2 f(x) dx = \int_1^2 2(x-1) dx$ $= 2 \left[\frac{x^2}{2} - x \right]_1^2$ $E(x) = 1$ $f(x) = 1$	
6	a	$[\vec{\alpha}, \vec{\beta}, \vec{\gamma}] = 0$	26	$(h, k) = (-3, -4), r = 3$ $(x-h)^2 + (y-k)^2 = r^2$ $(x+3)^2 + (y+4)^2 = 3^2$ $x^2 + y^2 + 6x + 8y + 14 = 0$	
7	d	$\frac{d^2y}{dn^2} - y = 0$			
8	b	$y = 0$			
9	d	$\frac{1}{(1+x)^2} dx$			
10	a	$x^2 + y^2$			
11	c	2			
12	d	புரவணமயம் / parabola			
13	c	$\begin{bmatrix} 5 & -2 \\ 3 & -1 \end{bmatrix}$			
14	a	45'			
15	d	$\frac{1}{\sqrt{5}}$			
16	b	$-\frac{2}{r}$			
17	a	0			
18	c	2			
19	d	$\text{adj}(AB) = (\text{adj}A)(\text{adj}B)$			
20	c	-4			

27 $A = \begin{bmatrix} -1 & 3 \\ 4 & -7 \\ 3 & -4 \end{bmatrix}$ $\rho(A) \leq \{\min 3, 2\} = 2$

$$\begin{vmatrix} -1 & 3 \\ 4 & -7 \end{vmatrix} = 7 - 12 = -5 \neq 0$$

எனவே $\rho(A) = 2$

32 $x^2 + 6x + 4y + 5 = 0$
 $(x+3)^2 = -4(y-1)$
 $x^2 = -4y$

மேலே உள்ள 4யின் சமன்பாடு
 $2x + y + 1 = 0$

கீழே உள்ள 4யின் சமன்பாடு
 $x - 2y - 7 = 0$

28 $\int_0^{\frac{\pi}{2}} \sin^{10} x dx = \frac{9}{10} \times \frac{7}{8} \times \frac{5}{6} \times \frac{3}{4} \times \frac{1}{2} \times \frac{\pi}{2}$

$$= \frac{63\pi}{512}$$

33 LHS = $[\vec{a} - \vec{b}, \vec{b} - \vec{c}, \vec{c} - \vec{a}]$
 $= (\vec{a} - \vec{b}) \cdot ((\vec{b} - \vec{c}) \times (\vec{c} - \vec{a}))$
 $= [\vec{a} \vec{b} \vec{c}] - 0 + 0 - 0 + 0 - [\vec{b} \vec{c} \vec{a}]$
 $= 0 = \text{RHS}$

29 $\lim_{x \rightarrow 1} \frac{x^2 - 3x + 2}{x^2 - 4x + 3}$
 $= \lim_{x \rightarrow 1} \left(\frac{2x - 3}{2x - 4} \right)$
 $= \frac{2 - 3}{2 - 4} = \frac{-1}{-2} = \frac{1}{2}$


34 $u(x, y) = \frac{x^2 + y^2}{\sqrt{x+y}}$
 $u(\lambda x, \lambda y) = \frac{\lambda^2 x^2 + \lambda^2 y^2}{\sqrt{\lambda} \sqrt{x+y}}$
 $= \lambda^{2 - \frac{1}{2}} u(x, y)$
 $= \lambda^{\frac{3}{2}} u(x, y)$

$u(x, y)$ -ன் மதிப்பு $\frac{3}{2}$ உள்ளது என்பதைக் காட்டுவோம்.
 சரிசெய்தல்: $x \frac{\partial u}{\partial x} + y \frac{\partial u}{\partial y} = \frac{3}{2} u$

30 $[\vec{a} \vec{b} \vec{c}] = \begin{vmatrix} 2 & -1 & 3 \\ 1 & -1 & 0 \\ 3 & -1 & 6 \end{vmatrix}$
 $= 2(-6-0) + 1(-6-18) + 3(-1+3)$
 $= -12 + 6 + 6 = 0$

எனவே இவற்றின் மதிப்பு 0.

PART - III



31 $\cos^{-1} \left(\frac{1}{\sqrt{x^2-1}} \right) = \alpha$ எனில்

$$\cos \alpha = \frac{1}{\sqrt{x^2-1}}$$

$$\sec \alpha = \frac{x}{1} = x$$

$$\alpha = \sec^{-1} x$$

$$\cos^{-1} \left(\frac{1}{\sqrt{x^2-1}} \right) = \sec^{-1} x$$

35 $x + y = 12$ — (1)
 $y = 12 - x$
 $P = xy = x(12 - x) = 12x - x^2$
 $\frac{dP}{dx} = 12 - 2x$
 $\frac{dP}{dx} = 0 \Rightarrow 12 - 2x = 0$
 $x = 6$

(1) $\Rightarrow y = 12 - 6$
 $y = 6$

எனவே இவற்றின் மதிப்பு $x = 6, y = 6$

36 $I = \int_{\frac{\pi}{8}}^{\frac{3\pi}{8}} \frac{1}{1 + \sqrt{\tan x}} dx$ — ①

$I = \int_{\frac{\pi}{8}}^{\frac{3\pi}{8}} \frac{\sqrt{\tan x}}{1 + \sqrt{\tan x}} dx$ — ②

Adding ①, ②

$$I + I = \int_{\frac{\pi}{8}}^{\frac{3\pi}{8}} \frac{1 + \sqrt{\tan x}}{1 + \sqrt{\tan x}} dx$$

$$2I = \int_{\frac{\pi}{8}}^{\frac{3\pi}{8}} 1 dx$$

$$= \left[x \right]_{\frac{\pi}{8}}^{\frac{3\pi}{8}}$$

$$2I = \frac{\pi}{4}$$

$$I = \frac{\pi}{8}$$

37 $\frac{1+i}{1-i} = i, \frac{1-i}{1+i} = -i$

$$\left(\frac{1+i}{1-i}\right)^3 - \left(\frac{1-i}{1+i}\right)^3 = i^3 - (-i)^3$$

$$= -i - (-i)$$

$$= -2i$$

38 $\frac{dy}{1+y^2} = \frac{dx}{1+x^2}$

$$\int \frac{1}{1+y^2} dy = \int \frac{1}{1+x^2} dx$$

$$\tan^{-1} y = \tan^{-1} x + C$$

$$\tan^{-1} \left(\frac{y-x}{1+xy}\right) = C$$

$$\frac{y-x}{1+xy} = \tan C = C$$

39

x	0	1	2	3
$f(x)$	$\frac{1}{8}$	$\frac{3}{8}$	$\frac{3}{8}$	$\frac{1}{8}$

$$f(x) = \begin{cases} \frac{1}{8}, & x = 0, 3 \\ \frac{3}{8}, & x = 1, 2 \end{cases}$$

40 $|\text{adj}(\text{adj} A)| = |A|^{(n-1)^2}$

$$\begin{vmatrix} 2 & -1 & 3 \\ -5 & 3 & 1 \\ -3 & 2 & 3 \end{vmatrix} = -1$$

$$|\text{adj}(\text{adj} A)| = (-1)^4 = 1$$

PART - IV

41 (a) given the point $\left(\frac{3}{2}, \frac{9}{4}\right)$

$$y = x^2$$

$$\frac{dy}{dx} = 2x, m_1 = 3$$

$$y = (x-3)^2$$

$$\frac{dy}{dx} = 2(x-3), m_2 = -3$$

$$\tan \theta = \left| \frac{3 - (-3)}{1 - 9} \right| = \frac{3}{4}$$

$$\theta = \tan^{-1} \left(\frac{3}{4}\right)$$

b $\tan^{-1} \left(\frac{x-1}{x-2}\right) + \tan^{-1} \left(\frac{x+1}{x+2}\right) = \frac{\pi}{4}$

$$\frac{\frac{x-1}{x-2} + \frac{x+1}{x+2}}{1 - \left(\frac{x-1}{x-2}\right)\left(\frac{x+1}{x+2}\right)} = 1$$

$$2x^2 - 4 = -3$$

$$x^2 = \frac{1}{2}$$

$$x = \pm \frac{1}{\sqrt{2}}$$

42

சொகுதியின் தொகை	2	3	4	5	6
பெரிசொகுதி	1	4	10	12	9

இதில் தரப்பட்ட தகவல்களைப் பற்றி

x	2	3	4	5	6
f(x)	1/36	4/36	10/36	12/36	9/36


இதில் தரப்பட்ட தகவல்களைப் பற்றி

$$f(x) = \begin{cases} 0 & ; -\infty < x < 2 \\ \frac{1}{36} & ; 2 \leq x < 3 \\ \frac{5}{36} & ; 3 \leq x < 4 \\ \frac{15}{36} & ; 4 \leq x < 5 \\ \frac{27}{36} & ; 5 \leq x < 6 \\ 1 & ; 6 \leq x < \infty \end{cases}$$

$$\frac{dv}{dt} = \frac{25\pi}{144} h^2 \frac{dh}{dt}$$

$$\frac{dh}{dt} = \frac{10 \times 144}{25\pi \times 64} = \frac{9}{10\pi} \text{ cm/sec}$$

b $\sin(\alpha - \beta) = \sin\alpha \cos\beta - \cos\alpha \sin\beta$



$$\hat{a} = \cos\alpha \hat{i} + \sin\alpha \hat{j}$$

$$\hat{b} = \cos\beta \hat{i} + \sin\beta \hat{j}$$

$$\hat{b} \times \hat{a} = (\sin\alpha \cos\beta - \cos\alpha \sin\beta) \hat{k}$$

$$\hat{b} \times \hat{a} = |\hat{b}| |\hat{a}| \sin(\alpha - \beta) \hat{k} = \sin(\alpha - \beta) \hat{k}$$

(1), (2)

$$\sin(\alpha - \beta) = \sin\alpha \cos\beta - \cos\alpha \sin\beta$$


b $\text{Im} \left(\frac{(2x+1)+2iy}{(1-y)+ix} \times \frac{(1-y)-ix}{(1-y)-ix} \right) = 0$

$$\frac{(2x+1)(-x) + 2y(1-y)}{(1-y)^2 + x^2} = 0$$

$$-2x^2 - x + 2y - 2y^2 = 0$$

$$2x^2 + 2y^2 + x - 2y = 0$$

44
(a)



$$x^2 = -4ay \quad \text{--- (1)}$$

$$(3, -2.5) \Rightarrow 3^2 = -4a(-2.5)$$

$$4a = \frac{9}{2.5}$$

$$x^2 = -\frac{9}{25} y \quad \text{--- (2)}$$


43
(a)

$$12r = 5h$$

$$r = \frac{5h}{12}$$

$$V = \frac{1}{3} \pi r^2 h$$

$$V = \frac{25\pi h^3}{3 \times 144}$$




$(x, -75) \Rightarrow$

$x_1 = 9(3)$

$x_2 = 3\sqrt{3}$

கீழ்க்கண்டிருக்கிற $3\sqrt{3}$ ல்
 கீழ்க்கண்டிருக்கிற $9(3)$ இரண்டின்
 மூலம் கிடைக்கிறது.

46
 (a)



$$A = 4 \int_0^a \frac{b}{a} \sqrt{a^2 - x^2} dx$$

$$= \pi ab$$

(b) $P = \frac{1}{x}, Q = \sin x$

$P = \log x, I \cdot F = x$

$y e^{\int P dx} = \int Q e^{\int P dx} dx + C$

$xy = \int \sin x \cdot x dx + C$

$xy = -x \cos x + \sin x + C$

$xy + x \cos x = \sin x + C$


(b)

முனை: (1, 2)

மேலே: (3, 2)

கீழே: $x = -1$

மேல்க்கண்டிருக்கிற கிடைக்கிறது = 8



45 $\int \frac{dA}{A} = A \int dt$

(a) $A = ce^{kt}$ — ①

$t=0$ க்கான $A = A_0$

$A = A_0 e^{kt}$ — ②

$t=5$ க்கான $A = 3A_0$

$e^{5k} = 3$

$t=10$ க்கான

$A = 9A_0$

47

P Q P ↔ Q

(a)

T T T

T F F

F T F

F F T

$((\sim P) \vee Q) \wedge ((\sim Q) \vee P) = TFFT$

b $\vec{r} = a\vec{i} + s\vec{j} + t\vec{k}$
 $\vec{r} = (2\hat{i} + 2\hat{j} + \hat{k}) + s(2\hat{i} + 3\hat{j} + 3\hat{k})$
 $+ t(3\hat{i} + 2\hat{j} + \hat{k})$

$$\begin{vmatrix} x-2 & y-2 & z-1 \\ 2 & 3 & 3 \\ 3 & 2 & 1 \end{vmatrix} = 0$$

$3x - 7y + 5z + 3 = 0$

(b)

$A = -15, A_1 = -15,$

$A_2 = -5, A_3 = -5$

$x = 1, y = 3, z = 3$

V. GURUNATHAN
 P. G. ASST. - Maths
 M. M. Hr. S. School
 Thirunagar
 Madurai - 6.