Marking Scheme 2016-17 Science Class - X (SA - II)

1 Hydrogenation 1 2 Moustaches, Beard, Hoarse voice, Thick growth of hair on the body (Any 1 two) 3 To prevent u-v radiation from reaching the earth. 1 f= -10cm (since the image is formed at the focus) 4 1/2 Now u= - 20cm, i,e. the candle flame is at C :. The image would form at C and would be of the same size 1/2 1 Ray diagram (Refer Fig. 10.7(c) Page 166 NCERT Text-book) 5 Conclusion: Water is polluted by sewage 1 Coliform is a group of bacteria found in human intestine whose 1 presence in river water indicates contamination by disease causing micro-organisms. 6 **Forests** 1/2 Range of different life forms plays an important role in maintaining 11/2 ecological balance as they all depend on each other for their survival. 7 Isomers: Compounds which have same molecular formula and different structural formula. 1 In first three members of alkane series, branching is not possible, therefore isomers are not possible. 1 Two isomers of butane C₄H₁₀ 1/2+1/2 1/2 8 X- Ethanol C₂H₅OH 1/2 Y- Ethanoic acid CH₃COOH Z- Ester CH₃COOC₂H₅ 1/2 · CH3CH2OH Alkaline KMm04+Heat CH3COOH or Acidified K2Cr2O2+ Heat (Y) 1 CH3COOH + CH3CH2OH Acid > CH3COOC2H5 1/2

Atomic number of A - 17

9

1/2

	 Electronic configuration of A = (2, 8,7) A is a non-metal. B →2,8,1 	½ ½
	B will give one electron to A to form an ionic compound- Ionic bond • Formula of the compound B ⁺ A ⁻	1 ½
10	(a) K, Electronic configuration of K is 2,8,8,1 (b) Be and Ca Be (4) = 2,2	1
	Ca $(20) = 2,8,8,2$ • $K^{+}X^{-}K = 2,8,8,1$ X = 2,8,7	1
	K will give one electron to X to form ionic bond. So the compound formed $K^{\scriptscriptstyle +}X^{\scriptscriptstyle -}$ will be an ionic compound.	1
11	 This process is called regeneration. Diagram (Refer Fig 8.3 Page 131 NCERT Text-book) When a Planaria is cut into three pieces, each of its pieces grow into separate individual; specialized cells proliferate; mass of cells 	½ 1
	 get differentiated into various cells and tissues. Another example of an organism showing the characteristics is Hydra 	1 ½
12	 Four methods of contraception (i) Mechanical/ Barrier method (ii) Use of hormonal preparations (iii) Use of loop/Copper-T/ IUCD (iv) Surgical method (Tubectomy/ Vasectomy) Effect on health and prosperity (Any two) Health of women is maintained 	4 x ½ 2 x ½
	Better attention of children by the parentsMore resources for better living standard	2 X / 2
13	 Two functions each of Ovaries: (i) Production of female sex hormone. (ii) Production of female gamete. 	2 x ½
	 Fallopian tube (i) Transfer of female gamete from ovary (ii) Site of fertilization. 	2 x ½
	 Uterus (i) Implantation of zygote (ii) Nourishment of developing embryo. 	2 x ½
14	 In some animals, the temperature at which fertilized eggs are kept determines whether the developing animal in egg is male or female In some animals like snail, individual can change sex. 	1½
	Genetical Cue - A child who inherits an x chromosome from her father will be a girl and one who inherits a y chromosome from the father will be a boy.	1½

Marking Scheme – 2 Pg2 of 6

		Acquired Traits	Inherited Traits	
	•	Does not bring about change in	Bring about change in the DNA of	
		the DNA of germ cells	the germ cells	
	•	Cannot be passed on to the	Can be passed on to the progeny	
		progeny Cannot direct evolution	Can direct evolution	
			(Any two)	2 x 1
	•	Examples: Acquiring knowledge/	Skin colour/ color of the eyes	
		less of body parts.		1
	_			
•	C	onverging lens (Convex lens)		1/2
	н	u+v=6m		
•	• • •	v = 3u		
		u + 3u = 6m		
	_			1
	_	$\Rightarrow u = 15m$		•
<u> </u>	_ 1_	$-\frac{1}{u} \Rightarrow f = \frac{uv}{u-v}$		1/
ŗ	- v	$u \xrightarrow{\longrightarrow} J - u - v$		1/2
		(1.5m)(1.5m)		
. j	^c = -	$\frac{(-1.5m)(4.5m)}{-1.5m - (4.5m)} = +1.125m$		1
	-	-1.5m - (4.5m)		
	a)	Hypermetropia		1/2
	•	Two causes: (i) The focal length	of the eye lens is too long	2 x ½
	D)	· · ·	•	2 X /2
	- \	(ii) The eye ball has		1/
	,	Ray diagram (Refer fig 11.3 (c) p	•	1/2
	` '	Importance of green plants, re	5 1 /	1 ½
		noise levels, aesthetic value, sa		
		plants in place of bouquet, creat		4 1/
	(D)	To protect and improve the		1 ½
		awareness, environmental conse	ervation, eco-mendiy approach	
		value for life (any three) Two reasons for large number of	carbon compounds	
	(a)	(i) Catenation: Unique ability		
		•	rise to long chains of different	
		types of compounds	The to tong chains of afficient	1
		(ii) Tetravalency: Since carbon h	has a valency of 4, it is capable	
			toms of carbon or atoms of the	
			Oxygen, Nitrogen, Hydrogen,	1
		Sulphur, chlorine etc.	,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,	
	(b)	,		
	. ,	(i) Carbon has 4 elements as its	outmost shell and needs to gain	
		• •	noble gas configuration. Losing	
		or gaming 4 electrons	is not possible for energy	1
			hares electrons to form the	ı
		covalent bands		
		(ii) The forces of attraction betw	-	
			compounds generally have low	1
		malting and bailing points		

melting and boiling points.

(c) (i) $2Na + 2CH_3CH_2OH \rightarrow 2CH_3CH_2ONa + H_2$ 1/2 1/2 (ii) $NaOH + CH_3COOH \rightarrow CH_3COONa + H_2O$ 20 A. Diagram: 1 Lebellings: 4x½ B. A - Pollen grain B - Pollen tube $4x\frac{1}{2}$ C - Ovary D - Female gamete 21 A. Speciation may take place by 1 (i) Migration (ii) Natural selection (iii) Mutation (iv) Genetic Drift (Any two) B. Segment of DNA which is functional and are made of nucleic acids and protein 1 (Any other definition) C. Given Red hair - Mother - Recessive : bb 1/2 Black hair - father Dominant : BB 1/2 Father 1 Mother **Parents** BB/Bb Χ Bb/bb F1 Bb 1 (Black) Thus, the child will have black hair 22 a) The water droplets act like small prisms. They refract and disperse the incident sunlight, then reflect it internally and finally refract it again when it comes out of rain drop. Due to 1 dispersion and internal reflection of light, different colours reach the observer's eye along different pairs.

light is comparatively less that results in lesser scattering. Thus sun looks white.

dispersion, internal reflection of light.

Therefore the three phenomena involved are refraction,

light. This makes sun orange. At noon, the distance travelled by

b) At sunrise / sunset, sunlight has to travel a larger distance. So, it come across more number of particles which scatters most of blue colour and the light reaching our eyes has more of reddish

11/2


1/2

2


- 23 (i) When an object is placed between the pole and focus of concave mirror a magnified, erect and virtual image is obtained.
- 1

1

(ii) For glass slab refer:



For prism refer:



1

In case of a rectangular glass slab, emergent rays of light are always parallel to the direction of incident rays. Whereas when an incident light passes through a prism, it bends towards the base of the prism hence incident ray and emergent ray are not parallel to each other.

1

(iii)
$$f = -50cm$$
 $p = \frac{100}{f}D = \frac{100}{-50} = -2D$


1

24 (i) No, magnified image of an object cannot be formed by a concave lens ever.

1

(ii) At 2f.

1



1

(iii)




Image obtained is virtual, erect and diminished in case of concave lens

1

		1
25	(a)	1
26	(a)	1
27	2	1
28	(d)	1
29	(c)	1
30	(d)	1
31	(b)	1
32	(b)	1
33	(d) 1 - plumule, 2- radicle	1
34	 Carbon dioxide/CO₂ line water terms milky when CO2 gas is passed through it/ the gas liberated extinguishes a boring splinter 	1 1
35	Turnip, Carrot, Sweet Potato - are modified root - Homologous organs Potato - modified stem.	2
36	He should move the lens towards the screen. As the distance of object increased, the image formed by a convex lens will be more close to the focus.	2

Marking Scheme – 2 Pg6 of 6