











### MARKING SCHEME

1	No $\alpha$ H is present	1
2	Ethanol will be converted into ethanoic acid.	1
3	$[\text{Cr}(\text{H}_2\text{O})_4\text{Cl}_2]\text{Cl}$ Tetraaquadichloridochromium(III) chloride	$\frac{1}{2} + \frac{1}{2}$
4	The Brownian movement has a stirring effect, which does not allow the particles to settle.	1
5	$e^{-E_a/RT}$ Corresponds to the fraction of molecules that have kinetic energy greater than $E_a$ .	1
6	(i) Vinyl chloride does not respond to NaOH and silver nitrate test because of partial double bond character due to resonance. (ii) Hydride ion / $\text{H}^-$	1 1
7	0.05 M $\text{Al}_2(\text{SO}_4)_3$ has higher freezing point. 0.05 M $\text{Al}_2(\text{SO}_4)_3$ : $i = 5$ , $\Delta T_f \propto$ No of particles ; $\Delta T_f = i \times$ concentration = $5 \times 0.05 = 0.25$ moles of ions 0.1 M $\text{K}_3[\text{Fe}(\text{CN})_6]$ : $i = 4$ , = $4 \times 0.1 = 0.4$ moles of ions	1 $\frac{1}{2}$ $\frac{1}{2}$
8	$2\text{Cr}(s) + 3\text{Fe}^{2+}(\text{aq.}) \rightarrow 3\text{Fe}(s) + 2\text{Cr}^{3+}(\text{aq.})$ $n = 6$ $E_{\text{Cell}} = E_{\text{Cell}}^0 - \frac{2.303RT}{nF} \log \frac{[\text{Cr}^{3+}]^2}{[\text{Fe}^{2+}]^3}$ $E_{\text{Cell}} = 0.30 - \frac{0.059}{6} \log \frac{[10^{-1}]^2}{[10^{-2}]^3}$ $E_{\text{Cell}} = 0.26 \text{ V}$ <p style="text-align: center;">OR</p> $\wedge_m = \frac{1000\kappa}{C}$ $\wedge_m = \frac{1000 \times 4.1 \times 10^{-5}}{10^{-3}} = 41 \text{ S cm}^2 \text{ mol}^{-1}$ $\alpha = \frac{\wedge_m^c}{\wedge_m^0}$ $\alpha = \frac{41}{390.5} = 0.105$	$\frac{1}{2}$ $\frac{1}{2}$ $\frac{1}{2}$ $\frac{1}{2}$ $\frac{1}{2}$ $\frac{1}{2}$
9	(i) Orthophosphorus acid on heating disproportionates to give orthophosphoric acid and phosphine gas.	1

	$4H_3PO_3 \xrightarrow{\text{heat}} PH_3 + 3H_3PO_4$ <p>(ii) When <math>XeF_6</math> undergoes complete hydrolysis, it forms <math>XeO_3</math>.  <math>XeF_6 + 3H_2O \rightarrow 6HF + XeO_3</math></p>	1
10	<p>(i) <math>Cr_2O_7^{2-}</math></p> <p>(ii) Cerium</p>	1 1
11	<p>(i) 2,5-Dimethylhexane.</p> <p>(ii) 1-Methyl-1-iodocyclohexane.</p> <p>(iii) Nitroethane.</p>	1+1+1
12	$\Delta T_f = i K_f m$ $2.12 = i \frac{5.12 \times 2.5 \times 1000}{122 \times 25}$ <p><math>i = 0.505</math> for association</p> $i = 1 - \frac{\alpha}{2}$ <p><math>\alpha = 0.99</math> Percentage association of benzoic acid is 99.0%</p>	$\frac{1}{2}$ 1 $\frac{1}{2}$ $\frac{1}{2}$ $\frac{1}{2}$
13	<p>(i) Because of H-bond formation between alcohol and water molecule.</p> <p>(ii) Nitro being the electron withdrawing group stabilises the phenoxide ion.</p> <p>(iii) side product formed in this reaction is acetone which is another important organic compound.</p>	1+1+1
14	$t = \frac{2.303}{k} \log \frac{[R]_0}{[R]}$ $t = \frac{2.303}{60} \log \frac{1}{0.0625}$ <p><math>t = 0.0462 \text{ s}</math></p>	1 1 1
15	<p>(i) 'B' is a strong electrolyte. A strong electrolyte is already dissociated into ions, but on dilution interionic forces are overcome, ions are free to move. So there is slight increase in molar conductivity on dilution.</p> <p>(ii) On anode water should get oxidised in preference to <math>Cl^-</math>, but due to overvoltage/ overpotential <math>Cl^-</math> is oxidised in preference to water.</p>	1 1 1
16	<p>(i) <math>\frac{x}{m} = kC^{1/n}</math></p> <p>(ii) The charge on the sol particles is due to</p> <ul style="list-style-type: none"> <li>• Electron capture by sol particles during electrodispersion.</li> <li>• Preferential adsorption of ions from solution.</li> <li>• Formulation of electrical double layer. (any one reason)</li> </ul> <p>(iii) Molybdenum acts as a promoter for iron.</p>	1 1 1

17	A		½ each
B			
C			
D			
E			
F			
18	(i) Vitamin D. (ii) Uracil. (iii) 5 OH groups are present.		1 1 1
19	(i) Addition (ii) Condensation/Hydrolysis (iii) Condensation		1 1 1
20	(i) Gold is leached with a dilute solution of NaCN in the presence of air (ii) Cryolite lowers the high melting point of alumina and makes it a good conductor of electricity. (iii) CO forms a volatile complex with metal Nickel which is further decomposed to give pure Ni metal.		1 1 1



25	(i)		1
	(ii)		$\frac{1}{2} + \frac{1}{2}$
	(iii)	Cl-CH <sub>2</sub> -COOH	1
	B(I)	NaHCO <sub>3</sub> test.	1
	(ii)	Iodoform test./Fehling's Test/ Tollen's Test	1
	<b>OR</b>		
	<b>A (i)</b>	steric and electronic factor.	
	(ii)	Inductive effect decreases with distance and hence the conjugate base of 2-Fluorobutanoic acid is more stable.	$\frac{1}{2} + \frac{1}{2}$
	b)		
	i)		
			1
	(ii)		1
	(c)		1

	$\text{HCN} + \text{OH}^- \rightleftharpoons \text{:}\bar{\text{C}}\text{N} + \text{H}_2\text{O}$ <p style="text-align: center;">Tetrahedral intermediate</p> <p style="text-align: center;">Cyanohydrin</p>	
26	<p>(i) Ferrimagnetism. These substances lose ferrimagnetism on heating and become paramagnetic.</p> <p>(ii) <math>r = 0.414 R</math></p> <p>(iii) <math>r = \frac{\sqrt{3}}{4} a</math> <math>r = \frac{\sqrt{3}}{4} \times 316.5</math> <math>r = 136.88 \text{ pm}</math></p> <p style="text-align: center;">OR</p> <p>(i) Schottky defect It is shown by ionic substances in which the cation and anion are of almost similar sizes.</p> <p>(ii) <math>r = \frac{\sqrt{3}}{4} a</math></p> <p>(iii) <math>\rho = \frac{z M}{a^3 N_A}</math></p> $8.92 = \frac{z \times 63}{(3.608 \times 10^{-8})^3 \times 6.022 \times 10^{23}}$ <p style="text-align: center;"><math>z = 4</math> So it is face centred cubic lattice</p>	<p>1 1 1 1 <math>\frac{1}{2}</math> <math>\frac{1}{2}</math></p> <p>1 1 1 <math>\frac{1}{2}</math> 1 <math>\frac{1}{2}</math></p>

**CBSE SAMPLE PAPER CHEMISTRY-2017-18**

**MM: 70**

**BLUE PRINT**

**TIME 3 HRS**

No	CHAPTER	VSA	SA-1	SA-11	VBQ	LA	TOTAL
1	SOLID STATE					1(5) (U)	9(23)
2	SOLUTIONS		1(2) (U)	1(3) (A)			
3	ELECTROCHEMISTRY		1(2) (A)	1(3) (U)			
4	CHEMICAL KINETICS	1(1) (R)		1(3) (A)			
5	SURFACE CHEMISTRY	1(1) (R)		1(3) (R)			
6	EXTRACTION OF METALS			1(3) (U)			7(19)
7	p-BLOCK		1(2) (U)			1(5) (A)	
8	d AND f BLOCK ELEMENTS		1(2) (R)	1(3) (E&MD)			
9	COORDINATION CHEMISTRY	1(1) Hots		1(3) Hots			
10	HALOALKANES AND HALOARENES		1(2) (A)	1(3) (A)			10(28)
11	ALCOHOLS, PHENOLS AND ETHERS	1(1) (E&MD)		1(3) (U)			
12	ALDEHYDES, KETONES AND CARBOXYLIC ACID	1(1)Hots				1(5) (E&MD)	
13	ORGANIC COMPOUNDS COTAINING NITROGEN			1(3) (A)			
14	BIOMOLECULES			1(3) (U)			
15	POLYMERS			1(3) (E&MD)			
16	CHEMISTRY IN EVERY DAY LIFE				1(4) (E&MD)		
	Total						26(70)

R-Recall; U-Understanding; A-Application, Hots- Higher Order Thinking Skills-;  
E&MD-Evaluation and multidisciplinary