

QM

QUARTERLY EXAM - SEPTEMBER - 2024 (Madurai)

12 - STD

Mathematics

(23/09/24)

Time : 3.00 Hours

Marks : 90

PART - I

Note : i) All questions are compulsory

20 × 1 = 20

ii) Choose the most appropriate answer from the given four alternatives and write the option code and the corresponding answer

1. If $P = \begin{bmatrix} 1 & x & 0 \\ 1 & 3 & 0 \\ 2 & 4 & -2 \end{bmatrix}$ is the adjoint of 3×3 matrix A and $|A| = 4$, then x is
 (1) 15 (2) 12 (3) 14 (4) 11
2. If $A = \begin{bmatrix} 3 & 5 \\ 1 & 2 \end{bmatrix}$, $B = \text{adj } A$ and $C = 3A$, then $\frac{|\text{adj } B|}{|C|} =$
 (1) $\frac{1}{3}$ (2) $\frac{1}{9}$ (3) $\frac{1}{4}$ (4) 1
3. If $\text{adj } A = \begin{bmatrix} 2 & 3 \\ 4 & -1 \end{bmatrix}$ and $\text{adj } B = \begin{bmatrix} 1 & -2 \\ -3 & 1 \end{bmatrix}$ then $\text{adj } (AB)$ is
 (1) $\begin{bmatrix} -7 & -1 \\ 7 & -9 \end{bmatrix}$ (2) $\begin{bmatrix} -6 & 5 \\ -2 & -10 \end{bmatrix}$ (3) $\begin{bmatrix} -7 & 7 \\ -1 & -9 \end{bmatrix}$ (4) $\begin{bmatrix} -6 & -2 \\ 5 & -10 \end{bmatrix}$
4. $i^n + i^{n+1} + i^{n+2} + i^{n+3}$ is (1) 0 (2) 1 (3) -1 (4) i
5. If $\left|z - \frac{3}{z}\right| = 2$, then the least value of $|z|$ is
 (1) 1 (2) 2 (3) 3 (4) 5
6. The principal argument of $(\sin 40^\circ + i \cos 40^\circ)^5$ is
 (1) -110° (2) -70° (3) 70° (4) 110°
7. If f and g are polynomials of degrees m and n respectively, and if $h(x) = (f \circ g)(x)$, then the degree of h is
 (1) mn (2) $m+n$ (3) m^n (4) n^m
8. The number of real numbers in $[0, 2\pi]$ satisfying $\sin^4 x - 2\sin^2 x + 1 = 0$ is
 (1) 2 (2) 4 (3) 1 (4) ∞
9. If $p + \sqrt{q}$ and $-i\sqrt{q}$ are the roots of a polynomial equation with rational Coefficients then the least possible degree of the equation is
 1) 2 2) 1 3) 3 4) 4 QM 12 MAT-EM P-1

10. If $\sin^{-1} x + \sin^{-1} y = \frac{2\pi}{3}$ then $\cos^{-1} x + \cos^{-1} y$ is equal to
 (1) $\frac{2\pi}{3}$ (2) $\frac{\pi}{3}$ (3) $\frac{\pi}{6}$ (4) π
11. $\cos(\cos^{-1} x) = x$ if
 1) $|x| < 1$ 2) $|x| \leq 1$ 3) $|x| \geq 1$ 4) $|x| = 0$
12. If the function $f(x) = \sin^{-1}(x^2 - 3)$, then x belongs to
 (1) $[-1, 1]$ (2) $[\sqrt{2}, 2]$ (3) $[-2, -\sqrt{2}] \cup [\sqrt{2}, 2]$ (4) $[-2, -\sqrt{2}]$
13. The radius of the circle $3x^2 + by^2 + 4bx - 6by + b^2 = 0$ is
 1) 1 2) 3 3) $\sqrt{10}$ 4) $\sqrt{11}$
14. If $x + y = k$ is a normal to the parabola $y^2 = 12x$, then the value of k is
 1) 3 2) -1 3) 1 4) 9
15. $y = mx + c$ is a tangent to the parabola $y^2 = 4ax$ then
 1) $c = \frac{a}{m}$ 2) $c = \frac{m}{a}$ 3) $c^2 = a^2 m^2 + m^2$ 4) $m = c$
16. The eccentricity of the ellipse $(x - 3)^2 + (y - 4)^2 = \frac{y^2}{9}$ is
 1) $\frac{\sqrt{3}}{2}$ 2) $\frac{1}{3}$ 3) $\frac{1}{3\sqrt{2}}$ 4) $\frac{1}{\sqrt{3}}$
17. If $\vec{a} \cdot \vec{b} = \vec{b} \cdot \vec{c} = \vec{c} \cdot \vec{a} = 0$, then the value of $[\vec{a}, \vec{b}, \vec{c}]$ is
 1) $|\vec{a}||\vec{b}||\vec{c}|$ 2) $\frac{1}{|\vec{a}||\vec{b}||\vec{c}|}$ 3) 1 4) -1
18. Which one is meaningful?
 1) $(\vec{a} \times \vec{b}) \times (\vec{b} \cdot \vec{c})$ 2) $\vec{a} \times (5 + \vec{b})$ 3) $(\vec{a} \cdot \vec{b}) \times (\vec{c} \cdot \vec{d})$ 4) $(\vec{a} \times \vec{b}) \cdot (\vec{c} \times \vec{d})$
19. If $\vec{a} = \hat{i} + \hat{j} + \hat{k}$, $\vec{b} = \hat{i} + \hat{j}$, $\vec{c} = \hat{i}$ and $(\vec{a} \times \vec{b}) \times \vec{c} = \lambda \vec{a} + \mu \vec{b}$, then the value of $\lambda + \mu$ is
 1) 0 2) 1 3) 6 4) 3
20. The distance between the planes $x + 2y + 3z + 7 = 0$ and $2x + 4y + 6z + 7 = 0$ is
 (1) $\frac{\sqrt{7}}{2\sqrt{2}}$ (2) $\frac{7}{2}$ (3) $\frac{\sqrt{7}}{2}$ (4) $\frac{7}{2\sqrt{2}}$

PART - II

- Note : i) Answer any Seven questions
 ii) Question number 30 is compulsory.

7 × 2 = 14

21. If $\text{adj}(A) = \begin{bmatrix} 0 & -2 & 0 \\ 6 & 2 & -6 \\ -3 & 0 & 6 \end{bmatrix}$, find A^{-1} .

22. Find the square root of $6 - 8i$.

23. If α, β, γ are the roots of the equation $x^3 + px^2 + qx + r = 0$, find the value of $\sum \frac{1}{\beta\gamma}$ in terms of the coefficients.

QM 12 MAT-EM P-2

24. Find the principal value of $\tan^{-1}(\sqrt{3})$.
25. Find the general equation of the circle whose diameter is the line segment joining the points $(-4, -2)$ and $(1, 1)$.
26. If $2\hat{i} - \hat{j} + 3\hat{k}$, $3\hat{i} + 2\hat{j} + \hat{k}$, $\hat{i} + m\hat{j} + 4\hat{k}$ are coplanar, find the value of m .
27. Find the product $\frac{3}{2} \left(\cos \frac{\pi}{3} + i \sin \frac{\pi}{3} \right) \cdot 6 \left(\cos \frac{5\pi}{6} + i \sin \frac{5\pi}{6} \right)$ in rectangular form.
28. Solve the following systems of linear equations by Cramer's rule:
 $5x - 2y + 16 = 0$, $x + 3y - 7 = 0$
29. Find the coordinates of the point where the straight line $\vec{r} = (2\hat{i} - \hat{j} + 2\hat{k}) + t(3\hat{i} + 4\hat{j} + 2\hat{k})$ intersects the plane $x - y + z - 5 = 0$.
30. The parabolic communications antenna has a focus at $4m$ distance from the vertex of the antenna. Write an equation of the parabolic arch.

Part - III

Note : i) Answer any Seven questions

7 × 3 = 21

ii) Question number 40 is compulsory.

31. Solve the following system of linear equations, using matrix inversion method:
 $5x + 2y = 3$, $3x + 2y = 5$. **MADURAI DISTRICT**
32. Write in polar form of the following complex numbers : $2 + i2\sqrt{3}$
33. Show that the equation $x^9 - 5x^5 + 4x^4 + 2x^2 + 1 = 0$ has atleast 6 imaginary solutions.
34. Find the value of $\tan^{-1}(-1) + \cos^{-1}\left(\frac{1}{2}\right) + \sin^{-1}\left(-\frac{1}{2}\right)$.
35. Find the equation of the parabola whose vertex is $(5, -2)$ and focus $(2, -2)$.
36. A particle acted on by constant forces $8\hat{i} + 2\hat{j} - 6\hat{k}$ and $6\hat{i} + 2\hat{j} - 2\hat{k}$ is displaced from the point $(1, 2, 3)$ to the point $(5, 4, 1)$. Find the total work done by the forces.
37. Construct a cubic equation with roots : 1, 2, and 3
38. Find the rank of each of the following matrices: $\begin{bmatrix} 3 & 2 & 5 \\ 1 & 1 & 2 \\ 3 & 3 & 6 \end{bmatrix}$.
39. Find the angle between the straight lines $\frac{x-4}{2} = \frac{y}{1} = \frac{z+1}{-2}$ and $\frac{x-1}{4} = \frac{y+1}{-4} = \frac{z-2}{2}$ and state whether they are parallel or perpendicular.
40. Find the value of the real numbers x and y , if the complex numbers $(2 + i)x + (1 - i)y - 5$ and $x + (-1 + 2i)y$ are equal.

QM 12 MAT-EM P-3

PART - IV

7×5 = 35

Note : i) Answer all the questions .

41. a) Prove by vector method that $\sin(\alpha - \beta) = \sin \alpha \cos \beta - \cos \alpha \sin \beta$. (OR)
- b) If $A = \begin{bmatrix} 8 & -6 & 2 \\ -6 & 7 & -4 \\ 2 & -4 & 3 \end{bmatrix}$, verify that $A(\text{adj } A) = (\text{adj } A)A = |A|I_3$.
42. a) If $z = x + iy$ and $\arg\left(\frac{z-1}{z+1}\right) = \frac{\pi}{2}$, show that $x^2 + y^2 = 1$. (OR)
- b) If $\tan^{-1} x + \tan^{-1} y + \tan^{-1} z = \pi$, show that $x + y + z = xyz$.
43. a) Identify the type of conic and find centre, foci, vertices, and directrices of each of the following : $18x^2 + 12y^2 - 144x + 48y + 120 = 0$. (OR)
- b) Find all cube roots of $\sqrt{3} + i$.
44. a) Solve the equation $(x - 2)(x - 7)(x - 3)(x + 2) + 19 = 0$. (OR)
- b) If the normal at the point ' t_1 ' on the parabola $y^2 = 4ax$ meets the parabola again at the point ' t_2 ' then prove that $t_2 = -\left(t_1 + \frac{2}{t_1}\right)$.
45. a) Test for consistency of the following system of linear equations and if possible, solve:
 $4x - 2y + 6z = 8$, $x + y - 3z = -1$, $15x - 3y + 9z = 21$. (OR)
- MADURAI DISTRICT**
- b) If $2 + i$ and $3 - \sqrt{2}$ are roots of the equation $x^6 - 13x^5 + 62x^4 - 126x^3 + 65x^2 + 127x - 140 = 0$ find the all roots .
46. a) Evaluate $\sin \left[\sin^{-1} \left(\frac{3}{5} \right) + \sec^{-1} \left(\frac{5}{4} \right) \right]$. (OR)
- b) Find the parametric, non parametric form of vector and Cartesian form of the equations of the plane passing through the three non-collinear points $(3, 6, -2)$, $(-1, -2, 6)$, and $(6, 4, -2)$.
47. a) If $\vec{a} = 2\hat{i} + 3\hat{j} - \hat{k}$, $\vec{b} = 3\hat{i} + 5\hat{j} + 2\hat{k}$, $\vec{c} = -\hat{i} - 2\hat{j} + 3\hat{k}$, verify that $\vec{a} \times (\vec{b} \times \vec{c}) = (\vec{a} \cdot \vec{c})\vec{b} - (\vec{a} \cdot \vec{b})\vec{c}$. (OR)
- b) On lighting a rocket cracker it gets projected in a parabolic path and reaches a maximum height of 4m when it is 6m away from the point of projection. Finally it reaches the ground 12m away from the starting point. Find the angle of projection.

QM 12 MAT-EM P-4