

# COMMON QUARTERLY EXAMINATION - 2024

## Standard XII

Reg.No.

## MATHEMATICS

Time : 3.00 hrs

Part - I

Marks : 90

I. Choose the correct answer:

20 x 1 = 20

1. If  $A = \begin{bmatrix} 2 & 0 \\ 1 & 5 \end{bmatrix}$  and  $B = \begin{bmatrix} 1 & 4 \\ 2 & 0 \end{bmatrix}$  then  $|\text{adj}(AB)| =$ 
  - a) -40
  - b) -80
  - c) -60
  - d) -20
2. If A, B and C are invertible matrices of some order, then which one of the following is not true?
  - a)  $\text{adj} A = |A| A^{-1}$
  - b)  $\text{adj}(AB) = (\text{adj} A) (\text{adj} B)$
  - c)  $\det A^{-1} = (\det A)^{-1}$
  - d)  $(ABC)^{-1} = C^{-1} B^{-1} A^{-1}$
3. If  $A = \begin{bmatrix} 2 & 3 \\ 5 & -2 \end{bmatrix}$  be such that  $\lambda A^{-1} = A$ , then  $\lambda$  is
  - a) 17
  - b) 14
  - c) 19
  - d) 21
4. The value of  $\sum_{n=1}^{13} (i^n + i^{n-1})$  is
  - a)  $1 + i$
  - b)  $i$
  - c) 1
  - d) 0
5. If  $|z_1| = 1$ ,  $|z_2| = 2$ ,  $|z_3| = 3$  and  $|9z_1z_2 + 4z_1z_3 + z_2z_3| = 12$ , then the value of  $|z_1 + z_2 + z_3|$  is
  - a) 1
  - b) 2
  - c) 3
  - d) 4
6. The principal argument of  $(\sin 40^\circ + i \cos 40^\circ)^5$  is
  - a)  $-110^\circ$
  - b)  $-70^\circ$
  - c)  $70^\circ$
  - d)  $110^\circ$
7. According to the rational root theorem, which number is not possible rational zero of  $4x^7 + 2x^4 - 10x^3 - 5$ ?
  - a) -1
  - b)  $\frac{5}{4}$
  - c)  $\frac{4}{5}$
  - d) 5
8. The polynomial  $x^3 - kx^2 + 9x$  has three real zeros if and only if, k satisfies
  - a)  $|k| \leq 6$
  - b)  $k = 0$
  - c)  $|k| > 6$
  - d)  $|k| \geq 6$
9. The polynomial  $x^3 + 2x + 3$  has
  - a) one negative and two imaginary zeros
  - b) one positive and two imaginary zeros
  - c) three real zeros
  - d) no zeros
10.  $\sin^{-1}(\cos x) = \frac{\pi - x}{2}$  is valid for
  - a)  $-\pi \leq x \leq 0$
  - b)  $0 \leq x \leq \pi$
  - c)  $-\frac{\pi}{2} \leq x \leq \frac{\pi}{2}$
  - d)  $-\frac{\pi}{4} \leq x \leq \frac{3\pi}{4}$
11.  $\tan^{-1}\left(\frac{1}{4}\right) + \tan^{-1}\left(\frac{2}{9}\right)$  is equal to
  - a)  $\frac{1}{2} \cos^{-1}\left(\frac{3}{5}\right)$
  - b)  $\frac{1}{2} \sin^{-1}\left(\frac{3}{5}\right)$
  - c)  $\frac{1}{2} \tan^{-1}\left(\frac{3}{5}\right)$
  - d)  $\tan^{-1}\left(\frac{1}{2}\right)$

12. If  $\cot^{-1}(\sqrt{\sin \alpha}) + \tan^{-1}(\sqrt{\sin \alpha}) = u$  then,  $\cos 2u$  is equal to  
 a)  $\tan^2 \alpha$       b) 0      c) -1      d)  $\tan 2\alpha$
13. The circle  $x^2 + y^2 = 4x + 8y + 5$  intersects the line  $3x - 4y = m$  at two distinct points if  
 a)  $15 < m < 65$       b)  $35 < m < 85$       c)  $-85 < m < -35$       d)  $-35 < m < 15$
14. Area of the greatest rectangle inscribed in the ellipse  $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$  is  
 a)  $2ab$       b)  $ab$       c)  $\sqrt{ab}$       d)  $\frac{a}{b}$
15. The locus of a point whose distance from  $(-2, 0)$  is  $\frac{2}{3}$  times its distance from the line  $x = \frac{-9}{2}$  is  
 a) a parabola      b) a hyperbola      c) an ellipse      d) a circle
16. If  $\vec{a}$  and  $\vec{b}$  are parallel vectors, then  $[\vec{a}, \vec{c}, \vec{b}]$  is equal to  
 a) 2      b) -1      c) 1      d) 0
17. If  $\vec{a} \cdot \vec{b} = \vec{b} \cdot \vec{c} = \vec{c} \cdot \vec{a} = 0$ , then the value of  $[\vec{a}, \vec{b}, \vec{c}]$  is  
 a)  $|\vec{a}| |\vec{b}| |\vec{c}|$       b)  $\frac{1}{3} |\vec{a}| |\vec{b}| |\vec{c}|$       c) 1      d) -1
18. If  $\vec{a}, \vec{b}, \vec{c}$  are three non-coplanar unit vectors such that  $\vec{a} \times (\vec{b} \times \vec{c}) = \frac{\vec{b} + \vec{c}}{\sqrt{2}}$ , then the angle between  $\vec{a}$  and  $\vec{b}$  is  
 a)  $\frac{\pi}{2}$       b)  $\frac{3\pi}{4}$       c)  $\frac{\pi}{4}$       d)  $\pi$
19. Find the  $|d_1 - d_2|$  between the parallel plane  $x + 2y - 2z + 1 = 0$  and  $2x + 4y - 4z + 5 = 0$   
 a) 2      b) -1      c) 1      d) -2
20. If  $5x + 2y = 3$  and  $3x + 2y = 5$  are the system of linear equations, then  $|A| =$   
 a) 1      b) 2      c) 3      d) 4

## Part - II

II. Answer any 7 questions. (Q.No.30 is compulsory)

7 x 2 = 14

21. If A is a non-singular matrix of odd order, prove that  $|\text{adj} A|$  is positive.

22. Prove that  $\begin{bmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{bmatrix}$  is orthogonal.

23. Simplifying the following :  $i^{50} + \frac{1}{i^{50}}$

24. If  $|z| = 2$ , show that  $3 \leq |z + 3 + 4i| \leq 7$

25. Find a polynomial equation of minimum degree with rational coefficients having  $2 - \sqrt{3}$  as a root.
26. Discuss the nature of the roots of the following polynomial :  
 $x^{2018} + 1947x^{1950} + 15x^8 + 26x^6 + 2019$
27. Find the principal value of  $\cos^{-1}\left(\frac{1}{2}\right)$
28. The maximum and minimum distances of the Earth from the Sun respectively are  $152 \times 10^6$  km and  $94.5 \times 10^6$  km. The sun is at one focus of the elliptical orbit. Find the distance from the sun to the other focus.
29. If  $2\hat{i} - \hat{j} + 3\hat{k}$ ,  $3\hat{i} + 2\hat{j} + \hat{k}$ ,  $\hat{i} + m\hat{j} + 4\hat{k}$  are coplanar, find the value of m.
30. Prove that  $[\vec{a} - \vec{b}, \vec{b} - \vec{c}, \vec{c} - \vec{a}] = 0$

## Part - III

III. Answer any 7 questions. (Q.No.40 is compulsory)

7 x 3 = 21

31. Find the adjoint of the following :  $\begin{bmatrix} -3 & 4 \\ 6 & 2 \end{bmatrix}$

32. Find the rank of the matrix  $\begin{bmatrix} 1 & 2 & 3 \\ 2 & 1 & 4 \\ 3 & 0 & 5 \end{bmatrix}$  by reducing it to a row-echelon form.

33. Write the following in the rectangular form :  $\frac{10 - 5i}{6 + 2i}$

34. Show that the equation  $z^2 = \bar{z}$  has four solutions.

35. If  $\alpha$  and  $\beta$  are the roots of the quadratic equation  $2x^2 - 7x + 13 = 0$ , construct a quadratic equation whose roots are  $\alpha^2$  and  $\beta^2$ .

36. If p is real, discuss the nature of the roots of the equation  $4x^2 + 4px + p + 2 = 0$  in terms of p.

37. Find the value of  $\sin^{-1}(-1) + \cos^{-1}\left(\frac{1}{2}\right) + \cot^{-1}(2)$

38. Find the general equation of the circle whose diameter is the line segment joining the points  $(-4, -2)$  and  $(1, 1)$

39. If  $\vec{a} = \hat{i} - \hat{j}$ ,  $\vec{b} = \hat{i} - \hat{j} - 4\hat{k}$ ,  $\vec{c} = 3\hat{i} - \hat{k}$  and  $\vec{d} = 2\hat{i} + 5\hat{j} + \hat{k}$ , verify that

$$(\vec{a} \times \vec{b}) \times (\vec{c} \times \vec{d}) = [\vec{a}, \vec{b}, \vec{d}] \vec{c} - [\vec{a}, \vec{b}, \vec{c}] \vec{d}$$

40. Show that the points  $(2, 3, 4)$ ,  $(-1, 4, 5)$  and  $(8, 1, 2)$  are collinear.

## Part - IV

## IV. Answer all the questions.

7 x 5 = 35

41. a) By using Gaussian elimination method, balance the chemical reaction equation  
 $C_2H_6 + O_2 \rightarrow H_2O + CO_2$  (OR)
- b) Investigate for what values of  $\lambda$  and  $\mu$  the system of linear equations  
 $x + 2y + z = 7$ ,  $x + y + \lambda z = \mu$ ,  $x + 3y - 5z = 5$  has  
 (i) no solution (ii) a unique solution (iii) an infinite number of solutions
42. a) Suppose  $z_1$ ,  $z_2$  and  $z_3$  are the vertices of an equilateral triangle inscribed in the circle  $|z| = 2$  if  $z_1 = 1 + i\sqrt{3}$ , then find  $z_2$  and  $z_3$ . (OR)
- b) If  $z = x + iy$  and  $\arg\left(\frac{z-i}{z+2}\right) = \frac{\pi}{4}$ , show that  $x^2 + y^2 + 3x - 3y + 2 = 0$
43. a) Solve the equation  $6x^4 - 35x^3 + 62x^2 - 35x + 6 = 0$  (OR)
- b) Solve the equation:  $(x-2)(x-7)(x-3)(x+2) + 19 = 0$
44. a) Find the value of  $\cos^{-1}\left(\cos\left(\frac{4\pi}{3}\right)\right) + \cos^{-1}\left(\cos\left(\frac{5\pi}{4}\right)\right)$  (OR)
- b) Solve:  $\tan^{-1}\left(\frac{x-1}{x-2}\right) + \tan^{-1}\left(\frac{x+1}{x+2}\right) = \frac{\pi}{4}$
45. a) Identify the type of conic and find centre, foci, vertices and directrices.  
 $18x^2 + 12y^2 - 144x + 48y + 120 = 0$  (OR)
- b) Find the equations of tangent and normal to the parabola  $x^2 + 6x + 4y + 5 = 0$  at  $(1, -3)$
46. a) Prove by vector method that the perpendiculars (altitudes) from the vertices to the opposite sides of a triangle are concurrent. (OR)
- b) Find the non-parametric form of vector equation and cartesian equation of the plane passing through the point  $(1, -2, 4)$  and perpendicular to the plane  $x + 2y - 3z = 11$  and parallel to the line  $\frac{x+7}{3} = \frac{y+3}{-1} = \frac{z}{1}$
47. a) If the equation of the ellipse is  $\frac{(x-11)^2}{484} + \frac{y^2}{64} = 1$  (x and y are measured in centimetres) where to the nearest centimetres, should the patient's kidney stone be placed so that the reflected sound hits the kidney stone? (OR)
- b) A fish tank can be filled in 10 minutes using both pumps A and B simultaneously. However, pump B can pump water in or out at the same rate. If pump B is inadvertently run in reverse, then the tank will be filled in 30 minutes. How long would it take each pump to fill the tank by itself? (Use Cramer's rule to solve the problem)

\*\*\*\*\*