



SRI RAGHAVENDRA TUITION CENTER

MATRIX

11th Standard

Maths

Date : 14-07-24

Reg.No. :

Exam Time : 01:30 Hrs

Total Marks : 60

T

EACHER NAME: P.DEEPAK M.Sc.,M.A.,B.Ed.,DCA.,TET-1.,TET-2.,

PHONE NUMBER : 9944249262

EMAIL: darthi99ktp@gmail.com

Centum Book Available

I. Multiple Choice Question

25 x 1 = 25

- 1) If $a_{ij} = \frac{1}{2}(3i - 2j)$ and $A = [a_{ij}]_{2 \times 2}$ is
 (a) $\begin{bmatrix} \frac{1}{2} & 2 \\ -\frac{1}{2} & 1 \end{bmatrix}$ (b) $\begin{bmatrix} \frac{1}{2} & -\frac{1}{2} \\ 2 & 1 \end{bmatrix}$ (c) $\begin{bmatrix} 2 & 2 \\ \frac{1}{2} & -\frac{1}{2} \end{bmatrix}$ (d) $\begin{bmatrix} -\frac{1}{2} & \frac{1}{2} \\ 1 & 2 \end{bmatrix}$
- 2) What must be the matrix X, if $2x + \begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix} = \begin{bmatrix} 3 & 8 \\ 7 & 2 \end{bmatrix}$?
 (a) $\begin{bmatrix} 1 & 3 \\ 2 & -1 \end{bmatrix}$ (b) $\begin{bmatrix} 1 & -3 \\ 2 & -1 \end{bmatrix}$ (c) $\begin{bmatrix} 2 & 6 \\ 4 & -2 \end{bmatrix}$ (d) $\begin{bmatrix} 2 & -6 \\ 4 & -2 \end{bmatrix}$
- 3) Which one of the following is not true about the matrix $\begin{bmatrix} 1 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 5 \end{bmatrix}$?
 (a) a scalar matrix (b) a diagonal matrix (c) an upper triangular matrix (d) a lower triangular matrix
- 4) If A and B are two matrices such that $A + B$ and AB are both defined, then
 (a) A and B are two matrices not necessarily of same order (b) A and B are square matrices of same order
 (c) Number of columns of A is equal to the number of rows of B (d) $A = B$.
- 5) If $A = \begin{bmatrix} \lambda & 1 \\ -1 & -\lambda \end{bmatrix}$, then for what value of λ , $A^2 = O$?
 (a) 0 (b) ± 1 (c) -1 (d) 1
- 6) If $A = \begin{bmatrix} 1 & 2 & 2 \\ 2 & 1 & -2 \\ a & 2 & b \end{bmatrix}$ is a matrix satisfying the equation $AA^T = 9I$, where I is 3×3 identity matrix, then the ordered pair (a, b) is equal to
 (a) (2, -1) (b) (-2, 1) (c) (2, 1) (d) (-2, -1)
- 7) If $A = \begin{bmatrix} 1 & -1 \\ 2 & -1 \end{bmatrix}$, $B = \begin{bmatrix} a & 1 \\ b & -1 \end{bmatrix}$ and $(A + B)^2 = A^2 + B^2$, then the values of a and b are
 (a) $a = 4, b = 1$ (b) $a = 1, b = 4$ (c) $a = 0, b = 4$ (d) $a = 2, b = 4$
- 8) If the points $(x, -2), (5, 2), (8, 8)$ are collinear, then x is equal to
 (a) -3 (b) $\frac{1}{3}$ (c) 1 (d) 3
- 9) If $A = \begin{bmatrix} a & x \\ y & a \end{bmatrix}$ and if $xy = 1$, then $\det(A A^T)$ is equal to
 (a) $(a - 1)^2$ (b) $(a^2 + 1)^2$ (c) $a^2 - 1$ (d) $(a^2 - 1)^2$
- 10) If A is a square matrix, then which of the following is not symmetric?
 (a) $A + A^T$ (b) AA^T (c) $A^T A$ (d) $A - A^T$

- 11) If A and B are symmetric matrices of order n, where $(A \neq B)$, then
 (a) A + B is skew-symmetric (b) A + B is symmetric (c) A + B is a diagonal matrix (d) A + B is a zero matrix
- 12) The value of x, for which the matrix $A = \begin{bmatrix} e^{x-2} & e^{7+x} \\ e^{2+x} & e^{2x+3} \end{bmatrix}$ is singular
 (a) 9 (b) 8 (c) 7 (d) 6
- 13) If the square of the matrix $\begin{bmatrix} \alpha & \beta \\ \gamma & -\alpha \end{bmatrix}$ is the unit matrix of order 2, then α, β and γ should satisfy the relation.
 (a) $1 + \alpha^2 + \beta\gamma = 0$ (b) $1 - \alpha^2 - \beta\gamma = 0$ (c) $1 - \alpha^2 + \beta\gamma = 0$ (d) $1 + \alpha^2 - \beta\gamma = 0$
- 14) If $\begin{vmatrix} 2a & x_1 & y_1 \\ 2b & x_2 & y_2 \\ 2c & x_3 & y_3 \end{vmatrix} = \frac{abc}{2} \neq 0$, then the area of the triangle whose vertices are $(\frac{x_1}{a}, \frac{y_1}{a}), (\frac{x_2}{b}, \frac{y_2}{b}), (\frac{x_3}{c}, \frac{y_3}{c})$ is
 (a) $\frac{1}{4}$ (b) $\frac{1}{4}abc$ (c) $\frac{1}{8}$ (d) $\frac{1}{8}abc$
- 15) If $a \neq b, b, c$ satisfy $\begin{vmatrix} a & 2b & 2c \\ 3 & b & c \\ 4 & a & b \end{vmatrix} = 0$, then abc =
 (a) a + b + c (b) 0 (c) b^3 (d) ab + bc
- 16) If $\Delta = \begin{vmatrix} a & b & c \\ x & y & z \\ p & q & r \end{vmatrix}$, then $\begin{vmatrix} ka & kb & kc \\ kx & ky & kz \\ kp & kq & kr \end{vmatrix}$ is
 (a) Δ (b) $k\Delta$ (c) $3k\Delta$ (d) $k^3\Delta$
- 17) A root of the equation $\begin{vmatrix} 3-x & -6 & 3 \\ -6 & 3-x & 3 \\ 3 & 3 & -6-x \end{vmatrix} = 0$ is
 (a) 6 (b) 3 (c) 0 (d) -6
- 18) The value of the determinant of $A = \begin{bmatrix} 0 & a & -b \\ -a & 0 & c \\ b & -c & 0 \end{bmatrix}$ is
 (a) $-2abc$ (b) abc (c) 0 (d) $a^2 + b^2 + c^2$
- 19) If x_1, x_2, x_3 as well as y_1, y_2, y_3 are in geometric progression with the same common ratio, then the points $(x_1, y_1), (x_2, y_2), (x_3, y_3)$ are
 (a) vertices of an equilateral triangle (b) vertices of a right angled triangle (c) vertices of a right angled isosceles triangle
 (d) collinear
- 20) If $A = \begin{vmatrix} -1 & 2 & 4 \\ 3 & 1 & 0 \\ -2 & 4 & 2 \end{vmatrix}$ and $B = \begin{vmatrix} -2 & 4 & 2 \\ 6 & 2 & 0 \\ -2 & 4 & 8 \end{vmatrix}$, then B is given by
 (a) $B = 4A$ (b) $B = -4A$ (c) $B = -A$ (d) $B = 6A$
- 21) If $\lfloor . \rfloor$ denotes the greatest integer less than or equal to the real number under consideration and $-1 \leq x < 0, 0 \leq y < 1, 1 \leq z < 2$, then the value of the determinant $\begin{vmatrix} \lfloor x \rfloor + 1 & \lfloor y \rfloor & \lfloor z \rfloor \\ \lfloor x \rfloor & \lfloor y \rfloor + 1 & \lfloor z \rfloor \\ \lfloor x \rfloor & \lfloor y \rfloor & \lfloor z \rfloor + 1 \end{vmatrix}$ is
 (a) $\lfloor z \rfloor$ (b) $\lfloor y \rfloor$ (c) $\lfloor x \rfloor$ (d) $\lfloor x \rfloor + 1$
- 22) If A is skew-symmetric of order n and C is a column matrix of order $n \times 1$, then $C^T AC$ is
 (a) an identity matrix of order n (b) an identity matrix of order 1 (c) a zero matrix of order 1
 (d) an identity matrix of order 2
- 23) The matrix A satisfying the equation $\begin{bmatrix} 1 & 3 \\ 0 & 1 \end{bmatrix} A = \begin{bmatrix} 1 & 1 \\ 0 & -1 \end{bmatrix}$ is
 (a) $\begin{bmatrix} 1 & 4 \\ -1 & 0 \end{bmatrix}$ (b) $\begin{bmatrix} 1 & -4 \\ 1 & 0 \end{bmatrix}$ (c) $\begin{bmatrix} 1 & 4 \\ 0 & -1 \end{bmatrix}$ (d) $\begin{bmatrix} 1 & -4 \\ 1 & 1 \end{bmatrix}$

- 24) If $A + I = \begin{bmatrix} 3 & -2 \\ 4 & 1 \end{bmatrix}$, then $(A + I)(A - I)$ is equal to
 (a) $\begin{bmatrix} -5 & -4 \\ 8 & -9 \end{bmatrix}$ (b) $\begin{bmatrix} -5 & 4 \\ -8 & 9 \end{bmatrix}$ (c) $\begin{bmatrix} 5 & 4 \\ 8 & 9 \end{bmatrix}$ (d) $\begin{bmatrix} -5 & -4 \\ -8 & -9 \end{bmatrix}$
- 25) Let A and B be two symmetric matrices of same order. Then which one of the following statement is not true?
 (a) A + B is a symmetric matrix (b) AB is a symmetric matrix (c) $AB = (BA)^T$ (d) $A^T B = AB^T$

II. Answer all question

5 x 2 = 10

- 26) Simplify : $\sec\theta \begin{bmatrix} \sec\theta & \tan\theta \\ \tan\theta & \sec\theta \end{bmatrix} - \tan\theta \begin{bmatrix} \tan\theta & \sec\theta \\ \sec\theta & \tan\theta \end{bmatrix}$
- 27) If a, b, c and x are positive real numbers, then show that $\begin{vmatrix} (a^x + a^{-x})^2 & (a^x - a^{-x})^2 & 1 \\ (b^x + b^{-x})^2 & (b^x - b^{-x})^2 & 1 \\ (c^x + c^{-x})^2 & (c^x - c^{-x})^2 & 1 \end{vmatrix}$ is zero.

- 28) Find the sum A + B + C if A, B, C are given by

$$A = \begin{bmatrix} \sin^2 \theta & 1 \\ \cot^2 \theta & 0 \end{bmatrix}, B = \begin{bmatrix} \cos^2 \theta & 0 \\ -\operatorname{cosec}^2 \theta & 1 \end{bmatrix} \text{ and } C = \begin{bmatrix} 0 & -1 \\ -1 & 0 \end{bmatrix}$$

- 29) Suppose that a matrix has 12 elements. What are the possible orders it can have? What if it has 7 elements?

- 30) Evaluate $\begin{vmatrix} 2014 & 2017 & 0 \\ 2020 & 2023 & 1 \\ 2023 & 2026 & 0 \end{vmatrix}$

III. ANSWER ALL QUESTION

5 x 3 = 15

- 31) If $A = \begin{bmatrix} 1 & 0 & 2 \\ 0 & 2 & 1 \\ 2 & 0 & 3 \end{bmatrix}$ and $A^3 - 6A^2 + 7A + KI = O$, find the value of k.

- 32) Find x, y, a, and b if $\begin{bmatrix} 3x + 4y & 6 & x - 2y \\ a + b & 2a - b & -3 \end{bmatrix} = \begin{bmatrix} 2 & 6 & 4 \\ 5 & -5 & -3 \end{bmatrix}$

- 33) If $A = \begin{bmatrix} 0 & c & b \\ c & 0 & a \\ b & a & 0 \end{bmatrix}$, compute A^2

- 34) If A and B are symmetric matrices of same order, prove that
 $AB - BA$ is a skew-symmetric matrix.

- 35) Find the value of x if $\begin{vmatrix} x - 1 & x & x - 2 \\ 0 & x - 2 & x - 3 \\ 0 & 0 & x - 3 \end{vmatrix} = 0$

III. ANSWER ALL QUESTION

2 x 5 = 10

- 36) a) Show that $\begin{vmatrix} a^2 + x^2 & ab & ac \\ ab & b^2 + x^2 & bc \\ ac & bc & c^2 + x^2 \end{vmatrix}$ is divisible by x^4 .

(OR)

- b) If $A = \begin{bmatrix} 1 & 2 & 2 \\ 2 & 1 & -2 \\ x & 2 & y \end{bmatrix}$ is a matrix such that $AA^T = 9I$, find the values of x and y.

(OR)

- c) Show that $\begin{vmatrix} x + 2a & y + 2b & z + 2c \\ x & y & z \\ a & b & c \end{vmatrix} = 0$.

37)

a) If $A = \begin{bmatrix} \frac{1}{2} & \alpha \\ 0 & \frac{1}{2} \end{bmatrix}$, prove that $\sum_{k=1}^n \det(A^k) = \frac{1}{3}(1 - \frac{1}{4^n})$.

(OR)

b) Prove that $\begin{vmatrix} 1+a & 1 & 1 \\ 1 & 1+b & 1 \\ 1 & 1 & 1+c \end{vmatrix} = abc \left(1 + \frac{1}{a} + \frac{1}{b} + \frac{1}{c}\right)$.

(OR)

c) Show that $\begin{vmatrix} 1 & 1 & 1 \\ x & y & z \\ x^2 & y^2 & z^2 \end{vmatrix} = (x-y)(y-z)(z-x)$.

ALL THE BEST

www.Padasalai.Net