

ONE MARKS

XI – MATHEMATICS

1. Sets, Relation and Functions

- If $A = \{(x, y) : y = e^x, x \in \mathbb{R}\}$ $B = \{(x, y) : y = e^{-x}, x \in \mathbb{R}\}$ then $n(A \cap B)$ is
 - Infinity
 - 0
 - 1
 - 2
- If $A = \{(x, y) : y = \sin x, x \in \mathbb{R}\}$ and $B = \{(x, y) : y = \cos x, x \in \mathbb{R}\}$ then $A \cap B$ contains
 - No element
 - infinitely many elements
 - Only one element
 - cannot be determined.
- The relation R defined on a set $A = \{0, -1, 1, 2\}$ by xRy if $|x^2 + y^2| \leq 2$, then which one of the following is true?
 - $R = \{(0, 0), (0, -1), (0, 1), (-1, 0), (-1, 1), (1, 2), (1, 0)\}$
 - $R^{-1} = \{(0, 0), (0, -1), (0, 1), (-1, 0), (-1, 0), (1, 0)\}$
 - Domain of R is $\{0, -1, 1, 2\}$
 - Range of R is $\{0, -1, 1\}$
- If $f(x) = |x - 2| + |x + 2|$, $x \in \mathbb{R}$, then
 - $f(x) = \begin{cases} -2x & \text{if } x \in (-\infty, -2] \\ 4 & \text{if } x \in (-2, 2] \\ 2x & \text{if } x \in (2, \infty) \end{cases}$
 - $f(x) = \begin{cases} 2x & \text{if } x \in (-\infty, -2] \\ 4x & \text{if } x \in (-2, 2] \\ -2x & \text{if } x \in (2, \infty) \end{cases}$
 - $f(x) = \begin{cases} -2x & \text{if } x \in (-\infty, -2] \\ -4x & \text{if } x \in (-2, 2] \\ 2x & \text{if } x \in (2, \infty) \end{cases}$
 - $f(x) = \begin{cases} -2x & \text{if } x \in (-\infty, -2] \\ 2x & \text{if } x \in (-2, 2] \\ 2x & \text{if } x \in (2, \infty) \end{cases}$
- Let \mathbb{R} be the set of all real numbers. Consider the following subsets of the plane $\mathbb{R} \times \mathbb{R}$: $S = \{(x, y) : y = x + 1 \text{ and } 0 < x < 2\}$ and $T = \{(x, y) : x - y \text{ is an integer}\}$
Then which of the following is true?
 - T is an equivalence relation but S is not an equivalence relation.
 - Neither S nor T is an equivalence relation
 - Both S and T are equivalence relation
 - S is an equivalence relation but T is not an equivalence relation.
- Let A and B be subsets of the universal set N , the set of natural numbers. Then $A' \cup [(A \cap B) \cup B']$ is
 - A
 - A'
 - B
 - N
- The number of students who take both the subjects Mathematics and Chemistry is 70. This represents 10% of the enrollment in Mathematics and 14% of the enrollment in Chemistry. The number of students take at least one of these two subjects, is
 - 1120
 - 1130
 - 1100
 - insufficient data
- If $n((A \times B) \cap (A \times C)) = 8$ and $n(B \cap C) = 2$, then $n(A)$ is
 - 6
 - 4
 - 8
 - 16
- If $n(A) = 2$ and $n(B \cup C) = 3$, then $n[(A \times B) \cup (A \times C)]$ is
 - 2^3
 - 3^2
 - 6
 - 5
- If two sets A and B have 17 elements in common, then the number of elements common to the set $A \times B$ and $B \times A$ is
 - 2^{17}
 - 17^2
 - 34
 - insufficient data
- For non-empty sets A and B , if $A \subset B$ then $(A \times B) \cap (B \times A)$ is equal to
 - $A \cap B$
 - $A \times A$
 - $B \times B$
 - none of these

ONE MARKS

XI – MATHEMATICS

12. The number of relations on a set containing 3 elements is
 1. 9 2. 81 3. 512 4. 1024
13. Let R be the universal relation on a set X with more than one element. Then R is
 1. Not reflexive 2. not symmetric 3. transitive 4. none of the above
14. Let $X = \{1, 2, 3, 4\}$ and $R = \{(1, 1), (1, 2), (1, 3), (2, 2), (3, 3), (2, 1), (3, 1), (1, 4), (4, 1)\}$. Then R is
 1. Reflexive 2. symmetric 3. transitive 4. equivalence
15. The range of the function $\frac{1}{1-2\sin x}$ is
 1. $(-\infty, -1) \cup (\frac{1}{3}, \infty)$ 2. $(-1, \frac{1}{3})$ 3. $[-1, \frac{1}{3}]$ 4. $(-\infty, -1] \cup [\frac{1}{3}, \infty)$
16. The range of the function $f(x) = ||x| - x|, x \in R$ is
 1. $[0, 1]$ 2. $[0, \infty)$ 3. $[0, 1)$ 4. $(0, 1)$
17. The rule $f(x) = x^2$ is a bijection if the domain and the co-domain are given by
 1. R, R 2. R, $(0, \infty)$ 3. $(0, \infty)$, R 4. $[0, \infty)$, $[0, \infty)$
18. The number of constant functions from a set containing m elements to a set containing n elements is
 1. mn 2. m 3. n 4. m + n
19. The function $f : [0, 2\pi] \rightarrow [-1, 1]$ defined by $f(x) = \sin x$ is
 1. one-to-one 2. onto 3. bijection 4. cannot be defined
20. If the function $f : [-3, 3] \rightarrow S$ defined by $f(x) = x^2$ is onto, then S is
 1. $[-9, 9]$ 2. R 3. $[-3, 3]$ 4. $[0, 9]$
21. Let $x = \{1, 2, 3, 4\}$, $Y = \{a, b, c, d\}$ and $f = \{(1, a), (4, b), (2, c), (3, d), (2, d)\}$. Then f is
 1. An one-to-one function 2. an onto function
 3. A function which is not one-to-one 4. not a function
22. The inverse of $f(x) = \begin{cases} x & \text{if } x < 1 \\ x^2 & \text{if } 1 \leq x \leq 4 \\ 8\sqrt{x} & \text{if } x > 4 \end{cases}$ is
 1. $f^{-1}(x) = \begin{cases} x & \text{if } x < 1 \\ \sqrt{x} & \text{if } 1 \leq x \leq 16 \\ \frac{x^2}{64} & \text{if } x > 16 \end{cases}$ 2. $f^{-1}(x) = \begin{cases} -x & \text{if } x < 1 \\ \sqrt{x} & \text{if } 1 \leq x \leq 16 \\ \frac{x^2}{64} & \text{if } x > 16 \end{cases}$
 3. $f^{-1}(x) = \begin{cases} x^2 & \text{if } x < 1 \\ \sqrt{x} & \text{if } 1 \leq x \leq 16 \\ \frac{x^2}{64} & \text{if } x > 16 \end{cases}$ 4. $f^{-1}(x) = \begin{cases} 2x & \text{if } x < 1 \\ \sqrt{x} & \text{if } 1 \leq x \leq 16 \\ \frac{x^2}{8} & \text{if } x > 16 \end{cases}$
23. Let $f : R \rightarrow R$ be defined by $f(x) = 1 - |x|$. Then the range of f is
 1. R 2. $(1, \infty)$ 3. $(-1, \infty)$ 4. $(-\infty, 1]$
24. The function $f : R \rightarrow R$ is defined by $f(x) = \sin x + \cos x$ is
 1. An odd function
 2. Neither an odd function nor an even function
 3. An even function
 4. Both odd function and even function
25. The function $f : R \rightarrow R$ is define by $f(x) = \frac{(x^2 + \cos x)(1 + x^4)}{(x - \sin x)(2x - x^3)}$
 1. An odd function 2. neither an odd function nor an even function

ONE MARKS

XI – MATHEMATICS

3. An even function 4. both odd function and even function.

2. Basic algebra

26. If $|x + 2| \leq 9$, then x belongs to
 1. $(-\infty, -7)$ 2. $[-11, 7]$ 3. $(-\infty, -7) \cup [11, \infty)$ 4. $(-11, 7)$
27. Given that x, y and b are real numbers $x < y, b > 0$, then
 1. $xb < yb$ 2. $xb > yb$ 3. $xb \leq yb$ 4. $\frac{x}{b} \geq \frac{y}{b}$
28. If $\frac{|x-2|}{x-2} \geq 0$, then x belongs to
 1. $[2, \infty)$ 2. $(2, \infty)$ 3. $(-\infty, 2)$ 4. $(-2, \infty)$
29. The solution of $5x - 1 < 24$ and $5x + 1 > -24$ is
 1. $(4, 5)$ 2. $(-5, -4)$ 3. $(-5, 5)$ 4. $(-5, 4)$
30. The solution set of the following inequality $|x - 1| \geq |x - 3|$ is
 1. $[0, 2]$ 2. $[2, \infty)$ 3. $(0, 2)$ 4. $(-\infty, 2)$
31. The value of $\log_{\sqrt{2}} 512$ is
 1. 16 2. 18 3. 9 4. 12
32. The value of $\log_3 \frac{1}{81}$ is
 1. -2 2. -8 3. -4 4. -9
33. If $\log_{\sqrt{x}} 0.25 = 4$, then the value of x is
 1. 0.5 2. 2.5 3. 1.5 4. 1.25
34. The value of $\log_a b \log_b c \log_c a$ is
 1. 2 2. 1 3. 3 4. 4
35. If 3 is the logarithm of 343, then the base is
 1. 5 2. 7 3. 6 4. 9
36. Find a so that the sum and product of the roots of the equation $2x^2 + (a - 3)x + 3a - 5 = 0$ are equal is
 1. 1 2. 2 3. 0 4. 4
37. If a and b are the roots of the equation $x^2 - kx + 16 = 0$ and satisfy $a^2 + b^2 = 32$, then the value of k is
 1. 10 2. -8 3. -8, 8 4. 6
38. The number of solutions of $x^2 + |x - 1| = 1$ is
 1. 1 2. 0 3. 2 4. 3
39. The equation whose roots are numerically equal but opposite in sign to the roots of $3x^2 - 5x - 7 = 0$ is
 1. $3x^2 - 5x - 7 = 0$ 2. $3x^2 + 5x - 7 = 0$ 3. $3x^2 - 5x + 7 = 0$ 4. $3x^2 + x - 7$
40. If 8 and 2 are the roots of $x^2 + ax + b = 0$ and 3, 3 are the roots of $x^2 + dx + b = 0$, then the roots of the equation $x^2 + ax + b = 0$ are
 1. 1, 2 2. -1, 1 3. 9, 1 4. -1, 2
41. If a and b are the real roots of the equation $x^2 - kx + c = 0$, then the distance between the points $(a, 0)$ and $(b, 0)$ is
 1. $\sqrt{k^2 - 4c}$ 2. $\sqrt{4k^2 - c}$ 3. $\sqrt{4c - k^2}$ 4. $\sqrt{k - 8c}$
42. If $\frac{kx}{(x+2)(x-1)} = \frac{2}{x+2} + \frac{1}{x-1}$, then the value of k is
 1. 1 2. 2 3. 3 4. 4
43. If $\frac{1-2x}{3+2x-x^2} = \frac{A}{3-x} + \frac{B}{x+1}$, then the value of $A + B$ is
 1. $\frac{-1}{2}$ 2. $\frac{-2}{3}$ 3. $\frac{1}{2}$ 4. $\frac{2}{3}$
44. The number of roots of $(x+3)^4 + (x+5)^4 = 16$ is
 1. 4 2. 2 3. 3 4. 0
45. The value of $\log_3 11 \cdot \log_{11} 13 \cdot \log_{13} 15 \cdot \log_{15} 27 \cdot \log_{27} 81$ is

ONE MARKS

XI – MATHEMATICS

1. 1 2. 2 3. 3 4. 4

3. Trigonometry

46. $\frac{1}{\cos 80^\circ} - \frac{\sqrt{3}}{\sin 80^\circ} =$
 1. $\sqrt{2}$ 2. $\sqrt{3}$ 3. 2 4. 4
47. If $\cos 28^\circ + \sin 28^\circ = k^3$, then $\cos 17^\circ$ is equal to
 1. $\frac{k^3}{\sqrt{2}}$ 2. $-\frac{k^3}{\sqrt{2}}$ 3. $\pm \frac{k^3}{\sqrt{2}}$ 4. $-\frac{k^3}{\sqrt{3}}$
48. The maximum value of $4\sin^2 x + 3\cos^2 x + \sin \frac{x}{2} + \cos \frac{x}{2}$ is
 1. $4 + \sqrt{2}$ 2. $3 + \sqrt{2}$ 3. 9 4. 4
49. $\left(1 + \cos \frac{\pi}{8}\right)\left(1 + \cos \frac{3\pi}{8}\right)\left(1 + \cos \frac{5\pi}{8}\right)\left(1 + \cos \frac{7\pi}{8}\right) =$
 1. $\frac{1}{8}$ 2. $\frac{1}{2}$ 3. $\frac{1}{\sqrt{3}}$ 4. $\frac{1}{\sqrt{2}}$
50. If $\pi < 2\theta < \frac{3\pi}{2}$, then $\sqrt{2 + \sqrt{2 + 2\cos 4\theta}}$ equals to
 1. $-2\cos\theta$ 2. $-2\sin\theta$ 3. $2\cos\theta$ 4. $2\sin\theta$
51. If $\tan 40^\circ = \lambda$, then $\frac{\tan 140^\circ - \tan 130^\circ}{1 + \tan 140^\circ \tan 130^\circ} =$
 1. $\frac{1-\lambda^2}{\lambda}$ 2. $\frac{1+\lambda^2}{\lambda}$ 3. $\frac{1+\lambda^2}{2\lambda}$ 4. $\frac{1-\lambda^2}{2\lambda}$
52. $\cos 1^\circ + \cos 2^\circ + \cos 3^\circ + \dots + \cos 179^\circ =$
 1. 0 2. 1 3. -1 4. 89
53. Let $f_k(x) = \frac{1}{k}[\sin^k x + \cos^k x]$ where $x \in \mathbb{R}$ and $k \geq 1$. The $f_4(x) - f_6(x) =$
 1. $\frac{1}{4}$ 2. $\frac{1}{12}$ 3. $\frac{1}{6}$ 4. $\frac{1}{3}$
54. Which of the following is not true?
 1. $\sin \theta = -\frac{1}{4}$ 2. $\cos \theta = -1$ 3. $\tan \theta = 25$ 4. $\sec \theta = \frac{1}{4}$
55. $\cos 2\theta \cos 2\phi + \sin^2(\theta - \phi) - \sin^2(\theta + \phi)$ is equal to
 1. $\sin 2(\theta + \phi)$ 2. $\cos 2(\theta + \phi)$ 3. $\sin 2(\theta - \phi)$ 4. $\cos 2(\theta - \phi)$
56. $\frac{\sin(A-B)}{\cos A \cos B} + \frac{\sin(B-C)}{\cos B \cos C} + \frac{\sin(C-A)}{\cos C \cos A}$ is
 1. $\sin A + \sin B + \sin C$ 3. 0
 2. 1 4. $\cos A + \cos B + \cos C$
57. If $\cos p\theta + \cos q\theta = 0$ and if $p \neq q$, then θ is equal to (n is any integer)
 1. $\frac{\pi(3n+1)}{p-q}$ 2. $\frac{\pi(2n+1)}{p+q}$ 3. $\frac{\pi(n+1)}{p+q}$ 4. $\frac{\pi(n+2)}{p+q}$
58. If $\tan \alpha$ and $\tan \beta$ are the roots of $x^2 + ax + b = 0$, then $\frac{\sin(\alpha+\beta)}{\sin \alpha \sin \beta}$ is equal to
 1. $\frac{b}{a}$ 2. $\frac{a}{b}$ 3. $-\frac{a}{b}$ 4. $-\frac{b}{a}$
59. In a triangle ABC, $\sin^2 A + \sin^2 B + \sin^2 C = 2$, then the triangle is
 1. Equilateral triangle 3. right triangle
 2. Isosceles triangle 4. scalene triangle
60. If $f(\theta) = |\sin \theta| + |\cos \theta|$, $\theta \in \mathbb{R}$, then $f(\theta)$ is in the interval
 1. $[0, 2]$ 2. $[1, \sqrt{2}]$ 3. $[1, 2]$ 4. $[0, 1]$
61. $\frac{\cos 6x + 6\cos 4x + 15\cos 2x + 10}{\cos 5x + 5\cos 3x + 10\cos x}$ is equal to
 1. $\cos 2x$ 2. $\cos x$ 3. $\cos 3x$ 4. $2\cos x$
62. The triangle of maximum area with constant perimeter 12 m
 1. Is an equilateral triangle with side 4m
 2. Is an isosceles triangle with sides 2m, 5m, 5m
 3. Is a triangle with sides 3m, 4m, 5m

XI – MATHEMATICS

ONE MARKS

XI – MATHEMATICS

1. 6 2. 9 3. 12 4. 18

78. Everybody in a room shakes hands with everybody else. The total number of shake hands is 66. The number of persons in the room is ____

1. 11 2. 12 3. 10 4. 6

79. Number of sides of a polygon having 44 diagonals is ____

1. 4 2. 4! 3. 11 4. 22

80. If 10 lines are drawn in a plane such that no two of them are parallel and no three are concurrent, then the total number of points of intersection are

1. 45 2. 40 3. 10! 4. 2^{10}

81. In a plane there are 10 points are there out of which 4 points are collinear, then the number of triangles formed is

1. 110 2. $^{10}C_3$ 3. 120 4. 116

82. In $^{2n}C_3 : ^nC_3 = 11 : 1$ then n is

1. 5 2. 6 3. 11 4. 7

83. $^{(n-1)}C_r + ^{(n-1)}C_{(r-1)}$ is

1. $^{(n+1)}C_r$ 2. $^{(n-1)}C_r$ 3. nC_r 4. $^nC_{r-1}$

84. The number of ways of choosing 5 cards out of a deck of 52 cards which include at least one king is

1. $^{52}C_5$ 2. $^{48}C_5$ 3. $^{52}C_5 + ^{48}C_5$ 4. $^{52}C_5 - ^{48}C_5$

85. The number of rectangles that a chess board has

1. 81 2. 9^9 3. 1296 4. 6561

86. The number of 10 digit number that can be written by using the digits 2 and 3 is

1. $^{10}C_2 + ^9C_2$ 2. 2^{10} 3. 2^{10-2} 4. $10!$

87. If P_r stands for rP_r then the sum of the series $1 + P_1 + 2P_2 + 3P_3 + \dots + nP_n$ is

1. P_{n+1} 2. $P_{n+1} - 1$ 3. $P_{n-1} + 1$ 4. $^{(n+1)}P_{(n-1)}$

88. The product of first n odd natural number equals

1. $^{2n}C_n \times ^nP_n$ 2. $\left(\frac{1}{2}\right)^n \times ^{2n}C_n \times ^nP_n$ 3. $\left(\frac{1}{4}\right)^n \times ^{2n}C_n \times ^{2n}P_n$ 4. $^nC_n \times ^nP_n$

89. If $^nC_4, ^nC_5, ^nC_6$ are in AP the value of n can be

1. 14 2. 11 3. 9 4. 5

90. $1 + 3 + 5 + 7 + \dots + 17$ is equal to

1. 101 2. 81 3. 71 4. 61

5. Binomial Theorem, Sequences and Series

91. The value of $2 + 4 + 6 + \dots + 2n$ is

1. $\frac{n(n-1)}{2}$ 2. $\frac{n(n+1)}{2}$ 3. $\frac{2n(2n+1)}{2}$ 4. $n(n+1)$

92. The coefficient of x^6 in $(2 + 2x)^{10}$ is

1. $^{10}C_6$ 2. 2^6 3. $^{10}C_6 2^6$ 4. $^{10}C_6 2^{10}$

93. The coefficient of $x^8 y^{12}$ in the expansion of $(2x + 3y)^{20}$ is

1. 0 2. $2^{83} 3^{12}$ 3. $2^{83} 3^{12} + 2^{12} 3^8$ 4. $^{20}C_8 2^{83} 3^{12}$

94. If $^nC_{10} > ^nC_r$ for all possible r, then a value of n is

1. 10 2. 21 3. 19 4. 20

95. If a is the arithmetic mean and g is the geometric mean of two numbers, then

1. $a \leq g$ 2. $a \geq g$ 3. $a = g$ 4. $a > g$

96. If $(1 + x^2)^2 (1 + x)^n = a_0 + a_1 x + a_2 x^2 + \dots + x^{n+4}$ and if a_0, a_1, a_2 are in AP, then n is

1. 1 2. 2 3. 3 4. 4

97. If a, 8, b are in AP, a, 4, b are in GP, and if a, x, b are in HP then x is

1. 2 2. 1 3. 4 4. 16

ONE MARKS

XI – MATHEMATICS

98. The sequence $\frac{1}{\sqrt{3}}, \frac{1}{\sqrt{3}+\sqrt{2}}, \frac{1}{\sqrt{3}+2\sqrt{2}} \dots$ form an
 1. AP 2. GP 3. HP 4. AGP.
99. The HM of two positive numbers whose AM and GM are 16, 8 respectively is
 1. 10 2. 6 3. 5 4. 4
100. If S_n denotes the sum of n terms of an AP whose common difference is d , the value of $S_n - 2S_{n-1} + S_{n-2}$ is
 1. 0 2. $2d$ 3. $4s$ 4. d^2
101. The remainder when 38^{15} is divided by 13 is
 1. 12 2. 1 3. 11 4. 5
102. The n^{th} term of the sequence 1, 2, 4, 7, 11, is
 1. $N^3 + 3n^2 + 2n$ 2. $n^3 - 3n^2 + 3n$ 3. $\frac{n(n+1)(n+2)}{3}$ 4. $\frac{n^2-n+2}{2}$
103. The sum up to n terms of the series $\frac{1}{\sqrt{1}+\sqrt{3}} + \frac{1}{\sqrt{3}+\sqrt{5}} + \frac{1}{\sqrt{5}+\sqrt{7}} + \dots$ is
 1. $\sqrt{2n+1}$ 2. $\frac{\sqrt{2n+1}}{2}$ 3. $\sqrt{2n+1} - 1$ 4. $\frac{\sqrt{2n+1}-1}{2}$
104. The n^{th} term of the sequence $\frac{1}{2}, \frac{3}{4}, \frac{7}{8}, \frac{15}{16}, \dots$ is
 1. $2^n - n - 1$ 2. $1 - 2^{-n}$ 3. $2^{-n} + n - 1$ 4. 2^{n-1}
105. The sum up to n terms of the series $\sqrt{2} + \sqrt{8} + \sqrt{18} + \sqrt{32} + \dots$ is
 1. $\frac{n(n+1)}{2}$ 2. $2n(n+1)$ 3. $\frac{n(n-1)}{2}$ 4. 1
106. The value of the series $\frac{1}{2} + \frac{7}{4} + \frac{13}{8} + \frac{19}{16} + \dots$ is
 1. 14 2. 7 3. 4 4. 6
107. The sum of an infinite GP is 18. If the first term is 6, the common ratio is
 1. $\frac{1}{3}$ 2. $\frac{2}{3}$ 3. $\frac{1}{6}$ 4. $\frac{3}{4}$
108. The coefficient of x^5 in the series e^{-2x} is
 1. $\frac{2}{3}$ 2. $\frac{3}{2}$ 3. $-\frac{4}{15}$ 4. $\frac{4}{15}$
109. The value of $\frac{1}{2!} + \frac{1}{4!} + \frac{1}{6!} + \dots$ is
 1. $\frac{e^2+1}{2e}$ 2. $\frac{(e+1)^2}{2e}$ 3. $\frac{(e-1)^2}{2e}$ 4. $\frac{e^2-1}{2e}$
110. The value of $1 - \frac{1}{2} \left(\frac{2}{3}\right) + \frac{1}{3} \left(\frac{2}{3}\right)^2 - \frac{1}{4} \left(\frac{2}{3}\right)^3 + \dots$ is
 1. $\log \left(\frac{5}{3}\right)$ 2. $\frac{3}{2} \log \left(\frac{5}{3}\right)$ 3. $\frac{5}{3} \log \left(\frac{5}{3}\right)$ 4. $\frac{2}{3} \log \left(\frac{2}{3}\right)$

6. Two Dimensional Analytical Geometry

111. The equation of the locus of the point whose distance from y - axis is half the distance from origin is
 1. $x^2 + 3y^2 = 0$ 2. $x^2 - 3y^2 = 0$ 3. $3x^2 + y^2 = 0$ 4. $3x^2 - y^2 = 0$
112. Which of the following equation is the locus of $(at^2, 2at)$
 1. $\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$ 2. $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$ 3. $x^2 + y^2 = a^2$ 4. $y^2 = 4ax$
113. Which of the following point lie on the locus of $3x^2 + 3y^2 - 8x - 12 + 17 = 0$
 1. (0, 0) 2. (-2, 3) 3. (1, 2) 4. (0, -1)
114. If the point (8, -5) lies on the locus $\frac{x^2}{16} - \frac{y^2}{25} = k$, then the value of k is
 1. 0 2. 1 3. 2 4. 3
115. Straight line joining the points (2, 3) and (-1, 4) passes through the point (α, β)
 1. $\alpha + 2\beta = 7$ 2. $3\alpha + \beta = 9$ 3. $\alpha + 3\beta = 11$ 4. $3\alpha + \beta = 11$

ONE MARKS

XI – MATHEMATICS

116. The slope of the line which makes an angle 45° with the line $3x - y = -5$ are
 1. 1, -1 2. $\frac{1}{2}, -2$ 3. $1, \frac{1}{2}$ 4. $2, -\frac{1}{2}$
117. Equation of the straight line that forms an isosceles triangle with coordinate axes in the I-quadrant with perimeter $4 + 2\sqrt{2}$ is
 1. $x + y + 2 = 0$ 2. $x + y - 2 = 0$ 3. $x + y - \sqrt{2} = 0$ 4. $x - y + \sqrt{2} = 0$
118. The coordinates of the four vertices of a quadrilateral are $(-2, 4)$, $(-1, 2)$, $(1, 2)$ and $(2, 4)$ taken in order. The equation of the line passing through the vertex $(-1, 2)$ and dividing the quadrilateral in the equal areas is
 1. $x + 1 = 0$ 2. $x + y = 1$ 3. $x + y + 3 = 0$ 4. $x - y = 3 = 0$
119. The intercepts of the perpendicular bisector of the line segment joining $(1, 2)$ and $(3, 4)$ with coordinate axes are
 1. 5, -5 2. 5, 5 3. 5, 3 4. 5, -4
120. The equation of the line with slope 2 and the length of the perpendicular from the origin equal to $\sqrt{5}$ is
 1. $x + 2y = \sqrt{5}$ 2. $2x + y = \sqrt{5}$ 3. $2x + y = 5$ 4. $x + 2y - 5 = 0$
121. A line perpendicular to the line $5x - y = 0$ forms a triangle with the coordinate axes. If the area of the triangle is 5 sq. units, then its equation is
 1. $x + 5y \pm 5\sqrt{2} = 0$ 2. $x - 5y \pm 5\sqrt{2} = 0$ 3. $5x + y \pm 5\sqrt{2} = 0$ 4. $5x - y \pm 5\sqrt{2} = 0$
122. Equation of the straight line perpendicular to the line $x - y + 5 = 0$, through the point of intersection the y-axis and the given line
 1. $x - y - 5 = 0$ 2. $x + y - 5 = 0$ 3. $x + y + 5 = 0$ 4. $x + y + 10 = 0$
123. If the equation of the base opposite to the vertex $(2, 3)$ of an equilateral triangle is $x + y = 2$, then the length of a side is
 1. $\frac{\sqrt{3}}{2}$ 2. 6 3. $\sqrt{6}$ 4. $3\sqrt{2}$
124. The line $(p + 2q)x + (p - 3q)y = p - q$ for different values of p and q passes through the point
 1. $(\frac{3}{2}, \frac{5}{2})$ 2. $(\frac{2}{5}, \frac{2}{5})$ 3. $(\frac{3}{5}, \frac{3}{5})$ 4. $(\frac{2}{5}, \frac{3}{5})$
125. The point on the line $2x - 3y = 5$ is equidistant from $(1, 2)$ and $(3, 4)$ is
 1. $(7, 3)$ 2. $(4, 1)$ 3. $(1, -1)$ 4. $(-2, 3)$
126. The image of the point $(2, 3)$ in the line $y = -x$ is
 1. $(-3, -2)$ 2. $(-3, 2)$ 3. $(-2, -3)$ 4. $(3, 2)$
127. The length of \perp from the origin to the line $\frac{x}{3} - \frac{y}{4} = 1$, is
 1. $\frac{11}{5}$ 2. $\frac{5}{12}$ 3. $\frac{12}{5}$ 4. $-\frac{5}{12}$
128. The y-intercept of the straight line passing through $(1, 3)$ and perpendicular to $2x - 3y + 1 = 0$ is
 1. $\frac{3}{2}$ 2. $\frac{9}{2}$ 3. $\frac{2}{3}$ 4. $\frac{2}{9}$
129. If the two straight lines $x + (2k - 7)y + 3 = 0$ and $3kx + 9y - 5 = 0$ are perpendicular then the value of k is
 1. $k = 3$ 2. $k = \frac{1}{3}$ 3. $k = \frac{2}{3}$ 4. $k = \frac{3}{2}$
130. If a vertex of a square is at the origin and its one side lies along the line $4x + 3y - 2 = 0$, then the area of the square is
 1. 20 sq. units 2. 16 sq. units 3. 25 sq. units 4. 4 sq. units
131. If the lines represented by the equation $6x^2 + 41xy - 7y^2 = 0$ make angles α and β with x-axis, then $\tan \alpha \tan \beta =$

ONE MARKS

XI – MATHEMATICS

1. $-\frac{6}{7}$

2. $\frac{6}{7}$

3. $-\frac{7}{6}$

4. $\frac{7}{6}$

132. The area of the triangle formed by the lines $x^2 - 4y^2 = 0$ and $x = a$ is

1. $2a^2$

2. $\frac{\sqrt{3}}{2}a^2$

3. $\frac{1}{2}a^2$

4. $\frac{2}{\sqrt{3}}a^2$

133. If one of the lines given by $6x^2 - xy + 4cy^2 = 0$ is $3x + 4y = 0$, then c equals to

1. -3

2. -1

3. 3

4. 1

134. θ is acute angle between the lines $x^2 - xy - 6y^2 = 0$, the $\frac{2\cos\theta + 3\sin\theta}{4\sin\theta + 5\cos\theta}$ is

1. 1

2. $-\frac{1}{9}$

3. $\frac{5}{9}$

4. $\frac{1}{9}$

135. The equation of one the line represented by the equation $x^2 + 2xy \cot \theta - y^2 = 0$ is

1. $x - y \cot \theta = 0$

2. $x + y \tan \theta = 0$

3. $x \cos \theta + y(\sin \theta + 1) = 0$

4. $x \sin \theta + y(\cos \theta + 1) = 0$

Answer key

1. Sets, Relation and Functions

1	2	3	4	5	6	7	8	9	10
3	2	4	1	1	4	2	2	3	2
11	12	13	14	15	16	17	18	19	20
2	3	3	2	4	3	4	3	2	4
21	22	23	24	25					
4	1	4	2	3					

2. Basic algebra

26	27	28	29	30	31	32	33	34	35
2	1	1	3	2	2	3	1	2	2
36	37	38	39	40	41	42	43	44	45
2	3	3	2	3	1	3	1	1	4

3. Trigonometry

46	47	48	49	50	51	52	53	54	55
4	1	1	1	4	4	1	2	4	2
56	57	58	59	60	61	62	63	64	65
3	2	3	3	2	4	1	1	1	1

ONE MARKS

XI – MATHEMATICS

4. Combinatorics and Mathematical Induction

66	67	68	69	70	71	72	73	74	75
2	2	3	2	4	2	1	4	2	2
76	77	78	79	80	81	82	83	84	85
3	4	2	3	1	4	2	3	4	3
86	87	88	89	90					
2	2	2	1	2					

5. Binomial Theorem, Sequences and Series

91	92	93	94	95	96	97	98	99	100
4	4	4	4	2	3	1	3	4	1
101	102	103	104	105	106	107	108	109	110
1	4	4	2	3	2	2	3	3	2

6. Two Dimensional Analytical Geometry

111	112	113	114	115	116	117	118	119	120
4	4	3	4	3	2	2	4	2	3
121	122	123	124	125	126	127	128	129	130
1	2	3	4	2	1	3	2	1	2
131	132	133	134	135					
1	3	1	3	4					

7. MATRICES AND DETERMINANTS

- If $a_{ij} = \frac{1}{2}(3i - 2j)$ and $A = [a_{ij}]_{2 \times 2}$ is
 - $\begin{bmatrix} \frac{1}{2} & 2 \\ -\frac{1}{2} & 1 \end{bmatrix}$
 - $\begin{bmatrix} \frac{1}{2} & -\frac{1}{2} \\ 2 & 1 \end{bmatrix}$
 - $\begin{bmatrix} 2 & 2 \\ \frac{1}{2} & -\frac{1}{2} \end{bmatrix}$
 - $\begin{bmatrix} -\frac{1}{2} & \frac{1}{2} \\ 1 & 2 \end{bmatrix}$
- What must be the matrix X, if $2X + \begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix} = \begin{bmatrix} 3 & 8 \\ 7 & 2 \end{bmatrix}$?
 - $\begin{bmatrix} 1 & 3 \\ 2 & -1 \end{bmatrix}$
 - $\begin{bmatrix} 1 & -3 \\ 2 & -1 \end{bmatrix}$
 - $\begin{bmatrix} 2 & 6 \\ 4 & -2 \end{bmatrix}$
 - $\begin{bmatrix} 2 & -6 \\ 4 & -2 \end{bmatrix}$
- Which one of the following is not true about the matrix $\begin{bmatrix} 1 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 5 \end{bmatrix}$?
 - a scalar matrix
 - a diagonal matrix
 - an upper triangular matrix
 - a lower triangular matrix
- If A and B are two matrices such that A + B and AB are both defined, then
 - A and B are two matrices not necessarily of same order
 - A and B are square matrices of same order
 - Number of columns of A is equal to the number of rows of B
 - A = B
- If $A = \begin{bmatrix} \lambda & 1 \\ -1 & -\lambda \end{bmatrix}$, then for what value of λ , $A^2 = O$?
 - 0
 - ± 1
 - 1
 - 1
- If $A = \begin{bmatrix} 1 & -1 \\ 2 & -1 \end{bmatrix}$, $B = \begin{bmatrix} a & 1 \\ b & -1 \end{bmatrix}$ and $(A + B)^2 = A^2 + B^2$, then the values of a and b are
 - a = 4, b = 1
 - a = 1, b = 4
 - a = 0, b = 4
 - a = 2, b = 4
- If $A = \begin{bmatrix} 1 & 2 & 2 \\ 2 & 1 & -2 \\ a & 2 & b \end{bmatrix}$ is a matrix satisfying the equation $AA^T = 9I$, where I is 3 x 3 identity matrix, then the ordered pair (a, b) is equal to
 - (2, -1)
 - (-2, 1)
 - (2, 1)
 - (-2, -1)
- If A is a square matrix, then which of the following is not symmetric?
 - $A + A^T$
 - AA^T
 - $A^T A$
 - $A - A^T$
- If A and B are symmetric matrices of order n, where $(A \neq B)$, then
 - $A + B$ is skew - symmetric
 - $A + B$ is symmetric
 - $A + B$ is a diagonal matrix
 - $A + B$ is a zero matrix
- If $A = \begin{bmatrix} a & x \\ y & a \end{bmatrix}$ and if $xy = 1$, then $\det(A A^T)$ is equal to
 - $(a - 1)^2$
 - $(a^2 + 1)^2$
 - $a^2 - 1$
 - $(a^2 - 1)^2$
- The value of x, for which the matrix $A = \begin{bmatrix} e^{x-2} & e^{7+x} \\ e^{2+x} & e^{2x+3} \end{bmatrix}$ is singular is
 - 9
 - 8
 - 7
 - 6
- If the points (x, -2), (5, 2), (8, 8) are collinear, then x is equal to
 - 3
 - $\frac{1}{3}$
 - 1
 - 3
- If $\begin{bmatrix} 2a & x_1 & y_1 \\ 2b & x_2 & y_2 \\ 2c & x_3 & y_3 \end{bmatrix} = \frac{abc}{2} \neq 0$, then the area of the triangle whose vertices are $\left(\frac{x_1}{a}, \frac{y_1}{a}\right), \left(\frac{x_2}{b}, \frac{y_2}{b}\right), \left(\frac{x_3}{c}, \frac{y_3}{c}\right)$ is

QUESTIONS BANK

XI - MATHS

1. $\frac{1}{4}$

2. $\frac{1}{4} abc$

3. $\frac{1}{8}$

4. $\frac{1}{8} abc$

14. If the square of the matrix $\begin{bmatrix} \alpha & \beta \\ \gamma & -\alpha \end{bmatrix}$ is the unit matrix of order 2, then α, β and γ should satisfy the relation.

1. $1 + \alpha^2 + \beta\gamma = 0$

2. $1 - \alpha^2 - \beta\gamma = 0$

3. $1 - \alpha^2 + \beta\gamma = 0$

4. $1 + \alpha^2 - \beta\gamma = 0$

15. If $\Delta = \begin{vmatrix} a & b & c \\ x & y & z \\ p & q & r \end{vmatrix}$, then $\begin{vmatrix} ka & kb & kc \\ kx & ky & kz \\ kp & kq & kr \end{vmatrix}$ is

1. Δ

2. $k\Delta$

3. $3k\Delta$

4. $k^3\Delta$

16. A root of the equation $\begin{vmatrix} 3-x & -6 & 3 \\ -6 & 3-x & 3 \\ 3 & 3 & -6-x \end{vmatrix} = 0$ is

1. 6

2. 3

3. 0

4. -6

17. The value of the determinant of $A = \begin{bmatrix} 0 & a & -b \\ -a & 0 & c \\ b & -c & 0 \end{bmatrix}$ is

1. $-2abc$

2. abc

3. 0

4. $a^2 + b^2 + c^2$

18. If x_1, x_2, x_3 as well as y_1, y_2, y_3 , are in geometric progression with the same common ratio then the points $(x_1, y_1), (x_2, y_2), (x_3, y_3)$ are

1. Vertices of an equilateral triangle

2. Vertices of a right angled triangle

3. Vertices of a right angled isosceles triangle

4. collinear

19. If $[.]$ denotes the greatest integer less than or equal to the real number under consideration and $-1 \leq x < 0, 0 \leq y < 1, 1 \leq z < 2$, then the value of the

determinant $\begin{vmatrix} [x]+1 & [y] & [z] \\ [x] & [y]+1 & [z] \\ [x] & [y] & [z]+1 \end{vmatrix}$

1. $[z]$

2. $[y]$

3. $[x]$

4. $[x] + 1$

20. If $a \neq b, b, c$ satisfy $\begin{vmatrix} a & 2b & 2c \\ 3 & b & c \\ 4 & a & b \end{vmatrix} = 0$, then $abc =$

1. $a + b + c$

2. 0

3. b^3

4. $ab + bc$

21. If $A = \begin{bmatrix} -1 & 2 & 4 \\ 3 & 1 & 0 \\ -2 & 4 & 2 \end{bmatrix}$ and $B = \begin{bmatrix} -2 & 4 & 2 \\ 6 & 2 & 0 \\ -2 & 4 & 8 \end{bmatrix}$, then B is given by

1. $B = 4A$

2. $B = -4A$

3. $B = -A$

4. $B = 6A$

22. If A is skew-symmetric of order n and C is a column matrix of order $n \times 1$, then $C^T AC$ is

1. an identity matrix of order n

2. an identity matrix of order 1

3. a zero matrix of order 1

4. an identity matrix of order 2

23. The matrix A satisfying the equation $\begin{bmatrix} 1 & 3 \\ 0 & 1 \end{bmatrix} A = \begin{bmatrix} 1 & 1 \\ 0 & -1 \end{bmatrix}$ is

1. $\begin{bmatrix} 1 & 4 \\ -1 & 0 \end{bmatrix}$

2. $\begin{bmatrix} 1 & -4 \\ 1 & 0 \end{bmatrix}$

3. $\begin{bmatrix} 1 & 4 \\ 0 & -1 \end{bmatrix}$

4. $\begin{bmatrix} 1 & -4 \\ 1 & 1 \end{bmatrix}$

24. If $A + I = \begin{bmatrix} 3 & -2 \\ 4 & 1 \end{bmatrix}$, then $(A + I)(A - I)$ is equal to

1. $\begin{bmatrix} -5 & -4 \\ 8 & -9 \end{bmatrix}$

2. $\begin{bmatrix} -5 & 4 \\ -8 & 9 \end{bmatrix}$

3. $\begin{bmatrix} 5 & 4 \\ 8 & 9 \end{bmatrix}$

4. $\begin{bmatrix} -5 & -4 \\ -8 & -9 \end{bmatrix}$

QUESTIONS BANK

XI - MATHS

25. Let A and B be two symmetric matrices of same order. Then which one of the following statement is not true?

1. $A + B$ is a symmetric matrix
2. AB is a symmetric matrix
3. $AB = (BA)^T$
4. $A^T B = AB^T$

8. VECTOR ALGEBRA

26. The value of $\overrightarrow{AB} + \overrightarrow{BC} + \overrightarrow{DA} + \overrightarrow{CD}$ is

1. \overrightarrow{AD}
2. \overrightarrow{CA}
3. $\vec{0}$
4. $-\overrightarrow{AD}$

27. If $\vec{a} + 2\vec{b}$ and $3\vec{a} + m\vec{b}$ are parallel, then the value of m is

1. 3
2. $\frac{1}{3}$
3. 6
4. $\frac{1}{6}$

28. The unit vector parallel to the resultant of the vectors $\hat{i} + \hat{j} - \hat{k}$ and $\hat{i} - 2\hat{j} + \hat{k}$ is

1. $\frac{\hat{i} - \hat{j} + \hat{k}}{\sqrt{5}}$
2. $\frac{2\hat{i} + \hat{j}}{\sqrt{5}}$
3. $\frac{2\hat{i} - \hat{j} + \hat{k}}{\sqrt{5}}$
4. $\frac{2\hat{i} - \hat{j}}{\sqrt{5}}$

29. A vector \overrightarrow{OP} makes 60° and 45° with the positive direction of the x and y axes respectively. Then the angle between \overrightarrow{OP} and the z-axis is

1. 45°
2. 60°
3. 90°
4. 30°

30. If $\overrightarrow{BA} = 3\hat{i} + 2\hat{j} + \hat{k}$ and the position vector B is $\hat{i} + 3\hat{j} - \hat{k}$ then the position vector A is

1. $4\hat{i} + 2\hat{j} + \hat{k}$
2. $4\hat{i} + 5\hat{j}$
3. $4\hat{i}$
4. $-4\hat{i}$

31. A vector makes equal angle with the positive direction of the coordinate axes. Then each angle is equal to

1. $\cos^{-1}\left(\frac{1}{3}\right)$
2. $\cos^{-1}\left(\frac{2}{3}\right)$
3. $\cos^{-1}\left(\frac{1}{\sqrt{3}}\right)$
4. $\cos^{-1}\left(\frac{2}{\sqrt{3}}\right)$

32. The vectors $\vec{a} - \vec{b}$, $\vec{b} - \vec{c}$, $\vec{c} - \vec{a}$ are

1. parallel to each other
2. unit vectors
3. mutually perpendicular vectors
4. coplanar vectors.

33. If ABCD is a parallelogram, then $\overrightarrow{AB} + \overrightarrow{AD} + \overrightarrow{CB} + \overrightarrow{CD}$ is equal to

1. $2(\overrightarrow{AB} + \overrightarrow{AD})$
2. $4\overrightarrow{AC}$
3. $4\overrightarrow{BD}$
4. $\vec{0}$

34. One of the diagonals of parallelogram ABCD with

\vec{a} and \vec{b} as adjacent sides is $\vec{a} + \vec{b}$. The other diagonal \overrightarrow{BD} is

1. $\vec{a} - \vec{b}$
2. $\vec{b} - \vec{a}$
3. $\vec{a} + \vec{b}$
4. $\frac{\vec{a} - \vec{b}}{3}$

35. If \vec{a} , \vec{b} , \vec{c} are the position vectors A and B, then which one of the following points whose position vector lies on AB, is

1. $\vec{a} + \vec{b}$
2. $\frac{2\vec{a} - \vec{b}}{2}$
3. $\frac{2\vec{a} + \vec{b}}{3}$
4. $\frac{\vec{a} + \vec{b}}{2}$

36. If \vec{a} , \vec{b} , \vec{c} are the position vectors of three collinear points, then which of the following is true?

1. $\vec{a} = \vec{b} + \vec{c}$
2. $2\vec{a} = \vec{b} + \vec{c}$
3. $\vec{b} = \vec{c} + \vec{a}$
4. $4\vec{a} + \vec{b} + \vec{c} = \vec{0}$

37. If $\vec{r} = \frac{9\vec{a} + 7\vec{b}}{16}$, then the point P whose position vector \vec{r} divides the line joining the points with position vectors \vec{a} and \vec{b} in the ratio

1. 7 : 9 internally
2. 9:7 internally
3. 9:7 externally
4. 7 : 9 externally

QUESTIONS BANK

XI - MATHS

38. If $\lambda \vec{i} + 2\lambda \vec{j} + 2\lambda \vec{k}$ is a unit vector, then the value of λ is
 1. $\frac{1}{3}$ 2. $\frac{1}{4}$ 3. $\frac{1}{9}$ 4. $\frac{1}{2}$
39. Two vertices of a triangle have position vectors $3\hat{i} + 4\hat{j} - 4\hat{k}$ and $2\hat{i} + 3\hat{j} + 4\hat{k}$. If the position vector of the centroid is $\hat{i} + 2\hat{j} + 3\hat{k}$, then the position vector of the third vertex is
 1. $-2\hat{i} - \hat{j} + 9\hat{k}$ 2. $-2\hat{i} - \hat{j} - 6\hat{k}$ 3. $2\hat{i} - \hat{j} + 6\hat{k}$ 4. $-2\hat{i} + \hat{j} + 6\hat{k}$
40. If $|\vec{a} + \vec{b}| = 60, |\vec{a} - \vec{b}| = 40$ and $|\vec{b}| = 46$, then $|\vec{a}|$ is
 1. 42 2. 12 3. 22 4. 32
41. If \vec{a} and \vec{b} having same magnitude and angle between them is 60° and their scalar product is $\frac{1}{2}$ then $|\vec{a}|$ is
 1. 2 2. 3 3. 7 4. 1
42. The value of $\theta \in (0, \frac{\pi}{2})$ for which the vectors $\vec{a} = (\sin \theta)\hat{i} + (\cos \theta)\hat{j}$ and $\vec{b} = \hat{i} - \sqrt{3}\hat{j} + 2\hat{k}$ are perpendicular, is equal to
 1. $\frac{\pi}{3}$ 2. $\frac{\pi}{6}$ 3. $\frac{\pi}{4}$ 4. $\frac{\pi}{2}$
43. If $|\vec{a}| = 13, |\vec{b}| = 5$ and $\vec{a} \cdot \vec{b} = 60^\circ$ then $|\vec{a} \times \vec{b}|$ is
 1. 15 2. 35 3. 45 4. 25
44. Vectors \vec{a} and \vec{b} are inclined at an angle $\theta = 120^\circ$. If $|\vec{a}| = 1, |\vec{b}| = 2$ then $[(\vec{a} + 3\vec{b}) \times (3\vec{a} - \vec{b})]^2$ is equal to
 1. 225 2. 275 3. 325 4. 300
45. If \vec{a} and \vec{b} are two vectors of magnitude 2 and inclined at an angle 60° , then the angle \vec{a} and $\vec{a} + \vec{b}$ is
 1. 30° 2. 60° 3. 45° 4. 90°
46. If the projection of $5\hat{i} - \hat{j} - 3\hat{k}$ on the vector $\hat{i} + 3\hat{j} + \lambda\hat{k}$ is same as the projection of $\hat{i} + 3\hat{j} + \lambda\hat{k}$ on $5\hat{i} - \hat{j} - 3\hat{k}$, then λ is equal to
 1. ± 4 2. ± 3 3. ± 5 4. ± 1
47. If $(1, 2, 4)$ and $(2, -3\lambda, -3)$ are the initial and terminal points of the vector $\hat{i} + 5\hat{j} - 7\hat{k}$, then the value of λ is equal to
 1. $\frac{7}{3}$ 2. $-\frac{7}{3}$ 3. $-\frac{5}{3}$ 4. $\frac{5}{3}$
48. If the points whose position vectors $10\hat{i} + 3\hat{j}$, $12\hat{i} - 5\hat{j}$ and $a\hat{i} + 11\hat{j}$ are collinear then a is equal to
 1. 6 2. 3 3. 5 4. 8
49. If $\vec{a} = \hat{i} + \hat{j} + \hat{k}, \vec{b} = 2\hat{i} + x\hat{j} + \hat{k}, \vec{c} = \hat{i} - \hat{j} + 4\hat{k}$ and $\vec{a} \cdot (\vec{b} + \vec{c}) = 70$, then x is equal
 1. 5 2. 7 3. 26 4. 10
50. If $\vec{a} = \hat{i} + 2\hat{j} + 2\hat{k}, |\vec{b}| = 5$ and the angle between \vec{a} and \vec{b} is $\frac{\pi}{6}$, then the area of the triangle formed by these two vectors as two sides, is
 1. $\frac{7}{4}$ 2. $\frac{15}{4}$ 3. $\frac{3}{4}$ 4. $\frac{17}{4}$

XI - MATHS

51. $\lim_{x \rightarrow \infty} \frac{\sin x}{x}$
 1. 1 2. 0 3. ∞ 4. $-\infty$

52. $\lim_{x \rightarrow \frac{\pi}{2}} \frac{2x - \pi}{\cos x}$
 1. 2 2. 1 3. -2 4. 0

53. $\lim_{x \rightarrow 0} \frac{\sqrt{1 - \cos 2x}}{x}$
 1. 0 2. 1 3. $\sqrt{2}$ 4. does not exist

54. $\lim_{\theta \rightarrow 0} \frac{\sin \sqrt{\theta}}{\sqrt{\sin \theta}}$
 1. 1 2. -1 3. 0 4. 2

55. $\lim_{x \rightarrow \infty} \left(\frac{x^2 + 5x + 3}{x^2 + x + 3} \right)^x$ is
 1. e^4 2. e^2 3. e^3 4. 1

56. $\lim_{x \rightarrow \infty} \frac{\sqrt{x^2 - 1}}{2x + 1} =$
 1. 1 2. 0 3. -1 4. $\frac{1}{2}$

57. $\lim_{x \rightarrow \infty} \frac{a^x - b^x}{x} =$
 1. $\log ab$ 2. $\log \left(\frac{a}{b} \right)$ 3. $\log \left(\frac{b}{a} \right)$ 4. $\left(\frac{a}{b} \right)$

58. $\lim_{x \rightarrow 0} \frac{8^x - 4^x - 2^x + 1^x}{x^2} =$
 1. $2 \log 2$ 2. $2(\log 2)^2$ 3. $\log 2$ 4. $3 \log 2$

59. If $f(x) = x(-1)^{\lfloor \frac{1}{|x|} \rfloor}$, $x \leq 0$, then the value of $\lim_{x \rightarrow 0} f(x)$ is equal to
 1. -1 2. 0 3. 2 4. 4

60. $\lim_{x \rightarrow 3} [x] =$
 1. 2 2. 3 3. does not exist 4. 0

61. Let the function f be defined by $f(x) = \begin{cases} 3x, & 0 \leq x \leq 1 \\ -3x + 5, & 1 < x \leq 2 \end{cases}$, then
 1. $\lim_{x \rightarrow 1} f(x) = 1$ 2. $\lim_{x \rightarrow 1} f(x) = 3$ 3. $\lim_{x \rightarrow 1} f(x) = 2$ 4. $\lim_{x \rightarrow 1} f(x)$ does not exist

62. If $f : \mathbb{R} \rightarrow \mathbb{R}$ is defined by $f(x) = [x - 3] + [x - 4]$ for $x \in \mathbb{R}$, then $\lim_{x \rightarrow 3^-} f(x)$ is equal to
 1. -2 2. -1 3. 0 4. 1

63. $\lim_{x \rightarrow 0} \frac{xe^x - \sin x}{x}$ is
 1. 1 2. 2 3. 3 4. 0

64. If $\lim_{x \rightarrow 0} \frac{\sin px}{\tan 3x} = 4$, then the value of p is
 1. 6 2. 9 3. 12 4. 4

65. $\lim_{\alpha \rightarrow \frac{\pi}{4}} \frac{\sin \alpha - \cos \alpha}{\alpha - \frac{\pi}{4}}$
 1. $\sqrt{2}$ 2. $\frac{1}{\sqrt{2}}$ 3. 1 4. 2

66. $\lim_{n \rightarrow \infty} \left(\frac{1}{n^2} + \frac{2}{n^2} + \frac{3}{n^2} + \dots + \frac{n}{n^2} \right)$ is
 1. $\frac{1}{2}$ 2. 0 3. 1 4. ∞

67. $\lim_{x \rightarrow 0} \frac{e^{\sin x} - 1}{x}$
 1. 1 2. e 3. $\frac{1}{e}$ 4. 0

68. $\lim_{x \rightarrow 0} \frac{e^{\tan x} - e^x}{\tan x - x} =$
 1. 1 2. e 3. $\frac{1}{2}$ 4. 0

QUESTIONS BANK

XI - MATHS

69. The value of $\lim_{x \rightarrow 0} \frac{\sin x}{\sqrt{x^2}}$ is
 1. 1 2. -1 3. 0 4. ∞
70. The value of $\lim_{x \rightarrow k^-} x - [x]$, where k is an integer is
 1. -1 2. 1 3. 0 4. 2
71. At $x = \frac{3}{2}$ the function $f(x) = \frac{|2x-3|}{2x-3}$ is
 1. Continuous 2. discontinuous 3. differentiable 4. non-zero
72. Let $f: \mathbb{R} \rightarrow \mathbb{R}$ be defined by $f(x) = \begin{cases} x & x \text{ is irrational} \\ 1-x, & x \text{ is rational} \end{cases}$ then f is
 1. discontinuous at $x = \frac{1}{2}$ 2. continuous at $x = \frac{1}{2}$
 3. continuous everywhere 4. discontinuous everywhere
73. The function $f(x) = \begin{cases} \frac{x^2-1}{x^3+1}, & x \neq -1 \\ p, & x = -1 \end{cases}$ is not defined for $x = -1$. The value of f(-1) so that the function extended by this value is continuous is
 1. $\frac{2}{3}$ 2. $-\frac{2}{3}$ 3. 1 4. 0
74. Let f be a continuous function on $[2, 5]$. If takes only rational values for all x and $f(3) = 12$ then $f(4.5)$ is equal to
 1. $\frac{f(3)+f(4.5)}{7.5}$ 2. 12 3. 17.5 4. $\frac{f(4.5)-f(3)}{1.5}$
75. Let a function f be defined by $f(x) = \frac{x-|x|}{x}$ for $x \neq 0$ and $f(0) = 2$. Then f is
 1. continuous nowhere 2. continuous everywhere
 3. continuous for all x except $x = 1$ 4. continuous for all x except $x = 0$.

10. Differential Calculus

76. $\frac{d}{dx} \left(\frac{2}{\pi} \sin x^\circ \right)$ is
 1. $\frac{\pi}{180} \sin x^\circ$ 2. $\frac{1}{90} \cos x^\circ$ 3. $\frac{\pi}{90} \cos x^\circ$ 4. $\frac{2}{\pi} \cos x^\circ$
77. If $y = f(x^2 + 2)$ and $f'(3) = 5$, then $\frac{dy}{dx}$ at $x = 1$ is
 1. 5 2. 25 3. 15 4. 10
78. If $y = \frac{1}{4}u^4$, $u = \frac{2}{3}x^3 + 5$, then $\frac{dy}{dx}$ is
 1. $\frac{1}{27}x^2(2x^3 + 15)^3$ 2. $\frac{2}{27}x(2x^3 + 5)^3$ 3. $\frac{2}{27}x^2(2x^3 + 15)^3$ 4. $-\frac{2}{27}x(2x^3 + 5)^3$
79. If $f(x) = x^2 - 3x$, then the point at which $f(x) = f'(x)$ are
 1. both positive integers 2. both negative integers
 3. both irrational 4. one rational and another irrational
80. If $y = \frac{1}{a-z}$, then $\frac{dz}{dy}$ is
 1. $(a-z)^2$ 2. $-(z-a)^2$ 3. $(z+a)^2$ 4. $-(z+a)^2$
81. If $y = \cos(\sin x^2)$, then $\frac{dy}{dx}$ at $x = \sqrt{\frac{\pi}{2}}$ is
 1. -2 2. 2 3. $-2\sqrt{\frac{\pi}{2}}$ 4. 0
82. If $y = mx + c$ and $f(0) = f'(0) = 1$, then $f(2)$ is
 1. 1 2. 2 3. 3 4. -3
83. If $f(x) = x \tan^{-1} x$, then $f'(1)$ is
 1. $1 + \frac{\pi}{4}$ 2. $\frac{1}{2} + \frac{\pi}{4}$ 3. $\frac{1}{2} - \frac{\pi}{4}$ 4. 2

QUESTIONS BANK

XI - MATHS

84. $\frac{d}{dx}(e^{x+5\log x})$ is

1. $e^x \cdot x^4(x+5)$ 2. $e^x \cdot x(x+5)$ 3. $e^x + \frac{5}{x}$ 4. $e^x - \frac{5}{x}$

85. If the derivative of $(ax - 5)e^{3x}$ at $x = 0$ is -13 , then the value of a is

1. 8 2. -2 3. 5 4. 2

86. $x = \frac{1-t^2}{1+t^2}$, $y = \frac{2t}{1+t^2}$ then $\frac{dy}{dx}$ is

1. $-\frac{y}{x}$ 2. $\frac{y}{x}$ 3. $-\frac{x}{y}$ 4. $\frac{x}{y}$

87. If $x = a \sin \theta$ and $y = b \cos \theta$, then $\frac{d^2y}{dx^2}$ is

1. $\frac{a}{b^2} \sec^2 \theta$ 2. $-\frac{b}{a} \sec^2 \theta$ 3. $\frac{b}{a^2} \sec^3 \theta$ 4. $\frac{b^2}{a^2} \sec^3 \theta$

88. The differential coefficient of $\log_{10} x$ with respect to $\log_x 10$ is

1. 1 2. $-(\log_{10} x)^2$ 3. $(\log_x 10)^2$ 4. $\frac{x^2}{100}$

89. If $f(x) = x + 2$, then $f'(f(x))$ at $x = 4$ is

1. 8 2. 1 3. 4 4. 5

90. If $y = \frac{(1-x)^2}{x^2}$, then $\frac{dy}{dx}$ is

1. $\frac{2}{x^2} + \frac{2}{x^3}$ 2. $-\frac{2}{x^2} + \frac{2}{x^3}$ 3. $-\frac{2}{x^2} - \frac{2}{x^3}$ 4. $-\frac{2}{x^3} + \frac{2}{x^2}$

91. If $pv = 81$, then $\frac{dp}{dv}$ at $v = 9$ is

1. 1 2. -1 3. 2 4. -2

92. If $f(x) = \begin{cases} x-5 & \text{if } x \leq 1 \\ 4x^2-9 & \text{if } 1 < x < 2 \\ 3x+4 & \text{if } x \geq 2 \end{cases}$, then the right hand derivative of $f(x)$ at $x = 2$ is

1. 0 2. 2 3. 3 4. 4

93. It is given that $f'(a)$ exists, then $\lim_{x \rightarrow a} \frac{xf(a) - af(x)}{x-a}$ is

1. $f(a) - af'(a)$ 2. $f'(a)$ 3. $-f'(a)$ 4. $f(a) + af'(a)$

94. If $f(x) = \begin{cases} x+1, & \text{when } x < 2 \\ 2x-1 & \text{when } x \geq 2 \end{cases}$ then $f'(2)$ is

1. 0 2. 1 3. 2 4. does not exist

95. If $g(x) = (x^2 + 2x + 3)f(x)$ and $f(0) = 5$ and $\lim_{x \rightarrow 0} \frac{f(x)-5}{x} = 4$, then $g'(0)$ is

1. 20 2. 14 3. 18 4. 12

96. If $f(x) = \begin{cases} x+2 & -1 < x < 3 \\ 5 & x = 3 \\ 8-x & x > 3 \end{cases}$, then at $x = 3$, $f'(x)$ is

1. 1 2. -1 3. 0 4. does not exist

97. The derivative of $f(x) = x|x|$ at $x = -3$ is

1. 6 2. -6 3. does not exist 4. 0

98. If $f(x) = \begin{cases} 2a-x, & \text{for } -a < x < a \\ 3x-2a & \text{for } x \geq a \end{cases}$, then which one of the following is true?

1. $f(x)$ is not differentiable at $x = a$ 2. $f(x)$ is discontinuous at $x = a$
3. $f(x)$ is continuous for all x in \mathbb{R} 4. $f(x)$ is differentiable for all $x \geq a$

99. If $f(x) = \begin{cases} ax^2 - b & -1 < x < 1 \\ \frac{1}{|x|} & \text{elsewhere} \end{cases}$, is differentiable at $x = 1$, then

1. $a = \frac{1}{2}, b = \frac{-3}{2}$ 2. $a = \frac{-1}{2}, b = \frac{3}{2}$ 3. $a = -\frac{1}{2}, b = -\frac{3}{2}$ 4. $a = \frac{1}{2}, b = \frac{3}{2}$

QUESTIONS BANK

XI - MATHS

100. The number of points in \mathbb{R} in which the function $f(x) = |x - 1| + |x - 3| + \sin x$ is not differentiable, is

1. 3

2. 2

3. 1

4. 4

11. Integral Calculus

101. If $\int f(x)dx = g(x) + c$, then $\int f(x)g'(x)dx$

1. $\int f(x))^2 dx$ 2. $\int f(x)g(x) dx$ 3. $\int f'(x) g(x) dx$ 4. $\int (g(x))^2 dx$

102. If $\int \frac{3^x}{x^2} dx = k(3^{\frac{1}{x}}) + c$, then the value of k is

1. $\log 3$ 2. $-\log 3$ 3. $-\frac{1}{\log 3}$ 4. $\frac{1}{\log 3}$

103. If $\int f'(x) e^{x^2} dx = (x - 1) e^{x^2} + c$, then $f(x)$ is

1. $2x^3 - \frac{x^2}{2} + x + c$ 2. $\frac{x^2}{2} + 3x^2 + 4x + c$ 3. $x^3 + 4x^2 + 6x + c$ 4. $\frac{2x^3}{3} - x^2 + x + c$

104. The gradient (slope) of a curve at any point (x, y) is $\frac{x^2 - 4}{x^2}$. If the curve passes through the point $(2, 7)$, then the equation of the curve is

1. $y = x + \frac{4}{x} + 3$ 2. $y = x + \frac{4}{x} + 4$ 3. $y = x^2 + 3x + 4$ 4. $y = x^2 - 3x + 6$

105. $\int \frac{e^x(1+x)}{\cos^2(xe^x)} dx$ is

1. $\cot(xe^x) + c$ 2. $\sec(xe^x) + c$ 3. $\tan(xe^x) + x$ 4. $\cos(xe^x) + c$

106. $\int \frac{\sqrt{\tan x}}{\sin 2x} dx$ is

1. $\sqrt{\tan x} + c$ 2. $2\sqrt{\tan x} + c$ 3. $\frac{1}{2}\sqrt{\tan x} + c$ 4. $\frac{1}{4}\sqrt{\tan x} + c$

107. $\int \sin^3 x dx$ is

1. $\frac{-3}{4} \cos x - \frac{\cos 3x}{12} + c$ 2. $\frac{3}{4} \cos x + \frac{\cos 3x}{12} + c$ 3. $\frac{-3}{4} \cos x + \frac{\cos 3x}{12} + c$ 4. $\frac{-3}{4} \sin x - \frac{\sin 3x}{12} + c$

108. $\int \frac{e^{6 \log x} - e^{5 \log x}}{e^{4 \log x} - e^{3 \log x}} dx$ is

1. $x + c$ 2. $\frac{x^3}{3} + c$ 3. $\frac{3}{x^3} + c$ 4. $\frac{1}{x^2} + c$

109. $\int \frac{\sec x}{\sqrt{\cos 2x}} dx$ is

1. $\tan^{-1}(\sin x) + c$ 2. $2 \sin^{-1}(\tan x) + c$ 3. $\tan^{-1}(\cos x) + c$ 4. $\sin^{-1}(\tan x) + c$

110. $\int \tan^{-1} \sqrt{\frac{1 - \cos 2x}{1 + \cos 2x}} dx$ is

1. $x^2 + c$ 2. $2x^2 + c$ 3. $\frac{x^2}{2} + c$ 4. $-\frac{x^2}{2} + c$

111. $\int 2^{3x+5} dx$ is

1. $\frac{3(2^{3x+5})}{\log 2} + c$ 2. $\frac{2^{3x+5}}{2 \log(3x+5)} + c$ 3. $\frac{2^{3x+5}}{2 \log 3} + c$ 4. $\frac{2^{3x+5}}{3 \log 2} + c$

QUESTIONS BANK

XI - MATHS

112. $\int \frac{\sin^8 x - \cos^8 x}{1 - 2\sin^2 x \cos^2 x} dx$ is

1. $\frac{1}{2} \sin 2x + c$ 2. $-\frac{1}{2} \sin 2x + c$ 3. $\frac{1}{2} \cos 2x + c$ 4. $-\frac{1}{2} \cos 2x + c$

113. $\int \frac{e^{x(x^2 \tan^{-1} x + \tan^{-1} x + 1)}}{x^2 + 1} dx$ is

1. $\cos x + \sin^{-1} x + c$ 2. $-\cot x + \tan^{-1} x + c$
3. $-\tan x + \cot^{-1} x + c$ 4. $-\cot x - \tan^{-1} x + c$

114. $\int \frac{x^2 + \cos^2 x}{x^2 + 1} \operatorname{cosec}^2 x dx$ is

1. $\cos x + \sin^{-1} x + c$ 2. $-\cot x + \tan^{-1} x + c$
3. $-\tan x + \cot^{-1} x + c$ 4. $-\cot x - \tan^{-1} x + c$

115. $\int x^2 \cos x dx$ is

1. $x^2 \sin x + 2x \cos x - 2 \sin x + c$ 2. $x^2 \sin x - 2x \cos x - 2 \sin x + c$
3. $-x^2 \sin x + 2x \cos x + 2 \sin x + c$ 4. $-x^2 \sin x - 2x \cos x + 2 \sin x + c$

116. $\int \sqrt{\frac{1-x}{1+x}} dx$ is

1. $\sqrt{1-x^2} + \sin^{-1} x + c$ 2. $\sin^{-1} x - \sqrt{1-x^2} + c$
3. $\log |x + \sqrt{1-x^2}| - \sqrt{1-x^2} + c$ 4. $\sqrt{1-x^2} + \log |x + \sqrt{1-x^2}| + c$

117. $\int \frac{dx}{e^x - 1}$ is

1. $\log |e^x| - \log |e^x - 1| + c$ 2. $\log |e^x| - \log |e^x - 1| + c$
3. $-\log |e^x| + \log |e^x - 1| + c$ 4. $\log |1 + e^x| - \log |e^x| + c$

118. $\int e^{-4x} \cos x dx$ is

1. $\frac{e^{-4x}}{17} [4 \cos x - \sin x] + c$ 2. $\frac{e^{-4x}}{17} [-4 \cos x + \sin x] + c$
3. $\frac{e^{-4x}}{17} [4 \cos x + \sin x] + c$ 4. $\frac{e^{-4x}}{17} [-4 \cos x - \sin x] + c$

119. $\int \frac{\sec^2 x}{\tan^2 x - 1} dx$

1. $2 \log \left| \frac{1 - \tan x}{1 + \tan x} \right| + c$ 2. $\log \left| \frac{1 + \tan x}{1 - \tan x} \right| + c$ 3. $\frac{1}{2} \log \left| \frac{\tan x + 1}{\tan x - 1} \right| + c$ 4. $\frac{1}{2} \log \left| \frac{\tan x - 1}{\tan x + 1} \right| + c$

120. $\int e^{-7x} \sin 5x dx$ is

1. $\frac{e^{-7x}}{74} [-7 \sin 5x - 5 \cos 5x] + c$ 2. $\frac{e^{-7x}}{74} [7 \sin 5x + 5 \cos 5x] + c$
3. $\frac{e^{-7x}}{74} [7 \sin 5x - 5 \cos 5x] + c$ 4. $\frac{e^{-7x}}{74} [-7 \sin 5x + 5 \cos 5x] + c$

121. $\int x^2 e^{\frac{x}{2}} dx$ is

1. $x^2 e^{\frac{x}{2}} - 4x e^{\frac{x}{2}} - 8e^{\frac{x}{2}} + c$ 2. $2x^2 e^{\frac{x}{2}} - 8x e^{\frac{x}{2}} - 16e^{\frac{x}{2}} + c$
3. $2x^2 e^{\frac{x}{2}} - 8x e^{\frac{x}{2}} + 16e^{\frac{x}{2}} + c$ 4. $x^2 \frac{e^{\frac{x}{2}}}{2} - \frac{x e^{\frac{x}{2}}}{4} + \frac{e^{\frac{x}{2}}}{8} + c$

QUESTIONS BANK

XI - MATHS

122. $\int \frac{x+2}{\sqrt{x^2-1}} dx$ is

1. $\sqrt{x^2-1} - 2 \log |x + \sqrt{x^2-1}| + c$

2. $\sin^{-1} x - 2 \log |x + \sqrt{x^2-1}| + c$

3. $2 \log |x + \sqrt{x^2-1}| - \sin^{-1} x + c$

4. $\sqrt{x^2-1} + 2 \log |x + \sqrt{x^2-1}| + c$

123. $\int \frac{1}{x\sqrt{(\log x)^2-5}} dx$ is

1. $\log |x + \sqrt{x^2-5}| + c$

2. $\log |\log + \sqrt{\log x - 5}| + c$

3. $\log |\log x + \sqrt{(\log x)^2 - 5}| + c$

4. $\log |\log - \sqrt{(\log x)^2 - 5}| + c$

124. $\int \sin \sqrt{x} dx$ is

1. $2(-\sqrt{x} \cos \sqrt{x} + \sin \sqrt{x}) + c$

2. $2(-\sqrt{x} \cos \sqrt{x} - \sin \sqrt{x}) + c$

3. $2(-\sqrt{x} \sin \sqrt{x} - \cos \sqrt{x}) + c$

4. $2(-\sqrt{x} \sin \sqrt{x} + \cos \sqrt{x}) + c$

125. $\int e^{\sqrt{x}} dx$ is

1. $2\sqrt{x} (1 - e^{\sqrt{x}}) + c$

2. $2\sqrt{x} (e^{\sqrt{x}} - 1) + c$

3. $2e^{\sqrt{x}}(1 - \sqrt{x}) + c$

4. $2e^{\sqrt{x}}(\sqrt{x} - 1) + c$

126. Four persons are selected at random from a group of 3 men, 2 women and 4 children. The probability that exactly two of them are children is

1. $\frac{3}{4}$

2. $\frac{10}{23}$

3. $\frac{1}{2}$

4. $\frac{10}{21}$

127. A number is selected from the set $\{1, 2, 3, \dots, 20\}$. The probability that the selected number is divisible by 3 or 4 is

1. $\frac{2}{5}$

2. $\frac{1}{8}$

3. $\frac{1}{2}$

4. $\frac{2}{3}$

128. A, B, and C try to hit a target simultaneously but independently. Their respective probabilities of hitting the target are $\frac{3}{4}, \frac{1}{2}, \frac{5}{8}$. The probability that the target is hit by A or B but not by C is

1. $\frac{21}{64}$

2. $\frac{7}{32}$

3. $\frac{9}{64}$

4. $\frac{7}{8}$

129. If A and B are any two events, then the probability that exactly one of them occur is

1. $P(A \cup \bar{B}) + P(\bar{A} \cup B)$

2. $P(A \cap \bar{B}) + P(\bar{A} \cap B)$

3. $P(A) + P(B) - P(A \cap B)$

4. $P(A) + P(B) + 2P(A \cap B)$

130. Let A and B be two events such that $\overline{(A \cup B)} = \frac{1}{6}$, $P(A \cap B) = \frac{1}{4}$ and $P(\bar{A}) = \frac{1}{4}$. Then the events A and B are

1. Equally likely but not independent

2. Independent but not equally likely

3. Independent and equally likely

4. Mutually inclusive and dependent

131. Two items are chosen from a lot containing twelve items of which four are defective, then the probability that at least one of the item is defective

1. $\frac{19}{33}$

2. $\frac{17}{33}$

3. $\frac{23}{33}$

4. $\frac{13}{33}$

132. A man has 3 fifty rupee notes, 4 hundred rupees notes and 6 five hundred rupees notes in his pocket. If 2 notes are taken at random, what are the odds in favour of both notes being of hundred rupee denomination?

1. 1:12

2. 12:1

3. 13:1

4. 1:13

QUESTIONS BANK

XI - MATHS

133. A letter is taken at random from the letters of the word 'ASSISTANT' and another letter is taken at random from the letters of the word 'STATISTICS'. The probability that the selected letters are the same is

1. $\frac{7}{45}$ 2. $\frac{17}{90}$ 3. $\frac{29}{90}$ 4. $\frac{19}{90}$

134. A matrix is chosen at random from a set of all matrices of order 2, with elements 0 or 1 only. The probability that the determinant of the matrix chosen is non zero will be

1. $\frac{3}{16}$ 2. $\frac{3}{8}$ 3. $\frac{1}{4}$ 4. $\frac{5}{8}$

135. A bag contains 5 white and 3 black balls. Five balls are drawn successively without replacement. The probability that they are alternately of different colours is

1. $\frac{3}{14}$ 2. $\frac{5}{14}$ 3. $\frac{1}{14}$ 4. $\frac{9}{14}$

136. If A and B are two events such that $A \subset B$ and $P(B) \neq 0$, then which of the following is correct?

1. $P(A/B) = \frac{P(A)}{P(B)}$ 2. $P(A/B) < P(A)$ 3. $P(A/B) \geq P(A)$ 4. $P(A/B) > P(B)$

137. A bag contains 6 green, 2 white, and 7 black balls. If two balls are drawn simultaneously, then the probability that both are different colours is

1. $\frac{68}{105}$ 2. $\frac{71}{105}$ 3. $\frac{64}{105}$ 4. $\frac{73}{105}$

138. If X and Y be two events such that $P(X/Y) = \frac{1}{2}$, $P(Y/X) = \frac{1}{3}$ and $P(X \cap Y) = \frac{1}{6}$, then $P(X \cup Y)$ is

1. $\frac{1}{3}$ 2. $\frac{2}{5}$ 3. $\frac{1}{6}$ 4. $\frac{2}{3}$

139. An urn contains 5 red and 5 black balls. A ball is drawn at random, its colour is noted and is returned to the urn. Moreover, 2 additional balls of the colour drawn are put in the urn and then a ball is drawn at random. The probability that the second ball drawn is red will be

1. $\frac{5}{12}$ 2. $\frac{1}{2}$ 3. $\frac{7}{12}$ 4. $\frac{1}{4}$

140. A number x is chosen at random from the first 100 natural numbers. Let A be the event of numbers which satisfies $\frac{(x-10)(x-50)}{x-30} \geq 0$, then $P(A)$ is

1. 0.20 2. 0.51 3. 0.71 4. 0.70

141. If two events A and B are independent such that $P(A) = 0.35$ and $P(A \cup B) = 0.6$, then $P(B)$ is

1. $\frac{5}{13}$ 2. $\frac{1}{13}$ 3. $\frac{4}{13}$ 4. $\frac{7}{13}$

142. If two events A and B are such that $P(\bar{A}) = \frac{3}{10}$ and $P(A \cap \bar{B}) = \frac{1}{2}$, then $P(A \cap B)$ is

1. $\frac{1}{2}$ 2. $\frac{1}{3}$ 3. $\frac{1}{4}$ 4. $\frac{1}{5}$

143. If A and B are two events such that $P(B) = \frac{1}{2}$ and $P(B/A) = 0.6$, then $P(\bar{A} \cap B)$ is

1. 0.96 2. 0.24 3. 0.56 4. 0.66

144. There are three events A, B and C of which one and only one can happen. If the odds are 7 to 4 against A and 5 to 3 against B, then odds against C is

1. 23: 65 2. 65: 23 3. 23: 88 4. 88: 23

QUESTIONS BANK

XI - MATHS

145. If a and b are chosen randomly from the set $\{1, 2, 3, 4\}$ with replacement, then the probability of the real roots of the equation $x^2 + ax + b = 0$ is
1. $\frac{3}{16}$ 2. $\frac{5}{16}$ 3. $\frac{7}{16}$ 4. $\frac{11}{16}$
146. It is given that the events A and B are such that $P(A) = \frac{1}{4}$, $P(A/B) = \frac{1}{2}$ and $P(B/A) = \frac{2}{3}$. Then P(B) is
1. $\frac{1}{6}$ 2. $\frac{1}{3}$ 3. $\frac{2}{3}$ 4. $\frac{1}{2}$
147. In a certain college 4% of the boys and 1% of the girls are taller than 1.8 meter. Further 60% of the students are girls. If a student is selected at random and is taller than 1.8 meters, then the probability that the student is a girl is
1. $\frac{2}{11}$ 2. $\frac{3}{11}$ 3. $\frac{5}{11}$ 4. $\frac{7}{11}$
148. Ten coins are tossed. The probability of getting at least 8 heads is
1. $\frac{7}{64}$ 2. $\frac{7}{32}$ 3. $\frac{7}{16}$ 4. $\frac{7}{128}$
149. The probability of two events A and B are 0.3 and 0.6 respectively. The probability that both A and B occur simultaneously is 0.18. The probability that neither A nor B occurs is
1. 0.1 2. 0.72 3. 0.42 4. 0.28
150. If m is a number such that $m \leq 5$, then the probability that quadratic equation $2x^2 + 2mx + m + 1 = 0$ has real roots is
1. $\frac{1}{5}$ 2. $\frac{2}{5}$ 3. $\frac{3}{5}$ 4. $\frac{4}{5}$

Padasalai.Net

ANSWERS

7. MATRICES AND DETERMINANTS

1	2	3	4	5	6	7	8	9	10
2	1	2	2	2	2	4	4	2	4
11	12	13	14	15	16	17	18	19	20
2	4	3	2	4	3	3	4	1	3
21	22	23	24	25					
2	3	3	1	2					

8. VECTOR ALGEBRA

26	27	28	29	30	31	32	33	34	35
2	3	4	1	1	4	2	2	2	3
36	37	38	39	40	41	42	43	44	45
4	3	4	3	1	1	1	1	4	2
46	47	48	49	50					
2	2	2	2	4					

9. DIFFERENTIAL CALCULUS LIMITS AND CONTINUITY

51	52	53	54	55	56	57	58	59	60
2	4	3	3	1	4	3	2	1	4
61	62	63	64	65	66	67	68	69	70
3	3	2	2	4	2	3	1	4	2
71	72	73	74	75					
4	1	1	3	2					

10. DIFFERENTIAL CALCULUS DIFFERENTIABILITY AND METHODS OF DIFFERENTIATION

76	77	78	79	80	81	82	83	84	85
2	4	3	3	1	4	3	2	1	4
86	87	88	89	90	91	92	93	94	95
3	3	2	2	4	2	3	1	4	2
96	97	98	99	100					
4	1	1	3	2					

11. INTEGRAL CALCULUS

101	102	103	104	105	106	107	108	109	110
1	3	4	1	3	1	3	2	4	3
111	112	113	114	115	116	117	118	119	120
4	2	4	4	1	2	3	2	4	1
121	122	123	124	125					
3	4	3	1	4					

12. INTRODUCTION TO PROBABILITY THEORY

126	127	128	129	130	131	132	133	134	135
4	3	1	2	2	1	1	4	2	3
136	137	138	139	140	141	142	143	144	145
3	1	4	2	3	1	4	3	2	1
146	147	148	149	150					
2	2	4	4	3					