



Padalsalai's Telegram Groups!

(தலைப்பிற்கு கீழே உள்ள லிங்கை கிளிக் செய்து குழுவில் இணையவும்!)

- Padalsalai's NEWS - Group
https://t.me/joinchat/NIfCqVRBNj9hhV4wu6_NqA
- Padalsalai's Channel - Group
<https://t.me/padasalaichannel>
- Lesson Plan - Group
<https://t.me/joinchat/NIfCqVWwo5iL-21gpzrXLw>
- 12th Standard - Group
https://t.me/Padalsalai_12th
- 11th Standard - Group
https://t.me/Padalsalai_11th
- 10th Standard - Group
https://t.me/Padalsalai_10th
- 9th Standard - Group
https://t.me/Padalsalai_9th
- 6th to 8th Standard - Group
https://t.me/Padalsalai_6to8
- 1st to 5th Standard - Group
https://t.me/Padalsalai_1to5
- TET - Group
https://t.me/Padalsalai_TET
- PGTRB - Group
https://t.me/Padalsalai_PGTRB
- TNPSC - Group
https://t.me/Padalsalai_TNPSC

ILAHY ORIENTAL ARABIC HIGH SCHOOL - CUMBUM9TH MATHS QUARTERLY EXAM (2019-20)MODEL QUESTION PAPERMARKS : 100TIME : 2½ HRI CHOOSE THE CORRECT ANSWER

① Which of the following is correct?

- (1) $\{7\} \in \{1, 2, 3, 4, 5, 6, 7, 8, 9, 10\}$
 (2) $7 \in \{1, 2, 3, 4, 5, 6, 7, 8, 9, 10\}$ (3) $7 \notin \{1, 2, 3, 4, 5, 6, 7, 8, 9, 10\}$
 (4) $\{7\} \notin \{1, 2, 3, 4, 5, 6, 7, 8, 9, 10\}$

② If $B - A$ is B , then $A \cap B$ is

- (1) A (2) B (3) U (4) $\emptyset$

③ If $n(A \cup B \cup C) = 100$, $n(A) = 4x$, $n(B) = 6x$,
 $n(C) = 5x$, $n(A \cap B) = 20$, $n(B \cap C) = 15$, $n(A \cap C) = 25$
 and $n(A \cap B \cap C) = 10$, then the value of x is

- (1) 10 (2) 15 (3) 25 (4) 30


④ The shaded region in the Venn diagram is

(1) $Z - (X \cup Y)$

(2) $(X \cup Y) \cap Z$

(3) $Z - (X \cap Y)$

(4) $Z \cup (X \cap Y)$



⑤ Which one of the following is an irrational number

- (1) $\sqrt{25}$ (2) $\sqrt{\frac{9}{4}}$ (3) $\frac{7}{11}$ (4) π

⑥ If $\frac{1}{7} = 0.\overline{142857}$ then the value of $\frac{5}{7}$ is

- (1) $0.\overline{142857}$ (2) $0.\overline{714285}$ (3) $0.\overline{571428}$ (4) 0.714285

⑦ When written with a rational denominator, the expression $\frac{2\sqrt{3}}{3\sqrt{2}}$ can be simplified as:

- (1) $\frac{\sqrt{2}}{3}$ (2) $\frac{\sqrt{3}}{2}$ (3) $\frac{\sqrt{6}}{3}$ (4) $\frac{2}{3}$

⑧ $(0.000729)^{\frac{-3}{4}} \times (0.09)^{\frac{-3}{4}} =$ _____

- (1) $\frac{10^3}{3^3}$ (2) $\frac{10^5}{3^5}$ (3) $\frac{10^2}{3^2}$ (4) $\frac{10^6}{3^6}$

⑨ If $x^3 + 6x^2 + Kx + 6$ is exactly divisible by $(x+2)$, then $K = ?$

- (1) -6 (2) -7 (3) -8 (4) 11

⑩ The sum of polynomials $p(x) = x^3 - x^2 - 2$,
 $q(x) = x^2 - 3x + 1$

- (1) $x^3 - 3x - 1$ (2) $x^3 + 2x^2 - 1$ (3) $x^3 - 2x^2 - 3x$
 (4) $x^3 - 2x^2 + 3x - 1$

⑪ If $p(x) = 0$ then $(x - a)$ is a _____ of $p(x)$

(1) divisor (2) quotient (3) remainder (4) factor


⑫ $(a + b - c)^2$ is equal to

(1) $(a - b + c)^2$ (2) $(-a - b + c)^2$ (3) $(a + b + c)^2$

(4) $(a - b - c)^2$

⑬ ABCD is a square; diagonals AC and BD meet at O. The number of pairs of congruent triangles with vertex O are

(1) 6 (2) 8 (3) 4 (4) 12



⑭ If bisectors of $\angle A$ and $\angle B$ of a quadrilateral ABCD meet at O, then $\angle AOB$ is

(1) $\angle C + \angle D$ (2) $\frac{1}{2}(\angle C + \angle D)$

(3) $\frac{1}{2}\angle C + \frac{1}{3}\angle D$ (4) $\frac{1}{3}\angle C + \frac{1}{2}\angle D$

PART II

Answers any 10 Questions only

$10 \times 2 = 20$

(Q.no: 28 is compulsory)

⑮ If $A = \{6, 7, 8, 9\}$ and $B = \{8, 10, 12\}$, find $A \Delta B$.

- (16) Let $U = \{0, 1, 2, 3, 4, 5, 6, 7\}$, $A = \{1, 3, 5, 7\}$ and $B = \{0, 2, 3, 5, 7\}$ find the following sets.

(i) $A' \cup B'$ (ii) $(A \cap B)'$

- (17) Test for the Commutative property of union and intersection of the sets.

$P = \{x : x \text{ is a real number between 2 and 7}\}$ and

$Q = \{x : x \text{ is an irrational number between 2 and 7}\}$

- (18) If $n(A) = 25$, $n(B) = 40$, $n(A \cup B) = 50$ and $n(B') = 25$ find $n(A \cap B)$ and $n(U)$.

- (19) Express the following decimal expression into rational numbers.

(i) -5.132 (ii) $17.2\overline{15}$

- (20) Find any two irrational numbers between

$$\frac{6}{7} \text{ and } \frac{12}{3}$$

- (21) Simplify : $\sqrt{63} - \sqrt{175} + \sqrt{28}$


- (22) Find the value of a and b if $\frac{\sqrt{7}-2}{\sqrt{7}+2} = a\sqrt{7}+b$

- (23) The length of a rectangle is $(3x+2)$ units and its breadth is $(3x-2)$ units. Find its area in terms of x . What will be the area if $x = 20$ units.

(24) Find the value of m , if $(x-2)$ is a factor of the polynomial $2x^3 - 6x^2 + mx + 4$

(25) Factorise : $\frac{1}{x^2} + \frac{1}{y^2} + \frac{2}{xy}$

(26) In the figure, AB is parallel to CD , find x



(27) The lengths of the diagonals of a Rhombus are 12 cm and 16 cm. Find the side of the rhombus.

(28) If $\left[y - \frac{1}{y}\right]^3 = 27$, then find the value of $y^3 - \frac{1}{y^3}$

PART - III

Answers any 10 Questions only $10 \times 5 = 50$

(Q.no: 42 is compulsory)

(29) Verify the associative property of intersection of sets for $A = \{-11, \sqrt{2}, \sqrt{5}, 7\}$
 $B = \{\sqrt{3}, \sqrt{5}, 6, 13\}$ and $C = \{\sqrt{2}, \sqrt{3}, \sqrt{5}, 9\}$

(30) Verify $A \cap (B \cup C) = (A \cap B) \cup (A \cap C)$ using Venn diagrams.

(31) If $A = \{-2, 0, 1, 3, 5\}$, $B = \{-1, 0, 2, 5, 6\}$ and $C = \{-1, 2, 5, 6, 7\}$, then show that $A - (B \cup C) = (A - B) \cap (A - C)$

(32) A Survey was conducted among 200 magazine subscribers of three different magazines A, B and C. It was found that 75 members do not subscribe magazine A, 100 members do not subscribe magazine B, 50 members do not subscribe magazine C and 125 subscribe at least two of the three magazines. Find

(i) Number of members who subscribe exactly two magazines

(ii) Number of members who subscribe only one magazine.

(33) (a) Find any two irrational numbers between

(i) $0.3010011000111\dots$ and $0.3020020002\dots$

(ii) $\sqrt{2}$ and $\sqrt{3}$

(b) Find any two rational numbers between

$2.2360679\dots$ and $2.236505500\dots$

③④ (a) Simplify : $2\sqrt[3]{40} + 3\sqrt[3]{625} - 4\sqrt[3]{320}$

(b) Simplify : $\left[\sqrt{\frac{225}{729}} - \sqrt{\frac{25}{144}} \right] \div \sqrt{\frac{16}{81}}$

③⑤ Given $\sqrt{2} = 1.414$, find the value of

$\frac{8-5\sqrt{2}}{3-2\sqrt{2}}$ (to 3 places of decimals)

③⑥ Represent the following numbers in scientific notation:

$\left\{ (0.00003)^6 \times (0.00005)^4 \right\} \div \left\{ (0.009)^3 \times (0.05)^2 \right\}$


③⑦ Without actual division, prove that $f(x) = 2x^4 - 6x^3 + 3x^2 + 3x - 2$ is exactly divisible by $x^2 - 3x + 2$

③⑧ If $(x+a)(x+b)(x+c) = x^3 + 14x^2 + 59x + 70$, Find the value of $\frac{a}{bc} + \frac{b}{ac} + \frac{c}{ab}$

③⑨ Factorise : $x^3 - 5x^2 - 2x + 24$

④⑦ $\triangle ABC$ and $\triangle DEF$ are two triangles in which $AB = DF$, $\angle ACB = 70^\circ$, $\angle ABC = 60^\circ$, $\angle DEF = 70^\circ$ and $\angle EDF = 60^\circ$ Prove that the triangles are congruent.

- ④1 In the given Figure, if
 $AB=2$, $BC=6$, $AE=6$, $BF=8$,
 $CE=7$ and $CF=7$, compute
 the ratio of the area of
 quadrilateral $ABDE$ to the
 area of $\triangle CDF$.



- ④2 The sum of $(x+5)$ observation is
 (x^3+125) . Find the mean of the observations.

PART - IV

Answer any two Questions: $2 \times 8 = 16$

- ④3 (a) Construct the centroid of $\triangle PQR$
 whose sides are $PQ=8\text{ cm}$; $QR=6\text{ cm}$; $RP=7\text{ cm}$

[OR]

- (b) Draw the $\triangle ABC$, where $AB=6\text{ cm}$,
 $\angle B=110^\circ$ and $AC=9\text{ cm}$ and construct
 the centroid.

- ④4 (a) Draw the graph for the following:

$$y = \left[\frac{2}{3} \right] x + 3$$

[OR]

(b) (i) The base of a parallelogram is $(5x+4)$.
Find its height, if the area is $25x^2-16$.

(ii) If $x^2 + \frac{1}{x^2} = 23$, then find the value
of $x + \frac{1}{x}$ and $x^3 + \frac{1}{x^3}$.

Prepared By

M. ABBAS MANTHIRI

B.Sc, B.Ed, M.A, M.Phil

P.T. ASSISTANT

Cell: 894096 5432

Youtube: Ilahi high school.