

Eqn of Parabola is

$$x^2 = -4ay \rightarrow (1)$$

It Passes through (15, -10)

$$(15)^2 = -4a(-10)$$

$$225 = 40a$$

$$\Rightarrow a = \frac{225}{40}$$

$$(1) \Rightarrow x^2 = -4\left(\frac{225}{40}\right)y$$


It passes through (b, -y₁)

$$36 = -4\left(\frac{225}{40}\right)(-y_1)$$

$$\frac{36 \times 10}{225} = y_1$$

$$1.6 = y_1$$

The required height = 10 - y₁ = 10 - 1.6
= 8.4m.



www.TrbTnpSC.com
The required height = 4 - y₁ = 4 - 1 = 3m


(4)

(a) Eqn of Parabola is

$$y^2 = 4ax$$

$$a = 1.2$$

$$y^2 = 4.8x$$




(b) It passes through (x₁, 2.5)

$$(2.5)^2 = 4.8x_1$$

$$x_1 = \frac{2.5 \times 2.5}{4.8} = 1.3m$$

The depth of the satellite dish at the vertex is 1.3m.

(5)



Eqn of the parabola is

$$(x-h)^2 = 4a(y-k)$$

$$(x-0)^2 = 4a(y-3)$$

It passes through (30, 16)

$$(30)^2 = 4a(16-3)$$

$$a = \frac{30 \times 30}{13 \times 4}$$

$$x^2 = 4\left(\frac{30 \times 30}{13 \times 4}\right)(y-3)$$

It passes through (6, y₁)

$$6^2 = \frac{900}{13}(y_1-3)$$

$$(y_1-3) = \frac{36 \times 13}{900} = 0.52$$

$$y_1 = 3 + 0.52 = 3.52m$$

It also passes through (12, y₂)

$$12^2 = \frac{900}{13}(y_2-3)$$

$$y_2-3 = \frac{144 \times 13}{900} = 2.08$$

$$y_2 = 3 + 2.08 = 5.08m$$

(3)

CA = a = 8, CB = b = 6

Eqn of ellipse is

$$\frac{x^2}{b^2} + \frac{y^2}{a^2} = 1$$

$$\frac{x^2}{25} + \frac{y^2}{64} = 1$$

It passes through (4, y₁)


$$\frac{16}{25} + \frac{y_1^2}{64} = 1$$

$$\frac{y_1^2}{64} = 1 - \frac{16}{25}$$

$$y_1^2 = 64\left(\frac{9}{25}\right)$$

$$y_1 = \frac{8 \times 3}{5} = \frac{24}{5} = 4.8m$$

$\therefore$ The required width for the opening is 2y = 2(4.8) = 9.6m.



(3)

Eqn of the Parabola is

$$x^2 = -4ay \rightarrow (1)$$

It passes through (0.5, -4)

$$(0.5)^2 = -4a(-4)$$


$$a = \frac{0.25}{16}$$

$$(1) \Rightarrow x^2 = -4\left[\frac{0.25}{16}\right]y$$

It passes through (0.25, -y₁)

$$(0.25)^2 = -4\left[\frac{0.25}{16}\right](-y_1)$$

$$y_1 = \frac{4 \times 0.25}{0.25} = 4 \times 0.25 = 1m$$



$$3k = 150$$

$$k = 50$$

$(x_1, 50)$ lies on the parabola

$$\frac{x_1^2}{30^2} - \frac{50^2}{44^2} = 1$$

$$\frac{x_1^2}{30^2} = 1 + \frac{50^2}{44^2}$$

$$x_1^2 = \frac{30^2}{44^2} [1936 + 2500]$$

$$x_1 = \frac{30}{44} \sqrt{4436} = \frac{30}{44} (66.60)$$

$$x_1 = 45.40$$

Radius of the top of the tower is 45.40m

$(x_2, -100)$ lies on the hyperbola

$$\frac{x_2^2}{30^2} - \frac{100^2}{44^2} = 1$$

$$\frac{x_2^2}{30^2} = 1 + \frac{100^2}{44^2}$$

$$x_2^2 = \frac{30^2}{44^2} (1936 + 10000)$$

$$x_2 = \frac{30}{44} \sqrt{11936}$$

$$= \frac{30}{44} \times 109.25$$

$$x_2 = \frac{3 \times 277.5}{44} = 74.48 \text{ m}$$

Radius of the base of the tower is 74.48m.

$$\textcircled{7} \quad \triangle BAP \cong \triangle PCD$$

$$\text{Let } \angle ABP = \angle CPD = \theta$$

In $\triangle PAB$

$$\sin \theta = \frac{y}{0.3}$$

In $\triangle PCD$


$$\cos \theta = \frac{x}{0.9}$$

$$\cos^2 \theta + \sin^2 \theta = 1$$

$$\frac{x^2}{(0.9)^2} + \frac{y^2}{(0.3)^2} = 1$$

$$e = \sqrt{1 - \frac{b^2}{a^2}} = \sqrt{1 - \frac{0.3^2}{0.9^2}} = \sqrt{1 - \left(\frac{1}{3}\right)^2}$$

$$e = \sqrt{\frac{8}{9}} = \frac{2\sqrt{2}}{3} \text{ m.}$$



⑧

The eqn of the parabola is

$$x^2 = -4ay$$

It passes through $(3, -2.5)$

$$3^2 = -4a(-2.5)$$

$$a = \frac{9}{10}$$

$$x^2 = -4\left(\frac{9}{10}\right)y$$

$$x^2 = -\frac{18}{5}y$$


It passes through $(x_1, -7.5)$

$$x_1^2 = -\frac{18}{5}(-7.5)$$

$$x_1^2 = 27$$

$$x_1 = 3\sqrt{3} \text{ m}$$

The water strikes the ground $3\sqrt{3} \text{ m}$ beyond the vertical line.



⑨

The eqn of parabola is $x^2 = -4ay$

It passes through $(b, -4)$

$$b^2 = 16a$$

$$\Rightarrow a = \frac{b^2}{16} = \frac{9}{4}$$

$$x^2 = -4\left(\frac{9}{4}\right)y$$

$$x^2 = -9y$$

Diff. w.r. to x


$$2x = -9 \frac{dy}{dx}$$

$$\frac{dy}{dx} = -\frac{2x}{9}$$

At $(-b, -4)$

$$\tan \theta = \frac{dy}{dx} = -\frac{2}{9}(-b) = \frac{4}{3}$$

$$\theta = \tan^{-1} \frac{4}{3}$$



⑩

$$ae = 5$$

$$AP \cdot BP = 2a = 6$$

$$a = 3$$


$$b^2 = a^2(e^2 - 1) = (3^2)(\frac{25}{9} - 1) = 25 - 9$$

$$b^2 = 16$$

The eqn of the required hyperbola

$$\frac{x^2}{9} - \frac{y^2}{16} = 1$$

The locus of P is hyperbola.



Eg: 5.37

Sol:


Let $a = 17$, $b = 8$

$$c^2 = a^2 - b^2$$

$$= 289 - 64$$

$$c^2 = 225$$

$$c = 15$$

Foci are $F_1(15, 0)$, $F_2(-15, 0)$ 

Eg: 5.38

Eqn of ellipse is $\frac{(x-11)^2}{484} + \frac{y^2}{64} = 1$

$$a^2 = 484, b^2 = 64$$

$$c^2 = a^2 - b^2$$


$$= 484 - 64$$

$$c^2 = 420$$

$$c \approx 20.5$$

$\therefore$ The patient's kidney stone should be placed 20.5 cm from the centre of the ellipse.

Eg: 5.39.

Centre: $C(0, 300)$

Eqn of hyperbola is

$$\frac{(y-300)^2}{a^2} - \frac{(x-0)^2}{b^2} = 1 \rightarrow (1)$$

It passes through $(0, 400)$

$$\frac{(400-300)^2}{a^2} - 0 = 1$$

$$a^2 = (100)^2$$

$$a = 100$$

In $\triangle ABP$

$$AP^2 = AB^2 + BP^2$$

$$(x_1 + 200)^2 = 600^2 + x_1^2$$

$$x_1^2 + 400x_1 + (200)^2 = (600)^2 + x_1^2$$

$$400x_1 = (600)^2 - (200)^2$$

$$= 360000 - 40000$$

$$x_1 = \frac{320000}{400}$$

$$x_1 = 800$$

$P(800, 600)$ lies on the hyperbola

$$(1) \Rightarrow \frac{(y-300)^2}{a^2} - \frac{x^2}{b^2} = 1$$

$$\frac{(600-300)^2}{a^2} - \frac{(800)^2}{b^2} = 1$$

$$\frac{(300)^2}{(100)^2} - 1 = \frac{(800)^2}{b^2}$$

$$\frac{90000}{10000} - 1 = \frac{(800)^2}{b^2}$$

$$b^2 = \frac{800000}{8}$$

$$b^2 = 80000$$

The required eqn of hyperbola is

$$\frac{(y-300)^2}{10000} - \frac{x^2}{80000} = 1$$

The exact location can be determined using data from a 3rd station.

Eg: 5.40

$$F_1F_2 = 14 - 2 = 12$$

$$2ae = 12$$

$$ae = 6$$

$$V_2F_1 = CF_2 - CV_2$$

$$1 = ae - a$$

$$1 = 6 - a$$

$$a = 5$$

$$b^2 = a^2(e^2 - 1)$$


$$= a^2e^2 - a^2$$

$$= 36 - 25$$

$$b^2 = 11$$

The eqn of the hyperbola is

$$\frac{y^2}{25} - \frac{x^2}{11} = 1$$



Eg: 5.30

Let $a = 6$, $b = 3$ Eqn of the ellipse $(-6, 0)$

$$\text{is } \frac{x^2}{36} + \frac{y^2}{9} = 1 \quad \text{--- (1)}$$

When $x = 1.5$, $y = ?$ or $x = \frac{3}{2}$

$$\text{(1)} \Rightarrow \left(\frac{3}{2}\right)^2 + \frac{y^2}{9} = 1$$


$$y^2 = 9 \left[1 - \frac{9}{144} \right]$$

$$= 9 \times \frac{135}{144} = \frac{135}{16}$$

$$y = \frac{\sqrt{135}}{16} = \frac{11.62}{16}$$

$$y = 2.90$$

The truck will clear the archway.



Eg: 5.33

Ssn of Parabola is

$$y^2 = 4ax$$

$$a = 2$$


$$y^2 = 8x$$

It passes through $P(3, y_1)$

$$y_1^2 = 8 \times 3$$

$$y_1 = 2\sqrt{2} \times \sqrt{3}$$

$$y_1 = 2\sqrt{6}$$

Required width $2y_1 = 4\sqrt{6} \text{ m}$.

Eg: 5.34

Sol:


Ssn of the parabola

$$y = \frac{1}{32} x^2$$

$$x^2 = 32y$$

$$x^2 = 4(8)y$$

$$\Rightarrow a = 8$$



The heating tube needs to be placed 8 units above the vertex of the parabola.

Eg: 5.31

Sol:

$$\text{Let } SA = 94.5 \times 10^6 \text{ km}$$


$$SA' = 152 \times 10^6 \text{ km}$$

$$SA = a - c = 94.5 \times 10^6 \rightarrow \text{(1)}$$

$$SA' = a + c = 152 \times 10^6 \rightarrow \text{(2)}$$

$$\text{(2)} - \text{(1)} \Rightarrow 2c = 57.5 \times 10^6 = 575 \times 10^5 \text{ km}$$

$$SS' = 575 \times 10^5 \text{ km}$$



Eg: 5.32

The eqn of parabola is

$$x^2 = -4ay \rightarrow \text{(1)}$$

It passes through

$$(20, -15)$$


$$(20)^2 = -4a(-15)$$

$$\Rightarrow 4a = \frac{400}{15} = \frac{80}{3}$$

$$\text{(1)} \Rightarrow x^2 = -\frac{80}{3}y$$

The required eqn is

$$3x^2 = -80y$$



Eg: 5.35

Sol:

Ssn of Parabola is

$$y^2 = 4ax \rightarrow \text{(1)}$$

(1) It passes through $(30, 20)$


$$(20)^2 = 4a(30)$$

$$\frac{400}{30} = 4a$$

$$\text{(1)} \Rightarrow y^2 = \frac{40}{3}x$$

$$4a = \frac{40}{3}$$

$$a = \frac{10}{3}$$

The bulb is at focus $(\frac{10}{3}, 0)$ 

Eg: 5.36


Sol:

Ssn of ellipse $\frac{x^2}{16} + \frac{y^2}{9} = 1$

$$a^2 = 16, b^2 = 9$$

$$c^2 = a^2 - b^2 = 7$$

$$c = \pm\sqrt{7}$$

FOCI: $F_1(\sqrt{7}, 0), F_2(-\sqrt{7}, 0)$. Focus of the parabola $(\sqrt{7}, 0) \Rightarrow a = \sqrt{7}$ Ssn of the parabola is $y^2 = 4\sqrt{7}x$ 

$$\frac{2x}{3} - \frac{y}{2} = 1.$$

$$4x - 3y - 6 = 0$$

Normal equation at 'O'

$$\frac{ax}{\sec \theta} + \frac{by}{\tan \theta} = a^2 + b^2$$

$$\theta = \pi/3$$

$$\frac{3x}{2} + \frac{2\sqrt{3}y}{\sqrt{3}} = 9 + 12.$$

$$\frac{3x}{2} + 2y = 21.$$

$$3x + 4y - 42 = 0$$

7. Solu:

Tangent equation at 't₁' is

$$yt_1 = x + at_1^2 \rightarrow (1)$$

at 't₂' is

$$yt_2 = x + at_2^2 \rightarrow (2)$$

$$(1) - (2) \Rightarrow y(t_1 - t_2) = a(t_1^2 - t_2^2)$$

$$y = \frac{a(t_1 + t_2)(t_1 - t_2)}{t_1 - t_2}$$

$$y = a(t_1 + t_2)$$

Using $y = a(t_1 + t_2)$ in (1), we get

$$a(t_1 + t_2)t_1 = x + at_1^2$$

$$at_1^2 + at_1t_2 = x + at_1^2$$

$$x = at_1t_2$$

$\therefore$ The point of intersection is

$$[at_1t_2, a(t_1 + t_2)]$$

8. Solu:

Normal equation of a parabola at $P(at_1^2, 2at_1)$ is

$$y + xt_1 = at_1^3 + 2at_1$$

It meets the parabola again at $Q(at_2^2, 2at_2)$

$$2at_2 + (at_1^2)t_1 = at_1^3 + 2at_1$$

$$2at_2 + at_1t_2^2 = at_1^3 + 2at_1$$

$$2at_2 - 2at_1 = at_1^3 - at_1t_2^2$$

$$2a(t_2 - t_1) = at_1(t_1^2 - t_2^2)$$

$$2a(t_2 - t_1) = at_1(t_1 - t_2)(t_1 + t_2)$$

$$2a(t_2 - t_1) = -at_1(t_2 - t_1)(t_1 + t_2)$$

$$-2 = t_1(t_1 + t_2)$$

$$t_1 + t_2 = -\frac{2}{t_1}$$

$$t_2 = -\frac{2}{t_1} - t_1$$

$$t_2 = -(t_1 + \frac{2}{t_1})$$

Hence proved.

2. Solu:-

$$\frac{x^2}{16} - \frac{y^2}{64} = 1$$

$$a^2 = 16, b^2 = 64.$$

$$\text{slope of the tangent } m = -\frac{10}{3}$$

$$m = \frac{10}{3}.$$

Any tangent to the hyperbola

$$\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1 \text{ is of the form,}$$

$$y = mx \pm \sqrt{a^2 m^2 - b^2}$$

$$y = \frac{10}{3}x \pm \sqrt{16\left(\frac{100}{9}\right) - 64}$$

$$y = \frac{10}{3}x \pm \sqrt{\frac{1024}{9}}$$

$$y = \frac{10}{3}x \pm \frac{32}{3}$$

$$3y = 10x \pm 32.$$

$$10x - 3y \pm 32 = 0.$$

Tangent equations are

$$10x - 3y + 32 = 0$$

$$10x - 3y - 32 = 0.$$

3. Solu:-

$$x - y + 4 = 0$$

$$y = x + 4.$$

$$m = 1, c = 4$$

$$x^2 + 3y^2 = 12$$

$$\frac{x^2}{12} + \frac{y^2}{4} = 1$$

$$a^2 = 12, b^2 = 4$$

Condition for tangency

$$c^2 = a^2 m^2 + b^2$$

$$c^2 = 16, a^2 m^2 + b^2 = 12(1)^2 + 4 = 16.$$

$\therefore$ The line is a tangent to the ellipse.

point of contact:

$$\left(-\frac{a^2 m}{c}, \frac{b^2}{c}\right) = \left[-\frac{12(1)}{4}, \frac{4}{4}\right]$$

$$= (-3, 1)$$

4. Solu:

$$y^2 = 16x$$

$$4a = 16 \Rightarrow a = 4$$

Any line $\perp$ r to $2x + 2y + 3 = 0$

$$\text{is } 2x - 2y + k = 0$$

$$x - y + \frac{k}{2} = 0 \rightarrow \textcircled{1}$$

$$y = x + \frac{k}{2}.$$

$$m = 1, c = \frac{k}{2}.$$

The condition for the line $y = mx + c$ be a tangent to the parabola $y^2 = 4ax$ is $c = \frac{a}{m}$

$$c = \frac{4}{1}$$

$$\frac{k}{2} = 4 \Rightarrow \boxed{k = 8}$$

Substitute $k = 8$ in $\textcircled{1}$ Required $\perp$ r line is

$$\boxed{x - y + 4 = 0}$$

5. Solu:

$$y^2 = 8x$$

$$4a = 8 \Rightarrow a = 2$$

$$\text{Here } t = 2, a = 2.$$

The equation of tangent at 't' on the parabola is

$$yt = x + at^2$$

$$y(2) = x + 2(4)$$

$$2y = x + 8.$$

$$\boxed{x - 2y + 8 = 0}$$

$$6. \text{ Solu: } 12x^2 - 9y^2 = 108$$

$$\div \text{ by } 108 \Rightarrow \frac{x^2}{9} - \frac{y^2}{12} = 1$$

$$a^2 = 9, b^2 = 12$$

$$a = 3, b = 2\sqrt{3}.$$

Tangent equation at 'O'

$$\frac{x \sec \theta}{a} - \frac{y \tan \theta}{b} = 1.$$

$$\theta = \pi/2, \frac{x \sec(\pi/2)}{3} - \frac{y \tan \pi/2}{2\sqrt{3}} = 1$$

$$\frac{2x}{3} - \frac{\sqrt{3}y}{2\sqrt{3}} = 1$$

Note: Tangent equation at (x_1, y_1) for all the curves,
 Replace x^2 by xx_1 , y^2 by yy_1 ,
 x by $\frac{1}{2}(x+x_1)$, y by $\frac{1}{2}(y+y_1)$
 and xy by $\frac{1}{2}(xy_1+yx_1)$.

Example: 5.27.

Solu:-

$$x^2 + 6x + 4y + 5 = 0.$$

Tangent equation at (x_1, y_1) is

$$xx_1 + 6\left(\frac{x+x_1}{2}\right) + 4\left(\frac{y+y_1}{2}\right) + 5 = 0.$$

$$xx_1 + 3(x+x_1) + 2(y+y_1) + 5 = 0.$$

$$(x_1, y_1) = (1, -3).$$

$$x(1) + 3(x+1) + 2(y-3) + 5 = 0$$

$$x + 3x + 3 + 2y - 6 + 5 = 0$$

$$4x + 2y + 2 = 0$$

$$\div \text{ by } 2, \quad 2x + y + 1 = 0.$$

Normal Equation: at $(1, -3)$

$$x - 2y + k = 0$$

$$1 - 2(-3) + k = 0$$

$$1 + 6 + k = 0$$

$$7 + k = 0$$

$$k = -7.$$

$$\therefore x - 2y - 7 = 0$$

Example: 5.28.

Solu:-

$$x^2 + 4y^2 = 32$$

$$\frac{x^2}{32} + \frac{y^2}{8} = 1.$$

$$a^2 = 32, \quad b^2 = 8$$

$$a = 4\sqrt{2}, \quad b = 2\sqrt{2}$$

Tangent equation at 'O'

$$\frac{x \cos \theta}{a} + \frac{y \sin \theta}{b} = 1$$

$$\theta = \pi/4,$$

$$\frac{x \cos \pi/4}{4\sqrt{2}} + \frac{y \sin \pi/4}{2\sqrt{2}} = 1$$

$$\frac{x}{8} + \frac{y}{4} = 1 \Rightarrow x + 2y - 8 = 0$$

Normal Equation at 'O'

$$\frac{ax}{\cos \theta} - \frac{by}{\sin \theta} = a^2 - b^2$$

$$\theta = \pi/4,$$

$$\frac{4\sqrt{2}x}{\cos \pi/4} - \frac{2\sqrt{2}y}{\sin \pi/4} = 32 - 8$$

$$8x - 4y = 24.$$

$$\therefore 2x - y - 6 = 0$$

Exercise: 5.4.

1. Solu:

$$2x^2 + 7y^2 = 14, \quad (5, 2)$$

$$\div \text{ by } 14, \quad \frac{x^2}{7} + \frac{y^2}{2} = 1.$$

$$a^2 = 7, \quad b^2 = 2.$$

Any tangent to the Ellipse

$\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$ is of the form,

$$y = mx \pm \sqrt{a^2 m^2 + b^2}$$

$$2 = m(5) \pm \sqrt{7m^2 + 2}.$$

$$2 - 5m = \pm \sqrt{7m^2 + 2}$$

Squaring on both sides, we get

$$4 - 20m + 25m^2 = 7m^2 + 2.$$

$$18m^2 - 20m + 2 = 0$$

$$\div \text{ by } 2, \quad 9m^2 - 10m + 1 = 0$$

$$(9m-1)(m-1) = 0.$$

$$\therefore m = \frac{1}{9} \text{ or } m = 1.$$

Tangent Equations:

when $m=1$, $(x_1, y_1) = (5, 2)$

$$y - y_1 = m(x - x_1)$$

$$y - 2 = 1(x - 5)$$

$$x - y - 3 = 0$$

when $m = \frac{1}{9}$, $(x_1, y_1) = (5, 2)$

$$y - 2 = \frac{1}{9}(x - 5)$$

$$9y - 18 = x - 5$$

$$x - 9y + 13 = 0$$

Exercise 5.3.

1. $2x^2 - y^2 = 7.$

Solu:

$2x^2 - y^2 = 7$

Compared with

$Ax^2 + Bxy + Cy^2 + Dx + Ey + F = 0$

$A = 2, B = 0, C = -1$

$B^2 - 4AC = 0 - 4(2)(-1) = 8 > 0$

 $\therefore$ The given equation is a Hyperbola.

2. Solu:

$3x^2 + 3y^2 - 4x + 2y + 10 = 0$

$A = 3, B = 0, C = 3,$

$A = C = 3 \neq B = 0.$

The given equation is a circle.

3. Solu:

$3x^2 + 3y^2 = 14.$

$A = 3, B = 0, C = 3,$

$B^2 - 4AC = 0 - 4(3)(3) = -12 < 0$

 $\therefore$ The given equation is an Ellipse.

4. Solu:

$x^2 + y^2 + x - y = 0$

$A = 1, B = 0, C = 1, D = 1, E = -1,$

$F = 0$

Here $A = C = 1, B = 0.$

 $\therefore$ The given equation is a circle.

5. Solu:

$11x^2 - 25y^2 - 44x + 50y - 256 = 0$

$A = 11, B = 0, C = -25, D = -44,$

$E = 50, F = -256.$

$B^2 - 4AC = 0 - 4(11)(-25) = 1100 > 0$

The given equation is a Hyperbola.

b. Solu:

$y^2 + 4x + 2y + 4 = 0.$

$A = 0, B = 0, C = 1, D = 4, E = 2,$

$F = 4.$

$B^2 - 4AC = 0 - 4(0)(1) = 0.$

The given equation is a Parabola.

Example: 5.2.6.

i. Solu:

$16y^2 = -4x^2 + 64.$

$4x^2 + 16y^2 - 64 = 0.$

$A = 4, B = 0, C = 16,$

$B^2 - 4AC = 0 - 4(4)(16) = -256 < 0$

 $\therefore$ The given equation is an Ellipse.

ii. Solu:

$x^2 + y^2 = -4x - y + 4$

$x^2 + y^2 + 4x + y - 4 = 0$

$A = 1, B = 0, C = 1, D = 4, E = 1, F = -4$

$A = C = 1, B = 0$

 $\therefore$ The given equation is a circle.

iii) $x^2 - 2y = x + 3$

$x^2 - x - 2y - 3 = 0$

$A = 1, B = 0, C = 0, D = -1, E = -2,$

$F = -3$

$B^2 - 4AC = 0 - 4(1)(0) = 0.$

 $\therefore$ The given equation is a Parabola.

iv. Solu:

$4x^2 - 9y^2 - 16x + 18y - 29 = 0$

$A = 4, B = 0, C = -9, D = -16, E = 18,$

$F = -29$

$B^2 - 4AC = 0 - 4(4)(-9) = 144 > 0$

 $\therefore$ The given equation is a Hyperbola.

Centre $(h, k) = (0, 0)$
 Focus $(h, k \pm ae) = (0, \pm 5)$
 Vertices $(h, k \pm a) = (0, \pm 4)$.

$$e = \sqrt{\frac{a^2 + b^2}{a^2}} = \frac{5}{4}$$

$$\text{Directrix } y = \pm \frac{a}{e} = \frac{(4)}{(5/4)}$$

$$y = \pm \frac{16}{5}$$

EXE 5.2 (S)

$$(i) \frac{(x-3)^2}{225} + \frac{(y-4)^2}{289} = 1.$$

$$b^2 = 225 \quad a^2 = 289$$

$$(ae)^2 = a^2 - b^2 = 289 - 225$$

$$(ae)^2 = 64$$

$$\boxed{ae = 8}$$

Centre $(h, k) = (3, 4)$

Focus $(h, k \pm ae) = (3, 4 \pm 8)$

$F_1(3, 12) \quad F_2(3, -4)$

Vertices $(h, k \pm a) = (3, 4 \pm 17)$

$A(3, 21), A'(3, -13)$.

$$e = \sqrt{\frac{a^2 - b^2}{a^2}} = \frac{8}{17}$$

$$\text{Directrix } y - 4 = \pm \frac{a}{e}$$

$$y - 4 = \pm \frac{17}{(8/17)}$$

$$y - 4 = \pm \frac{289}{8}$$

$$y = 4 \pm \frac{289}{8}$$

$$(ii) \frac{(x+1)^2}{100} + \frac{(y-2)^2}{64} = 1.$$

$$a^2 = 100, \quad b^2 = 64.$$

$$(ae)^2 = a^2 - b^2 = 100 - 64 = 36$$

$$\boxed{ae = 6}$$

Centre $(h, k) = (-1, 2)$.

Focus $(h \pm ae, k) = (-1 \pm 6, 2)$

$F_1(5, 2) \text{ \& } F_2(-7, 2)$.

Vertices $(h \pm a, k) = (-1 \pm 10, 2)$

$A(9, 2), A'(-11, 2)$.

$$e = \sqrt{\frac{a^2 - b^2}{a^2}} = \frac{6}{10} = \frac{3}{5}$$

Equation of Directrices $x = \pm \frac{a}{e}$

$$x + 1 = \pm \frac{a}{e}$$

$$x + 1 = \pm \frac{10}{(3/5)}$$

$$x + 1 = \pm \frac{50}{3}$$

$$x = -1 \pm \frac{50}{3}.$$

$$(iii) \frac{(x+3)^2}{225} - \frac{(y-4)^2}{64} = 1.$$

$$a^2 = 225, \quad b^2 = 64$$

$$(ae)^2 = a^2 + b^2 = 225 + 64 = 289$$

$$\boxed{ae = 17}$$

Centre $(h, k) = (-3, 4)$

Focus $(h \pm ae) = (-3 \pm 17, 4)$

$F_1(14, 4), F_2(-20, 4)$

Vertices $(h \pm a, k) = (-3 \pm 15, 4)$

$A(12, 4), A'(-18, 4)$

$$\text{Directrix } x = \pm \frac{a}{e} = \frac{15}{17/15} = \frac{225}{17}$$

$$x + 3 = \pm \frac{225}{17}$$

$$x = -3 \pm \frac{225}{17}$$

$$(i) y^2 - 4y - 8x + 12 = 0$$

$$y^2 - 4y + 4 = +8x - 8$$

$$(y-2)^2 = +8(x-1)$$

$$y^2 = +4ax \therefore a=2$$

The parabola open ^{Right} leftward

$$\text{Vertices } (h, k) = (1, 2)$$

$$\text{Focus } (h+a, 0+k) = (3, 2)$$

$$\text{Latus rectum } x = +a$$

$$x-1 = +2$$

$$x = +1+2$$

$$\boxed{x=3}$$

$$\text{Directrix } x = -a$$

$$x-1 = -2$$

$$\boxed{x=-1}$$

Length of the latus rectum.

$$4a = 4(2) = 8.$$

Ex 5.2 (5).

$$(i) \frac{x^2}{25} + \frac{y^2}{9} = 1$$

$$a^2 = 25 \Rightarrow a = \pm 5$$

$$b^2 = 9 \Rightarrow b = \pm 3.$$

$$(ae)^2 = a^2 - b^2 = 25 - 9 = 16$$

$$\boxed{ae=4}$$

$$\text{Centre } (h, k) = (0, 0)$$

$$\text{Focus } (h \pm ae, k) = (\pm 4, 0).$$

$$\text{Vertices } (h \pm a, k) = (\pm 5, 0).$$

$$\text{Directrix } x = \pm \frac{a}{e}$$

$$e = \sqrt{\frac{a^2 - b^2}{a^2}} = \sqrt{\frac{25 - 9}{25}} = \sqrt{\frac{16}{25}} = \frac{4}{5}$$

$$x = \pm \frac{5}{(4/5)} = \pm \frac{25}{4}.$$

$$(ii) \frac{x^2}{3} + \frac{y^2}{10} = 1.$$

$$a^2 = 10, b^2 = 3.$$

$$(ae)^2 = a^2 - b^2 = 10 - 3 = 7$$

$$ae = \sqrt{7}$$

$$\text{Centre } (h, k) = (0, 0).$$

$$\text{Vertices } (h, \pm a) = (0, \pm \sqrt{10}).$$

$$\text{Focus } (h, \pm ae) = (0, \pm \sqrt{7})$$

$$e = \sqrt{\frac{a^2 - b^2}{a^2}} = \sqrt{\frac{10 - 3}{10}} = \frac{\sqrt{7}}{\sqrt{10}}$$

$$\text{Directrix } = \pm \frac{a}{e} = \pm \frac{\sqrt{10}}{(\sqrt{7}/\sqrt{10})}$$

$$y = \pm \frac{10}{\sqrt{7}}$$

$$(iii) \frac{x^2}{25} - \frac{y^2}{144} = 1.$$

$$a^2 = 25, b^2 = 144.$$

$$(ae)^2 = a^2 + b^2 = 25 + 144 = 169$$

$$ae = 13.$$

$$\text{Centre } (h, k) = (0, 0)$$

$$\text{Focus } (h \pm ae, k) = (\pm 13, 0).$$

$$\text{Vertices } (h \pm a, k) = (\pm 5, 0).$$

$$e = \sqrt{\frac{a^2 + b^2}{a^2}} = \frac{13}{5}$$

$$\text{Directrix } x = \pm \frac{a}{e} = \pm \frac{5}{(13/5)}$$

$$x = \pm \frac{25}{13}.$$

$$(iv) \frac{y^2}{16} - \frac{x^2}{9} = 1$$

$$a^2 = 16, b^2 = 9$$

$$(ae)^2 = a^2 + b^2 = 16 + 9 = 25$$

$$\boxed{ae=5}$$

Equation of the hyperbola

$$\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$$

At (5, -2) $\Rightarrow a = 4$

$$\frac{25}{16} - \frac{4}{b^2} = 1$$

$$\frac{25}{16} - 1 = \frac{4}{b^2}$$

$$\frac{25-16}{16} = \frac{4}{b^2}$$

$$\frac{9}{16} = \frac{4}{b^2}$$

$$\frac{b^2}{4} = \frac{16}{9}$$

$$b^2 = \frac{64}{9}$$

Equation of the hyperbola.

$$\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$$

$$\frac{x^2}{16} - \frac{y^2}{64/9} = 1$$

$$\frac{x^2}{16} - \frac{9y^2}{64} = 1$$

Ex 5.2 (4).

$$(i) y^2 = 16x \Rightarrow y^2 = 4ax$$

$$y^2 = 4(4)x \quad \boxed{a=4}$$

The parabola open rightward

vertices (0,0)

Focus (a,0) = (4,0)

Latus rectum $x = a \Rightarrow x = 4$

Directrix $x = -a \Rightarrow x = -4$

length of L.R $4a = 4(4) = 16$

$$(ii) x^2 = 24y$$

$$x^2 = 4ay$$

$$x^2 = 4(6)y \quad a = 6$$

The parabola open upward

vertices (0,0) = (h,k)

focus (0,a) = (0,6)

Latus rectum $y = a \Rightarrow y = 6$

Directrix $y = -a \Rightarrow y = -6$

length of L.R $4a = 4(6) = 24$

$$(iii) y^2 = -8x \quad y^2 = -4ax$$

$$y^2 = -4(2)x \quad a = 2$$

The parabola open leftward

vertices (0,0) = (h,k)

focus (-a,0) = (-2,0)

Latus rectum $x = -a \Rightarrow x = -2$

Directrix $x = a \Rightarrow x = 2$

length of the Latus rectum $4a = 4(2) = 8$

$$iv) x^2 - 2x + 8y + 17 = 0$$

$$x^2 - 2x + 17 = -8y$$

$$x^2 - 2x + 1 = -8y - 16$$

$$(x-1)^2 = -8(y+2)$$

The parabola open downward

Centre (h,k) = (1, -2)

Vertices (0th, -a+k) = (1, -4)

Latus rectum $y = -a$
 $y+2 = -2$
 $\boxed{y = -4}$

Directrix $y = a$
 $y+2 = +2$
 $\boxed{y = 0}$

length of the latus rectum $4a = 4(2) = 8$

EXAMPLE: 5.25.

Astronomical unit long $2a = 36.18$
 wide (or) short $2b = 9.12$

$$b^2 = (ae)^2 - a^2$$

$$a^2 + b^2 = a^2 e^2$$

$$e^2 = \frac{a^2 + b^2}{a^2}$$

$$e = \frac{\sqrt{a^2 + b^2}}{a}$$

$$= \frac{\sqrt{(18.09)^2 + (4.56)^2}}{18.09}$$

$$e \approx 0.97.$$

EXE 5.2 (3) (i)

$$\text{Foci } (\pm ae, 0) = (\pm 2, 0)$$

$$ae = 2$$

$$e = 3/2.$$

$$a(3/2) = 2$$

$$a = \frac{4}{3}$$

$$\therefore b^2 = (ae)^2 - a^2$$

$$= 4 - \frac{16}{9}$$

$$= \frac{36 - 16}{9}$$

$$b^2 = \frac{20}{9}$$

Equation of the hyperbola

$$\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$$

$$\frac{x^2}{16/9} - \frac{y^2}{20/9} = 1$$

$$\frac{9x^2}{16} - \frac{9y^2}{20} = 1$$

EXE 5.2 (3) (ii)

Centre (h, k) (2, 1), Foci (8, 1)

The distance between centre & focus

$$ae = \sqrt{(8-2)^2 + (1-1)^2}$$

$$= \sqrt{6^2}$$

$$ae = 6.$$

Corresponding directrix $x = 4$

(4, 1)

The distance between centre & directrix

$$\frac{a}{e} = \sqrt{(4-2)^2 + (1-1)^2}$$

$$= \sqrt{2^2}$$

$$\frac{a}{e} = 2$$

$$\rightarrow (ae) \left(\frac{a}{e} \right) = 6 \times 2$$

$$a^2 = 12$$

$$(ae)^2 = 36.$$

$$b^2 = (ae)^2 - a^2$$

$$= 36 - 12$$

$$b^2 = 24.$$

Equation of the hyperbola.

$$\frac{(x-2)^2}{a^2} - \frac{(y-1)^2}{b^2} = 1$$

$$\frac{(x-2)^2}{12} - \frac{(y-1)^2}{24} = 1.$$

EXE 5.2 (3) (iii).

passing through (5, 2) = (x, y).

length of transverse axis, long x-axis
length 8 units.

$$2a = 8$$

$$a = 4$$

Centre (h, k) = (0, 0)

EXAMPLE 5.20

$$4x^2 + 36y^2 + 40x - 288y + 532 = 0$$

$$4[x^2 + 10x] + 36[y^2 - 8y] = -532$$

$$4[(x+5)^2 - 25] + 36[(y-4)^2 - 16] = -532$$

$$4(x+5)^2 + 36(y-4)^2 = -532 + 100 + 576$$

$$4(x+5)^2 + 36(y-4)^2 = 144$$

$$\frac{(x+5)^2}{36} + \frac{(y-4)^2}{4} = 1$$

Centre $(-5, 4) = (h, k)$

$$a^2 = 36, \quad b^2 = 4$$

$$a = 6, \quad b = 2$$


Length of the major axis $2a = 12$

Length of the minor axis $2b = 4$

EXAMPLE 5.22.

vertices $(0, \pm a) = (0, \pm 4)$

Focus $(0, \pm ae) = (0, \pm 6)$



The midpoint of the foci $(0, 0)$

$$2a = 8 \quad 2ae = 12$$

$$a = 4 \quad ae = 6$$

$$b^2 = (ae)^2 - a^2$$

$$= 36 - (4)^2 = 36 - 16$$

$$b^2 = 20$$

Equation of the hyperbola

$$\frac{(y-k)^2}{a^2} - \frac{(x-h)^2}{b^2} = 1$$

$$\frac{y^2}{16} - \frac{x^2}{20} = 1$$

EXAMPLE 5.23:

$$9x^2 - 16y^2 = 144$$

$$\frac{x^2}{16} - \frac{y^2}{9} = 1$$

$$a^2 = 16 \Rightarrow a = \pm 4$$

$$b^2 = 9 \Rightarrow b = \pm 3$$

vertices $(\pm a, 0) = (\pm 4, 0)$

$$b^2 = (ae)^2 - a^2$$

$$(ae)^2 = a^2 + b^2$$

$$= 16 + 9$$

$$(ae)^2 = 25$$

$$ae = \pm 5$$

Focus $(\pm ae, 0) = (\pm 5, 0)$

EXAMPLE 5.24.

$$11x^2 - 25y^2 - 44x + 50y - 256 = 0$$

$$11[x^2 - 4x] - 25[y^2 - 2y] = 256$$

$$11[(x-2)^2 - 4] - 25[(y-1)^2 - 1] = 256$$

$$11(x-2)^2 - 25(y-1)^2 = 256 + 44 - 25$$

$$11(x-2)^2 - 25(y-1)^2 = 275$$

$$\frac{(x-2)^2}{25} - \frac{(y-1)^2}{11} = 1$$

$$a^2 = 25, \quad a = \pm 5$$

$$b^2 = 11, \quad b = \pm \sqrt{11}$$

$$\therefore b^2 = (ae)^2 - a^2$$

$$(ae)^2 = a^2 + b^2 = 25 + 11 = 36$$

$$ae = \pm 6$$

$$ae = 6$$

$$e = \frac{6}{5}$$

Focus $(\pm ae, 0) = (\pm 6, 0)$

$$x-2 = \pm 6 \mid y-1 = 0$$

$$x = \pm 6 + 2 \mid y = 1$$

Focus $(\pm 6 + 2, 1)$

$(8, 1) (-4, 1)$

Foci $ae = 2$, vertices $a = 3$.

$$b^2 = a^2 - (ae)^2$$

$$= 9 - 4$$

$$b^2 = 5$$

Equation of the ellipse

$$\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$$

$$\frac{x^2}{9} + \frac{y^2}{5} = 1.$$

EXAMPLE: 5.19.

$$\frac{FM}{PM} = e$$

$$FM^2 = e^2 PM^2$$

$$(x - ae)^2 + (y - 0)^2 = e^2 \left[\left(x - \frac{a}{e} \right)^2 + 0 \right]$$

$$\text{Foci } (ae, 0) = (2, 3), \quad e = \frac{1}{2}.$$

$$\text{directrix } x = \frac{a}{e} = 7.$$

$$(x - 2)^2 + (y - 3)^2 = \left(\frac{1}{2} \right)^2 [(x - 7)^2 - 0]$$

$$(x - 2)^2 + (y - 3)^2 = \frac{1}{4} (x - 7)^2$$

$$x^2 - 4x + 4 + y^2 - 6y + 9 = \frac{1}{4} (x^2 - 14x + 49)$$

$$4x^2 + 4y^2 - 16x - 24y + 52 = x^2 - 14x + 49$$

$$3x^2 + 4y^2 - 2x - 24y + 3 = 0.$$

$$3\left(x - \frac{1}{3}\right)^2 + 4(y - 3)^2 = \frac{1}{3} + 36 - 3$$

$$3\left(x - \frac{1}{3}\right)^2 + 4(y - 3)^2 = \frac{100}{3}.$$

$$\frac{\left(x - \frac{1}{3}\right)^2}{\frac{100}{9}} + \frac{4(y - 3)^2}{\frac{100}{3}} = 1$$

$$\frac{\left(x - \frac{1}{3}\right)^2}{100/9} + \frac{(y - 3)^2}{100/12} = 1.$$

length of the major axis $= 2a \Rightarrow 2\sqrt{\frac{100}{9}}$

length of the minor axis $= 2b \Rightarrow 2\sqrt{\frac{100}{12}}$

EXAMPLE 5.20.

$$4x^2 + y^2 + 24x - 2y + 21 = 0$$

$$4[x^2 + 6x] + 1[y^2 - 2y] = -21$$

$$4[(x+3)^2 - 9] + 1[(y-1)^2 - 1] = -21$$

$$4(x+3)^2 + 1(y-1)^2 = -21 + 36 + 1$$

$$4(x+3)^2 + (y-1)^2 = 16$$


$$4 \frac{(x+3)^2}{16} + \frac{(y-1)^2}{16} = 1.$$

$$\frac{(x+3)^2}{4} + \frac{(y-1)^2}{16} = 1.$$

$(h, k) = (-3, 1)$ centre

$$a^2 = 16$$

$$b^2 = 4$$



vertices $(0, \pm a)$ $A(-3, 5)$
 $A'(-3, -3)$

$$b^2 = a^2 - (ae)^2$$

$$4 = 16 - 16e^2$$

$$-12 = -16e^2$$

$$\frac{3}{4} = e^2 \quad \boxed{e = \frac{\sqrt{3}}{2}}$$

Foci $(0, \pm ae)$, $ae = 4\left(\frac{\sqrt{3}}{2}\right) = 2\sqrt{3}$

$$x + 3 = 0, \quad y - 1 = \pm 2\sqrt{3}$$

$$x = -3$$

$$y = \pm 2\sqrt{3} + 1$$

Foci $(0, \pm ae) = (-3, \pm 2\sqrt{3} + 1)$

$$ae = 4, \quad a = 5$$

$$5e = 4$$

$$e = 4/5$$

$$\therefore b^2 = a^2 - (ae)^2$$

$$= 25 - 16$$


$$b^2 = 9$$

Equation of the ellipses.

$$\frac{x^2}{b^2} + \frac{y^2}{a^2} = 1.$$

$$\frac{x^2}{9} + \frac{y^2}{25} = 1.$$

Ex E (5.2) (2) (iii).



Length of the Latus rectum $\frac{2b^2}{a} = 8$

$$2b^2 = 8a$$

$$b^2 = 4a$$

$$e = 3/5$$

$$\therefore b^2 = a^2 - (ae)^2$$

$$4a = a^2 [1 - e^2]$$

$$4 = a [1 - 9/25]$$

$$4 = a \left[\frac{16}{25} \right]$$

$$\boxed{a = \frac{25}{4}}$$

$$\therefore b^2 = 4a$$

$$b^2 = 4 \left(\frac{25}{4} \right) = 25$$


Equation of the ellipses

$$\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1.$$

$$\frac{x^2}{\left(\frac{625}{16} \right)} + \frac{y^2}{25} = 1.$$

$$\frac{16x^2}{625} + \frac{y^2}{25} = 1.$$

Ex E 5.2 (2) (iv).



Equation of the ellipses $\frac{x^2}{b^2} + \frac{y^2}{a^2} = 1$

Length of the latus rectum $\frac{2b^2}{a^2} = 4$

$$2b^2 = 4a$$

$$\boxed{b^2 = 2a}$$

distance between foci

$$2ae = 4\sqrt{2}$$

$$ae = 2\sqrt{2}$$

$$\therefore b^2 = a^2 - (ae)^2$$

$$2a = a^2 [1 - e^2]$$

$$2a = a^2 - (2\sqrt{2})^2$$

$$2a = a^2 - 8$$

$$a^2 - 2a - 8 = 0.$$

$$(a-4)(a+2) = 0$$


$$\therefore a = 4 \text{ and } a = -2 \text{ (not possible)}$$

$$b^2 = 2(4) = 8.$$


Equation of the ellipses.

$$\frac{x^2}{8} + \frac{y^2}{16} = 1.$$

EXAMPLE 5.18.



EXAMPLE : 5.15



The parabola open leftward

$$(y-k)^2 = -4a(x-h)$$

$$V(h, k) = (5, -2), F(-a, 0) = (-1, -2)$$

$$a=3$$

$$(y+2)^2 = -4(3)(x-5)$$

$$(y+2)^2 = -12(x-5)$$

$$\text{The parabola } x^2 - 4x - 5y - 1 = 0$$

$$x^2 - 4x - 1 + 5 = 5y + 5$$

$$x^2 - 4x + 4 = 5y + 5$$

$$(x-2)^2 = 5(y+1)$$

 $4a=5$, length of latus rectum is 5


$$a = 5/4$$

$$\text{Focus } (0, a), x-2=0 \quad \left| \begin{array}{l} y+1 = 5/4 \\ y = 1/4 \end{array} \right.$$

$$\text{Focus } (0, a) = (2, 1/4)$$

$$\text{vertices } (h, k) = (2, -1)$$

EXAMPLE : 5.16



The parabola open ward (up).

$$(x-h)^2 = 4a(y-k)$$

$$V(h, k) = (-1, -2), \text{ passes } (3, 6)$$

$$(3+1)^2 = 4a(6+2)$$


$$16 = 4a(8)$$

$$a = 1/2$$

$$(x+1)^2 = 4(1/2)(y+2)$$

$$(x+1)^2 = 2(y+2)$$

EXE 5.2 (2) (i)



The equation of the ellipses.

$$\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$$

$$ae = 3, e = 1/2$$

$$a(1/2) = 3$$

$$a = 6$$

$$\therefore b^2 = a^2 - (ae)^2$$

$$= 36 - 9$$

$$b^2 = 27$$

$$\text{Equation } \frac{x^2}{36} + \frac{y^2}{27} = 1$$

EXE 5.2 (2) (ii)


$$C(0, 0)$$

$$F(0, 3)$$


$$F(0, -3)$$

$$A(0, 4)$$

$$A'(0, -4)$$



EXE 5.2 (i) (i)

Focus $(4,0)$ and directrix $x=-4$ 

The parabola open rightward.

$$(y-k)^2 = 4a(x-h)$$


$$(h,k) = (0,0)$$

$$(a,0) = (4,0)$$

$$(y-0)^2 = 4(4)(x-0)$$

$$y^2 = 16x.$$

EXE 5.2 (i) (ii)



The parabola open downward.

$$(x-h)^2 = -4a(y-k)$$

$$V(h,k) = (0,0)$$

$$(x,y) = (2,-3)$$

$$(2-0)^2 = -4a(-3-0)$$

$$4 = 4a(3)$$


$$a = \frac{1}{3}$$

$$\text{Equation. } (x-0)^2 = -4\left(\frac{1}{3}\right)(y-0)$$

$$x^2 = -\frac{4}{3}y.$$

$$3x^2 = -4y.$$

EXE 5.2 (i) (iii)



The parabola open rightward.


$$(y-k)^2 = 4a(x-h)$$

$$\therefore a=3, V(h,k) = (1,-2)$$

$$(y+2)^2 = 4(3)(x-1)$$

$$(y+2)^2 = 12(x-1)$$

EXE 5.2 (i) (iv)



The parabola open rightward.


$$(y-k)^2 = 4a(x-h)$$

$$V(h,k) = (0,0), \boxed{a=4} \therefore 8a=8$$

$$(y-0)^2 = 4(4)(x-0)$$

$$y^2 = 16x.$$

EXAMPLE 5.14.



The parabola open leftward.

Symmetric as x-axis.

$$F(a,0) = (-\sqrt{2},0), \therefore a=\sqrt{2}.$$

$$V(h,k) = (0,0).$$

$$(y-k)^2 = -4a(x-h)$$

$$(y-0)^2 = -4(\sqrt{2})(x-0)$$

$$y^2 = -4\sqrt{2}x.$$

CONICS:

A conic is the locus of a point which moves in a plane, so that its distance from a fixed point bears a constant ratio to its distance from a fixed line not containing the fixed point.

fixed point $\rightarrow$ focus

fixed line $\rightarrow$ directrix

constant ratio $\rightarrow$ eccentricity.

$e = 1$, parabola $B^2 - 4AC = 0$

$e < 1$, ellipse $B^2 - 4AC < 0$

$e > 1$, hyperbola $B^2 - 4AC > 0$

General second grade equation.

$$Ax^2 + Bxy + Cy^2 + Dx + Ey + F = 0$$

ELLIPSE

$a^2 > b^2$

$b^2 > a^2$

Equation	$\frac{(x-h)^2}{a^2} + \frac{(y-k)^2}{b^2} = 1$	$\frac{(x-h)^2}{b^2} + \frac{(y-k)^2}{a^2} = 1$
Diagram		
Centre	$(0,0)$	$(0,0)$
Focus	$(\pm a, 0)$	$(0, \pm a)$
vertices	$(\pm a, 0)$	$(0, \pm a)$
eccentricity	$e = \sqrt{1 - \frac{b^2}{a^2}}$	$e = \sqrt{1 - \frac{a^2}{b^2}}$
Directrix	$x = \pm \frac{a}{e}$	$y = \pm \frac{a}{e}$

Hyperbola

Equation	$\frac{(x-h)^2}{a^2} - \frac{(y-k)^2}{b^2} = 1$	$\frac{(y-k)^2}{a^2} - \frac{(x-h)^2}{b^2} = 1$
Diagram		
Pocus	$(0, \pm ae)$ $(\pm ae, 0)$	$(0, \pm ae)$
vertices	$(\pm a, 0)$	$(0, \pm a)$
eccentricity	$e = \sqrt{1 + \frac{b^2}{a^2}}$	$e = \sqrt{1 + \frac{b^2}{a^2}}$
Directrix	$x = \pm \frac{a}{e}$	$y = \pm \frac{a}{e}$

Equation	$y^2 = 4ax$	$y^2 = -4ax$	$x^2 = 4ay$	$x^2 = -4ay$
Graph				
vertex	$(0,0)$	$(0,0)$	$(0,0)$	$(0,0)$
focus	$(a,0)$	$(-a,0)$	$(0,a)$	$(0,-a)$
Axis	$y=0$	$y=0$	$x=0$	$x=0$
Equation Directrix	$x=-a$	$x=a$	$y=-a$	$y=a$
Length of the latus rectum	$4a$	$4a$	$4a$	$4a$

⑥

From the diagram


Centre : (0,0)

Radius : $r=1$

Equation of the Circle

$$(x-h)^2 + (y-k)^2 = r^2$$

$$x^2 + y^2 = 1$$



⑦

Area of the Circle = 9π

$$\pi r^2 = 9\pi \Rightarrow \boxed{r^2 = 9}$$

Two diameters are

$$x+y=5 \rightarrow \textcircled{1} \text{ and } x-y=1 \rightarrow \textcircled{2}$$

$$\textcircled{1} + \textcircled{2} \Rightarrow 2x = 6 \Rightarrow \boxed{x = 3}$$

$$\textcircled{1} \Rightarrow 3+y=5 \Rightarrow \boxed{y=2}$$

 $\therefore$ Centre : $(h,k) = (3,2)$

Equation of the Circle

$$(x-h)^2 + (y-k)^2 = r^2$$

$$(x-3)^2 + (y-2)^2 = 9$$

$$x^2 + 9 - 6x + y^2 + 4 - 4y = 9$$

$$x^2 + y^2 - 6x - 4y + 4 = 0$$

⑧

$$y = 2\sqrt{2}x + c \quad \left| \quad x^2 + y^2 = 16 \right.$$

$$m = 2\sqrt{2} \quad \left| \quad a^2 = 16 \right.$$

$$m^2 = (2\sqrt{2})^2 = 8$$

$$\text{Condition : } c^2 = a^2(1+m^2)$$

$$c^2 = 16(1+8) = 16(9)$$

$$c^2 = 144$$

$$\boxed{c = \pm 12}$$

⑨

$$x^2 + y^2 - 6x + 6y - 8 = 0$$

Eqn. of the tangent

$$xx_1 + yy_1 - 6\left(\frac{x+x_1}{2}\right) + 6\left(\frac{y+y_1}{2}\right) - 8 = 0$$

$$xx_1 + yy_1 - 3(x+x_1) + 3(y+y_1) - 8 = 0$$

Here $(x_1, y_1) = (2, 2)$

$$2x + 2y - 3(x+2) + 3(y+2) - 8 = 0$$

$$2x + 2y - 3x - 6 + 3y + 6 - 8 = 0$$

$$\boxed{-x + 5y - 8 = 0}$$

$$\Rightarrow x - 5y + 8 = 0$$

Eqn. of the Normal

$$5x + y + k = 0$$

It passes through (2,2)

$$5(2) + 2 + k = 0$$

$$10 + 2 + k = 0 \Rightarrow \boxed{k = -12}$$

Eqn. of the normal is

$$5x + y - 12 = 0$$

⑩

$$x^2 + y^2 - 5x + 2y - 5 = 0$$

At $(-2, 1)$

$$\textcircled{1} \Rightarrow 4 + 1 + 10 - 2 - 5 = 12 > 0$$

 $\therefore (-2, 1)$ lies outside the circleAt $(0, 0)$

$$\textcircled{1} \Rightarrow 0 + 0 - 0 + 0 - 5 = -5 < 0$$

 $\therefore (0, 0)$ lies inside the circleAt $(-4, -3)$

$$\textcircled{1} \Rightarrow 16 + 9 + 20 - 6 - 5 = 34 > 0$$

 $\therefore (-4, -3)$ lies outside the circle

$$\textcircled{11} \text{ i) } x^2 + (y+2)^2 = 0 \quad [(x-h)^2 + (y-k)^2 = r^2]$$

 $\therefore$ Centre = $(0, -2)$ and $r = 0$

$$\text{ii) } x^2 + y^2 + 2gx + 2fy + c = 0$$

$$x^2 + y^2 + 6x - 4y + 4 = 0$$

$$\therefore 2g = 6 \Rightarrow \boxed{g = 3} \text{ and } 2f = -4 \Rightarrow \boxed{f = -2}$$

$$\boxed{c = 4}$$

 $\therefore$ Centre = $(-g, -f) = (-3, 2)$

$$\text{radius } r = \sqrt{g^2 + f^2 - c} = \sqrt{9 + 4 - 4} = \sqrt{9} = 3$$

$$\text{iii) } x^2 + y^2 - x + 2y - 3 = 0$$

$$2g = -1 \quad | \quad 2f = 2 \quad | \quad c = -3$$

$$g = -1/2 \quad | \quad f = 1$$

Centre : $(-g, -f) = (1/2, -1)$

$$\text{radius } r = \sqrt{g^2 + f^2 - c} = \sqrt{1/4 + 1 + 3} = \sqrt{17/4}$$

$$\boxed{r = \sqrt{17}/2}$$

$$\text{iv) } 2x^2 + 2y^2 - 6x + 4y + 2 = 0$$

$$\div 2 \Rightarrow x^2 + y^2 - 3x + 2y + 1 = 0$$

$$2g = -3 \quad | \quad 2f = 2 \quad | \quad c = 1$$

$$g = -3/2 \quad | \quad f = 1$$

Centre = $(-g, -f) = (3/2, -1)$

$$\text{radius} = \sqrt{g^2 + f^2 - c} = \sqrt{9/4 + 1 - 1} = \sqrt{9/4}$$

$$r = 3/2$$

$$\textcircled{12} \text{ Sol: } 3x^2 + (3-f)xy + 9y^2 - 2fx = 8f^2 \quad \textcircled{1}$$

 $\therefore$ Coeff. of x^2 = Coeff. of y^2

$$3 = 9 \Rightarrow \boxed{f = 3}$$

Again, Coeff. of $xy = 0 \Rightarrow 3-f=0 \Rightarrow \boxed{f=3}$

$$\textcircled{1} \Rightarrow 3x^2 + 3y^2 - 6x - 72 = 0$$

$$\div 3 \Rightarrow x^2 + y^2 - 2x - 24 = 0$$

$$2g = -2 \quad | \quad 2f = 0 \quad | \quad c = -24$$

$$g = -1 \quad | \quad f = 0$$

Centre = $(-g, -f) = (1, 0)$

$$\text{radius } r = \sqrt{g^2 + f^2 - c} = \sqrt{1 + 0 + 24} = \sqrt{25} = 5$$

Example 5.13

Let $O_1(12, 0)$ and $O_2(34, 0)$ be the centres of the semi circle and radius $r = 10$.

Equation of the semicircle

$$(x-h)^2 + (y-k)^2 = r^2$$

$$(x-h)^2 + (y-k)^2 = 100 \quad [r=10]$$

Case (i) : Centre $O_1(12, 0) = (h, k)$

$$(x-12)^2 + (y-0)^2 = 100$$

$$x^2 + 144 - 24x + y^2 - 100 = 0$$

$$x^2 + y^2 - 24x + 44 = 0, \quad y > 0.$$

Case (ii) : Centre $O_2(h, k) = (34, 0)$

$$(x-34)^2 + (y-0)^2 = 100$$

$$x^2 + 1156 - 68x + y^2 - 100 = 0$$

$$x^2 + y^2 - 68x + 1056 = 0, \quad y > 0.$$


Ex 5.1

1) Solution :

$$r = 5, \text{ Centre } = (0, \pm 5)$$

Equation of the circle

$$(x-h)^2 + (y-k)^2 = r^2$$



Here $(h, k) = (0, \pm 5), r = 5$

$$(x-0)^2 + (y \pm 5)^2 = 25$$

$$x^2 + y^2 + 25 \pm 10y = 25$$

$$\boxed{x^2 + y^2 \pm 10y = 0}$$

② Equation of the circle

$$(x-h)^2 + (y-k)^2 = r^2$$

Here $(h, k) = (2, -1)$

$$(x-2)^2 + (y+1)^2 = r^2$$

It passes through $(3, 6)$

$$(3-2)^2 + (6+1)^2 = r^2$$

$$1 + 49 = r^2 \Rightarrow \boxed{r^2 = 50}$$

Equation of the circle

$$(x-2)^2 + (y+1)^2 = 50.$$

③ Equation of the circle

$$(x-h)^2 + (y-k)^2 = r^2$$

Centre is $(h, k) = (-r, -r)$

$$(x+r)^2 + (y+r)^2 = r^2 \rightarrow ①$$

But it passes through $(-4, -2)$

$$(-4+r)^2 + (-2+r)^2 = r^2$$

$$16 + r^2 - 8r + 4 + r^2 - 4r = r^2$$

$$r^2 - 12r + 20 = 0$$

$$(r-10)(r-2) = 0$$

$$r = 10, r = 2$$

$$\begin{array}{r} 20 \\ -12 \\ \hline -2 \end{array}$$

Case (i) $r = 10$

$$\textcircled{1} \Rightarrow (x+10)^2 + (y+10)^2 = 10^2$$

$$x^2 + 100 + 20x + y^2 + 100 + 20y = 100$$

$$x^2 + y^2 + 20x + 20y + 100 = 0$$

Case (ii) $r = 2$

$$\textcircled{1} \Rightarrow (x+2)^2 + (y+2)^2 = 2^2$$

$$x^2 + 4 + 4x + y^2 + 4 + 4y = 4$$

$$x^2 + y^2 + 4x + 4y + 4 = 0.$$

④ Centre $(h, k) = (2, 3)$

Equation of the circle

$$(x-h)^2 + (y-k)^2 = r^2$$

$$(x-2)^2 + (y-3)^2 = r^2 \rightarrow ①$$

Given Lines : $3x - 2y - 1 = 0 \rightarrow ②$

$$4x + y - 27 = 0 \rightarrow ③$$

$$\therefore ② \times 1 \Rightarrow 3x - 2y - 1 = 0$$

$$③ \times 2 \Rightarrow \underline{8x + 2y - 54 = 0}$$

$$11x - 55 = 0$$

$$11x = 55 \Rightarrow \boxed{x = 5}$$

$$③ \Rightarrow 4(5) + y - 27 = 0$$

$$20 + y - 27 = 0 \Rightarrow \boxed{y = 7}$$

$\therefore$ Point of intersection $(5, 7)$.

$$\therefore ① \Rightarrow (5-2)^2 + (7-3)^2 = r^2$$

$$9 + 16 = r^2 \Rightarrow \boxed{r^2 = 25}$$

$$\therefore ① \Rightarrow (x-2)^2 + (y-3)^2 = 25$$

$$x^2 + 4 - 4x + y^2 + 9 - 6y - 25 = 0$$

$$x^2 + y^2 - 4x - 6y - 12 = 0$$

⑤ Equation of the circle

$$(x-x_1)(x-x_2) + (y-y_1)(y-y_2) = 0$$


Here $(x_1, y_1) = (3, 4), (x_2, y_2) = (2, -7)$

$$(x-3)(x-2) + (y-4)(y+7) = 0$$

$$x^2 - 2x - 3x + 6 + y^2 + 7y - 4y - 28 = 0$$

$$x^2 + y^2 - 5x + 3y - 22 = 0$$

Example 5.8



Equation of circle
 $(x-h)^2 + (y-k)^2 = r^2$

i) Centre : $(h,k) = (3,3)$

$$(x-3)^2 + (y-3)^2 = 3^2$$

$$x^2 + 9 - 6x + y^2 + 9 - 6y = 9$$

$$x^2 + y^2 - 6x - 6y + 9 = 0$$

(ii) Centre : $(h,k) = (-3,3)$

$$(x+3)^2 + (y-3)^2 = 3^2$$

$$x^2 + 9 + 6x + y^2 + 9 - 6y = 9$$

$$x^2 + y^2 + 6x - 6y + 9 = 0$$

(iii) Centre : $(h,k) = (-3,-3)$

$$(x+3)^2 + (y+3)^2 = 3^2$$

$$x^2 + 9 + 6x + y^2 + 9 + 6y = 9$$

$$x^2 + y^2 + 6x + 6y + 9 = 0$$

(iv) Centre : $(h,k) = (3,-3)$

$$(x-3)^2 + (y+3)^2 = 3^2$$

$$x^2 + 9 - 6x + y^2 + 9 + 6y = 9$$

$$x^2 + y^2 - 6x + 6y + 9 = 0$$

∴ The req. equations of the circle

$$x^2 + y^2 \pm 6x \pm 6y + 9 = 0.$$

Example 5.9

$$\text{Circle : } 3x^2 + (a+1)y^2 + 6x - 9y + a + 4 = 0$$

∴ Coeff. of x^2 = Coeff. of y^2

$$3 = a + 1 \Rightarrow \boxed{a = 2}$$

∴ Equation of the circle

$$3x^2 + 3y^2 + 6x - 9y + 6 = 0$$

$$\div 3 \Rightarrow x^2 + y^2 + 2x - 3y + 2 = 0$$

$$\text{Here } 2g = 2 \mid 2f = -3$$

$$g = 1 \mid f = -3/2$$

$$\therefore \text{Centre} = (-g, -f) = (-1, 3/2)$$

$$\text{Radius } r = \sqrt{g^2 + f^2 - c} = \sqrt{1 + 9/4 - 2}$$

$$= \sqrt{\frac{4+9-8}{4}} = \sqrt{5/2}$$

Example : 5.10

General equation of the circle

$$x^2 + y^2 + 2gx + 2fy + c = 0 \rightarrow (1)$$

It passes through points

$$(1,1), (2,-1) \text{ and } (3,2)$$

$$(1,1) \Rightarrow 1+1+2g+2f+c=0$$

$$2g+2f+c=-2 \rightarrow (2)$$

$$(2,-1) \Rightarrow 4+1+4g-2f+c=0$$

$$4g-2f+c=-5 \rightarrow (3)$$

$$(3,2) \Rightarrow 9+4+6g+4f+c=0$$

$$6g+4f+c=-13 \rightarrow (4)$$

$$(2)-(3) \Rightarrow -2g+4f=3 \rightarrow (5)$$

$$(4)-(3) \Rightarrow 2g+6f=-8 \rightarrow (6)$$

$$\therefore (5)+(6) \Rightarrow 10f=-5$$

$$f = \frac{-5}{10} = -\frac{1}{2}$$

$$\boxed{f = -1/2}$$

$$(6) \Rightarrow 2g+6(-1/2)=-8$$

$$2g-3=-8 \Rightarrow 2g=-5$$

$$\boxed{g = -5/2}$$

$$(2) \Rightarrow 2(-5/2)+2(-1/2)+c=-2$$

$$-5-1+c=-2$$

$$c=-2+6$$

$$\boxed{c=4}$$

$$\therefore (1) \Rightarrow x^2 + y^2 + 2(-5/2)x + 2(-1/2)y + 4 = 0$$

$$x^2 + y^2 - 5x - y + 4 = 0.$$

Example 5.11

$$x^2 + y^2 = 25 \text{ at } P(-3,4)$$

Equation of the tangent

$$xx_1 + yy_1 = 25$$

$$(x_1, y_1) = (-3, 4)$$

$$x(-3) + y(4) = 25$$

$$\boxed{-3x + 4y = 25}$$

Equation of the normal

$$4x + 3y = k$$

It passes through $(-3,4)$

$$4(-3) + 3(4) = k$$

$$-12 + 12 = k \Rightarrow k = 0$$

∴ Equation of the normal is

$$\boxed{4x + 3y = 0}$$

Example 5.12

$$y = 4x + c \mid x^2 + y^2 = 9$$

$$m = 4 \mid a^2 = 9$$

The condition is $c^2 = a^2(1+m^2)$

$$c^2 = 9(1+16)$$

$$c^2 = 9(17) \Rightarrow \boxed{c = \pm 3\sqrt{17}}$$

5. TWO DIMENSIONAL ANALYTICAL GEOMETRY-II

Example : 5.1

Equation of the circle

$$(x-h)^2 + (y-k)^2 = r^2$$

$$(h,k) = (-3,-4) \text{ and } r=3$$

$$(x+3)^2 + (y+4)^2 = 3^2$$

$$x^2 + 6x + 9 + y^2 + 8y + 16 = 9$$

$$x^2 + y^2 + 6x + 8y + 16 = 0$$

Example 5.2

$$\text{Chord : } 3x + y + 5 = 0$$

$$\text{Circle : } x^2 + y^2 = 16 \Rightarrow x^2 + y^2 - 16 = 0$$

The req. equation of the circle

$$(x^2 + y^2 - 16) + \lambda(3x + y + 5) = 0$$

$$x^2 + y^2 - 16 + 3\lambda x + \lambda y + 5\lambda = 0 \quad \text{--- (1)}$$

$$\text{Here } 2g = 3\lambda \quad 2f = \lambda$$

$$g = \frac{3\lambda}{2} \quad f = \frac{\lambda}{2}$$

$$\text{Centre} = (-g, -f) = \left(-\frac{3\lambda}{2}, -\frac{\lambda}{2}\right)$$

$\therefore$ Centre lies on the chord

$$3x + y + 5 = 0.$$

$$\therefore 3\left(-\frac{3\lambda}{2}\right) + \left(-\frac{\lambda}{2}\right) + 5 = 0$$

$$-\frac{9}{2} - \frac{\lambda}{2} + 5 = 0$$

$$-\frac{10}{2}\lambda + 5 = 0$$

$$-5\lambda = -5 \Rightarrow \boxed{\lambda = 1}$$

$$\textcircled{1} \Rightarrow x^2 + y^2 - 16 + 3x + y + 5 = 0$$

$$x^2 + y^2 + 3x + y - 11 = 0$$

Example : 5.3

$$\text{Circle : } x^2 + y^2 - 6x + 4y + c = 0$$

$$\text{Here } 2g = -6 \quad 2f = 4$$

$$g = -3 \quad f = 2$$

$$\text{Centre } (-g, -f) = (3, -2) \text{ which}$$

lies on $x + y - 1 = 0$.

$\therefore x + y - 1 = 0$ passes through the centre.

$\therefore x + y - 1 = 0$ is a diameter of the circle for all possible value of c .

Example : 5.4.

Equation of the circle [diameter form]

$$(x-x_1)(x-x_2) + (y-y_1)(y-y_2) = 0$$

$$(x_1, y_1) = (-4, -2), (x_2, y_2) = (1, 1)$$

$$(x+4)(x-1) + (y+2)(y-1) = 0$$

$$x^2 - x + 4x - 4 + y^2 - y + 2y - 2 = 0$$

$$x^2 + y^2 + 3x + y - 6 = 0.$$

Example 5.5

point (2,3)

$$x^2 + y^2 - 6x - 8y + 12$$

$$= 4 + 9 - 6(2) - 8(3) + 12$$

$$= 4 + 9 - 12 - 24 + 12$$

$$= -11 < 0.$$

$\therefore (2,3)$ lies inside the circle.

Example 5.6.

$$3x + 4y = 12$$

$$\div 12 \Rightarrow \frac{x}{4} + \frac{y}{3} = 1$$

$$\therefore A(4,0), B(0,3)$$

Equation of the circle

$$(x-x_1)(x-x_2) + (y-y_1)(y-y_2) = 0$$

$$(x_1, y_1) = (4,0), (x_2, y_2) = (0,3)$$

$$(x-4)(x-0) + (y-0)(y-3) = 0$$

$$x^2 - 4x + y^2 - 3y = 0$$

$$\Rightarrow x^2 + y^2 - 4x - 3y = 0.$$

Example 5.7

$$\text{Centre} = (2,1)$$

$$3x + 4y + 10 = 0 \text{ cuts a}$$

Chord AB on the circle.

Let M be the midpoint of AB.

$$\therefore AM = MB = 3 \quad [\because AB = 6]$$

$$CM = \frac{|3(2) + 4(1) + 10|}{\sqrt{3^2 + 4^2}}$$

$$= \frac{|6 + 4 + 10|}{\sqrt{9 + 16}} = \frac{20}{5} = 4$$

$$\Delta BMC, AC^2 = AM^2 + MC^2 = 3^2 + 4^2 = 9 + 16 = 25$$

$$AC^2 = 25 \Rightarrow AC = 5 \text{ (radius)}$$

Equation of the circle $(x-h)^2 + (y-k)^2 = r^2$

$$(x-2)^2 + (y-1)^2 = 25 \Rightarrow x^2 + y^2 - 4x - 2y - 24 = 0$$

