



# Padalsalai's Telegram Groups!

( தலைப்பிற்கு கீழே உள்ள லிங்கை கிளிக் செய்து குழுவில் இணையவும்! )

- **Padalsalai's NEWS - Group**  
[https://t.me/joinchat/NIfCqVRBNj9hhV4wu6\\_NqA](https://t.me/joinchat/NIfCqVRBNj9hhV4wu6_NqA)
- **Padalsalai's Channel - Group**  
<https://t.me/padasalaichannel>
- **Lesson Plan - Group**  
<https://t.me/joinchat/NIfCqVWwo5iL-21gpzrXLw>
- **12th Standard - Group**  
[https://t.me/Padalsalai\\_12th](https://t.me/Padalsalai_12th)
- **11th Standard - Group**  
[https://t.me/Padalsalai\\_11th](https://t.me/Padalsalai_11th)
- **10th Standard - Group**  
[https://t.me/Padalsalai\\_10th](https://t.me/Padalsalai_10th)
- **9th Standard - Group**  
[https://t.me/Padalsalai\\_9th](https://t.me/Padalsalai_9th)
- **6th to 8th Standard - Group**  
[https://t.me/Padalsalai\\_6to8](https://t.me/Padalsalai_6to8)
- **1st to 5th Standard - Group**  
[https://t.me/Padalsalai\\_1to5](https://t.me/Padalsalai_1to5)
- **TET - Group**  
[https://t.me/Padalsalai\\_TET](https://t.me/Padalsalai_TET)
- **PGTRB - Group**  
[https://t.me/Padalsalai\\_PGTRB](https://t.me/Padalsalai_PGTRB)
- **TNPSC - Group**  
[https://t.me/Padalsalai\\_TNPSC](https://t.me/Padalsalai_TNPSC)

## Ionic Equilibrium



### EVALUATION

**Choose the correct answer:**

- Concentration of the  $\text{Ag}^+$  ions in a saturated solution of  $\text{Ag}_2\text{C}_2\text{O}_4$  is  $2.24 \times 10^{-4} \text{ mol L}^{-1}$  solubility product of  $\text{Ag}_2\text{C}_2\text{O}_4$  is (NEET – 2017)
  - $2.42 \times 10^{-8} \text{ mol}^3\text{L}^{-3}$
  - $2.66 \times 10^{-12} \text{ mol}^3\text{L}^{-3}$
  - $4.5 \times 10^{-11} \text{ mol}^3\text{L}^{-3}$
  - $5.619 \times 10^{-12} \text{ mol}^3\text{L}^{-3}$
- Following solutions were prepared by mixing different volumes of NaOH of HCl different concentrations. (NEET – 2018)
  - 60 mL  $\frac{M}{10}$  HCl + 40 mL  $\frac{M}{10}$  NaOH
  - 55 mL  $\frac{M}{10}$  HCl + 45 mL  $\frac{M}{10}$  NaOH
  - 75 mL  $\frac{M}{5}$  HCl + 25 mL  $\frac{M}{5}$  NaOH
  - 100 mL  $\frac{M}{10}$  HCl + 100 mL  $\frac{M}{10}$  NaOH

pH of which one of them will be equal to 1?

  - iv
  - i
  - ii
  - iii
- The solubility of  $\text{BaSO}_4$  in water is  $2.42 \times 10^{-3} \text{ g L}^{-1}$  at 298K. The value of its solubility product ( $K_{\text{sp}}$ ) will be (NEET – 2018). (Given molar mass of  $\text{BaSO}_4 = 233 \text{ g mol}^{-1}$ )
  - $1.08 \times 10^{-14} \text{ mol}^2\text{L}^{-2}$
  - $1.08 \times 10^{-12} \text{ mol}^2\text{L}^{-2}$
  - $1.08 \times 10^{-10} \text{ mol}^2\text{L}^{-2}$
  - $1.08 \times 10^{-8} \text{ mol}^2\text{L}^{-2}$
- pH of a saturated solution of  $\text{Ca}(\text{OH})_2$  is 9. The Solubility product ( $K_{\text{sp}}$ ) of  $\text{Ca}(\text{OH})_2$ 
  - $0.5 \times 10^{-15}$
  - $0.25 \times 10^{-10}$
  - $0.125 \times 10^{-15}$
  - $0.5 \times 10^{-10}$
- Conjugate base for Bronsted acids  $\text{H}_2\text{O}$  and  $\text{HF}$  are
  - $\text{OH}^-$  and  $\text{H}_2\text{FH}^+$ , respectively
  - $\text{H}_3\text{O}^+$  and  $\text{F}^-$ , respectively
  - $\text{OH}^-$  and  $\text{F}^-$ , respectively
  - $\text{H}_3\text{O}^+$  and  $\text{H}_2\text{F}^+$ , respectively
- Which will make basic buffer?
  - 50 mL of 0.1M NaOH + 25 mL of 0.1M  $\text{CH}_3\text{COOH}$
  - 100 mL of 0.1M  $\text{CH}_3\text{COOH}$  + 100 mL of 0.1M  $\text{NH}_4\text{OH}$
  - 100 mL of 0.1M HCl + 200 mL of 0.1M  $\text{NH}_4\text{OH}$
  - 100 mL of 0.1M HCl + 100 mL of 0.1M NaOH
- Which of the following fluoro compounds is most likely to behave as a Lewis base? NEET – 2016

- a)  $\text{BF}_3$                       b)  $\text{PF}_3$                       c)  $\text{CF}_4$                       d)  $\text{SiF}_4$
8. Which of these is not likely to act as Lewis base?
- a)  $\text{BF}_3$                       b)  $\text{PF}_3$                       c)  $\text{CO}$                       d)  $\text{F}^-$
9. What is the decreasing order of strength of bases?  
 $\text{OH}^-$ ,  $\text{NH}_2^-$ ,  $\text{H}-\text{C}\equiv\text{C}^-$  and  $\text{CH}_3-\text{CH}_2^-$
- a)  $\text{OH}^- > \text{NH}_2^- > \text{H}-\text{C}\equiv\text{C}^- > \text{CH}_3-\text{CH}_2^-$       b)  $\text{NH}_2^- > \text{OH}^- > \text{CH}_3-\text{CH}_2^- > \text{H}-\text{C}\equiv\text{C}^-$   
 c)  $\text{CH}_3-\text{CH}_2^- > \text{NH}_2^- > \text{H}-\text{C}\equiv\text{C}^- > \text{OH}^-$       d)  $\text{OH}^- > \text{H}-\text{C}\equiv\text{C}^- > \text{CH}_3-\text{CH}_2^- > \text{NH}_2^-$
10. The aqueous solutions of sodium formate, anilinium chloride and potassium cyanide are respectively
- a) acidic, acidic, basic                      b) basic, acidic, basic  
 c) basic, neutral, basic                      d) none of these
11. The percentage of pyridine ( $\text{C}_5\text{H}_5\text{N}$ ) that forms pyridinium ion ( $\text{C}_5\text{H}_5\text{NH}^+$ ) in a 0.10M aqueous pyridine solution ( $K_b$  for  $\text{C}_5\text{H}_5\text{N} = 1.7 \times 10^{-9}$ ) is
- a) 0.006%                      b) 0.013%                      c) 0.77%                      d) 1.6%
12. Equal volumes of three acid solutions of pH 1, 2 and 3 are mixed in a vessel. What will be the  $\text{H}^+$  ion concentration in the mixture?
- a)  $3.7 \times 10^{-2}$                       b)  $10^{-6}$                       c) 0.111                      d) none of these
13. The solubility of  $\text{AgCl (s)}$  with solubility product  $1.6 \times 10^{-10}$  in 0.1M  $\text{NaCl}$  solution would be
- a)  $1.26 \times 10^{-5}\text{M}$                       b)  $1.6 \times 10^{-9}\text{M}$                       c)  $1.6 \times 10^{-11}\text{M}$                       d) Zero
14. If the solubility product of lead iodide is  $3.2 \times 10^{-8}$ , its solubility will be
- a)  $2 \times 10^{-3}\text{M}$                       b)  $4 \times 10^{-4}\text{M}$                       c)  $1.6 \times 10^{-5}\text{M}$                       d)  $1.8 \times 10^{-5}\text{M}$
15. Using Gibb's free energy change,  $\Delta G^\circ = 57.34 \text{ kJ mol}^{-1}$ , for the reaction,  
 $\text{X}_2\text{Y(s)} \rightleftharpoons 2\text{X}^+(\text{aq}) + \text{Y}^{2-}(\text{aq})$  calculate the solubility product of  $\text{X}_2\text{Y}$  in water at 300 K ( $R = 8.3 \text{ J K}^{-1}\text{Mol}^{-1}$ )
- a)  $10^{-10}$                       b)  $10^{-12}$   
 c)  $10^{-14}$                       d) can not be calculated from the given data
16.  $\text{MY}$  and  $\text{NY}_3$ , are insoluble salts and have the same  $K_{sp}$  values of  $6.2 \times 10^{-13}$  at room temperature. Which statement would be true with regard to  $\text{MY}$  and  $\text{NY}_3$ ?
- a) The salts  $\text{MY}$  and  $\text{NY}_3$  are more soluble in 0.5M  $\text{KY}$  than in pure water  
 b) The addition of the salt of  $\text{KY}$  to the suspension of  $\text{MY}$  and  $\text{NY}_3$  will have no effect on their solubility's  
 c) The molar solubilities of  $\text{MY}$  and  $\text{NY}_3$  in water are identical  
 d) The molar solubility of  $\text{MY}$  in water is less than that of  $\text{NY}_3$

17. What is the pH of the resulting solution when equal volumes of 0.1M NaOH and 0.01M HCl are mixed?
- a) 2.0                      b) 3                      c) 7.0                      d) 12.65
18. The dissociation constant of a weak acid is  $1 \times 10^{-3}$ . In order to prepare a buffer solution with a pH = 4, the  $\frac{[\text{Acid}]}{[\text{Salt}]}$  ratio should be
- a) 4:3                      b) 3:4                      c) 10:1                      d) 1:10
19. The pH of  $10^{-5}$ M KOH solution will be
- a) 9                      b) 5                      c) 19                      d) none of these
20.  $\text{H}_2\text{PO}_4^-$  the conjugate base of
- a)  $\text{PO}_4^{3-}$                       b)  $\text{P}_2\text{O}_5$                       c)  $\text{H}_3\text{PO}_4$                       d)  $\text{HPO}_4^{2-}$
21. Which of the following can act as Lowry – Bronsted acid as well as base?
- a) HCl                      b)  $\text{SO}_4^{2-}$                       c)  $\text{HPO}_4^{2-}$                       d)  $\text{Br}^-$
22. The pH of an aqueous solution is Zero. The solution is
- a) slightly acidic      b) strongly acidic      c) neutral      d) basic
23. The hydrogen ion concentration of a buffer solution consisting of a weak acid and its salts is given by
- a)  $[\text{H}^+] = \frac{K_a[\text{acid}]}{[\text{salt}]}$       b)  $[\text{H}^+] = K_a[\text{salt}]$       c)  $[\text{H}^+] = K_a[\text{acid}]$       d)  $[\text{H}^+] = \frac{K_a[\text{salt}]}{[\text{acid}]}$
24. Which of the following relation is correct for degree of hydrolysis of ammonium acetate?
- a)  $h = \sqrt{\frac{K_h}{C}}$       b)  $h = \sqrt{\frac{K_a}{K_b}}$       c)  $h = \sqrt{\frac{K_h}{K_a \cdot K_b}}$       d)  $h = \sqrt{\frac{K_a \cdot K_b}{K_h}}$
25. Dissociation constant of  $\text{NH}_4\text{OH}$  is  $1.8 \times 10^{-5}$  the hydrolysis constant of  $\text{NH}_4\text{Cl}$  would be
- a)  $1.8 \times 10^{-19}$       b)  $5.55 \times 10^{-10}$       c)  $5.55 \times 10^{-5}$       d)  $1.80 \times 10^{-5}$

*Do not give up, the beginning is*

*always the hardest...*

## *Electro Chemistry*





### EVALUATION

Choose the correct answer:

1. The number of electrons that have a total charge of 9650 coulombs is

- |                           |                            |
|---------------------------|----------------------------|
| a) $6.22 \times 10^{23}$  | b) $6.022 \times 10^{24}$  |
| c) $6.022 \times 10^{22}$ | c) $6.022 \times 10^{-34}$ |

2. Consider the following half cell reactions:



The  $E^\circ$  for the reaction  $3\text{Mn}^{2+} \rightarrow \text{Mn} + 2\text{Mn}^{3+}$ , and the possibility of the forward reaction are respectively.

- |                          |                              |
|--------------------------|------------------------------|
| a) 2.69V and spontaneous | b) -2.69 and non spontaneous |
| c) 0.33V and Spontaneous | d) 4.18V and non spontaneous |

3. The button cell used in watches function as follows

$\text{Zn (s)} + \text{Ag}_2\text{O (s)} + \text{H}_2\text{O (l)} \rightleftharpoons 2\text{Ag (s)} + \text{Zn}^{2+}(\text{aq}) + 2\text{OH}^-(\text{aq})$  the half cell potentials are  $\text{Ag}_2\text{O (s)} + \text{H}_2\text{O (l)} + 2\text{e}^- \rightarrow 2\text{Ag (s)} + 2\text{OH}^-(\text{aq}) \quad E^\circ = 0.34\text{V}$  The cell potential will be

- a) 0.84V                      b) 1.34V                      c) 1.10V                      d) 0.42V
4. The molar conductivity of a  $0.5 \text{ mol dm}^{-3}$  solution of  $\text{AgNO}_3$  with electrolytic conductivity of  $5.76 \times 10^{-3} \text{ S cm}^{-1}$  at 298 K is
- a)  $2.88 \text{ S cm}^2 \text{ mol}^{-1}$                       b)  $11.52 \text{ S cm}^2 \text{ mol}^{-1}$   
 c)  $0.086 \text{ S cm}^2 \text{ mol}^{-1}$                       d)  $28.8 \text{ S cm}^2 \text{ mol}^{-1}$

5.

Electrolyte	KCl	$\text{KNO}_3$	HCl	NaOAC	NaCl
$\Lambda_{\infty}$ ( $\text{S cm}^2 \text{ mol}^{-1}$ )	149.9	145.0	426.2	91.0	126.5

Calculate  $\Lambda_{\text{HOAC}}^{\circ}$  using appropriate molar conductances of the electrolytes listed above at infinite dilution in water at  $25^{\circ}\text{C}$ .

- a) 517.2                      b) 552.7                      c) 390.7                      d) 217.5
6. Faradays constant is defined as
- a) charge carried by 1 electron  
 b) charge carried by one mole of electrons  
 c) charge required to deposit one mole of substance  
 d) charge carried by  $6.22 \times 10^{19}$  electrons.
7. How many faradays of electricity are required for the following reaction to occur  
 $\text{MnO}_4^- \rightarrow \text{Mn}^{2+}$
- a) 5F                      b) 3F                      c) 1F                      d) 7F
8. A current strength of 3.86 A was passed through molten Calcium oxide for 41 minutes and 40 seconds. The mass of Calcium in grams deposited at the cathode is (atomic mass of Ca is 40g / mol and  $1\text{F} = 96500\text{C}$ ).
- a) 4                      b) 2                      c) 8                      d) 6
9. During electrolysis of molten sodium chloride, the time required to produce 0.1mole of chlorine gas using a current of 3A is
- a) 55 minutes                      b) 107.2 minutes                      c) 220 minutes                      d) 330 minutes
10. The number of electrons delivered at the cathode during electrolysis by a current of 1A in 60 seconds is (charge of electron =  $1.6 \times 10^{-19} \text{ C}$ )
- a)  $6.22 \times 10^{23}$                       b)  $6.022 \times 10^{20}$                       c)  $3.75 \times 10^{20}$                       d)  $7.48 \times 10^{23}$
11. Which of the following electrolytic solution has the least specific conductance
- a) 2N                      b) 0.002N                      c) 0.02N                      d) 0.2N
12. While charging lead storage battery
- a)  $\text{PbSO}_4$  on cathode is reduced to Pb                      b)  $\text{PbSO}_4$  on anode is oxidised to  $\text{PbO}_2$

- c)  $\text{PbSO}_4$  on anode is reduced to Pb                      d)  $\text{PbSO}_4$  on cathode is oxidised to Pb
13. Among the following cells  
 I) Leclanche cell  
 II) Nickel – Cadmium cell  
 III) Lead storage battery  
 IV) Mercury cell  
 Primary cells are  
 a) I and IV                      b) I and III                      c) III and IV                      d) II and III
14. Zinc can be coated on iron to produce galvanized iron but the reverse is not possible. It is because  
 a) Zinc is lighter than iron  
 b) Zinc has lower melting point than iron  
 c) Zinc has lower negative electrode potential than iron  
 d) Zinc has higher negative electrode potential than iron
15. Assertion : pure iron when heated in dry air is converted with a layer of rust.  
 Reason : Rust has the composition  $\text{Fe}_3\text{O}_4$   
 a) if both assertion and reason are true and reason is the correct explanation of assertion.  
 b) if both assertion and reason are true but reason is not the correct explanation of assertion.  
 c) assertion is true but reason is false  
 d) both assertion and reason are false.
16. In  $\text{H}_2$ - $\text{O}_2$  fuel cell the reaction occurs at cathode is  
 a)  $\text{O}_2(\text{g}) + 2\text{H}_2\text{O}(\text{l}) + 4\text{e}^- \rightarrow 4\text{OH}^-(\text{aq})$   
 b)  $\text{H}^+(\text{aq}) + \text{OH}^-(\text{aq}) \rightarrow \text{H}_2\text{O}(\text{l})$   
 c)  $2\text{H}_2(\text{g}) + \text{O}_2(\text{g}) \rightarrow 2\text{H}_2\text{O}(\text{g})$   
 d)  $\text{H}^+ + \text{e}^- \rightarrow \frac{1}{2} \text{H}_2$
17. The equivalent conductance of  $\frac{M}{36}$  solution of a weak monobasic acid is 6 mho  $\text{cm}^2$  equivalent $^{-1}$  and at infinite dilution is 400 mho  $\text{cm}^2$  equivalent $^{-1}$ . The dissociation constant of this acid is  
 a)  $1.25 \times 10^{-6}$                       b)  $6.25 \times 10^{-6}$                       c)  $1.25 \times 10^{-4}$                       d)  $6.25 \times 10^{-5}$
18. A conductivity cell has been calibrated with a 0.01M, 1:1 electrolytic solution (specific conductance ( $\kappa = 1.25 \times 10^{-3} \text{ S cm}^{-1}$ ) in the cell and the measured resistance was 800  $\Omega$  at 25°C. The cell constant is,  
 a)  $10^{-1} \text{ cm}^{-1}$                       b)  $10^1 \text{ cm}^{-1}$                       c)  $1 \text{ cm}^{-1}$                       d)  $5.7 \times 10^{-12}$

19. Conductivity of a saturated solution of a sparingly soluble salt AB (1:1 electrolyte) at 298K is  $1.85 \times 10^{-5} \text{ S m}^{-1}$ . Solubility product of the salt AB at 298K  $(\Lambda_m^\circ)_{AB} = 14 \times 10^{-3} \text{ S m}^2 \text{ mol}^{-1}$ .
- a)  $5.7 \times 10^{-12}$       b)  $1.32 \times 10^{-12}$       c)  $7.5 \times 10^{-12}$       d)  $1.74 \times 10^{-12}$
20. In the electrochemical cell:  $\text{Zn}|\text{ZnSO}_4 (0.01\text{M})||\text{CuSO}_4 (1.0\text{M})|\text{Cu}$ , the emf of this Daniel cell is  $E_1$ . When the concentration of  $\text{ZnSO}_4$  is changed to 1.0M and that  $\text{CuSO}_4$  changed to 0.01M, the emf changes to  $E_2$ . From the above, which one is the relationship between  $E_1$  and  $E_2$ ?
- a)  $E_1 < E_2$       b)  $E_1 > E_2$       c)  $E_2 \geq E_1$       d)  $E_1 = E_2$
21. Consider the change in oxidation state of Bromine corresponding to different emf values as shown in the diagram below:
- $$\text{BrO}_4^- \xrightarrow{1.82\text{V}} \text{BrO}_3^- \xrightarrow{1.5\text{V}} \text{HBrO} \xrightarrow{1.595\text{V}} \text{Br}_2 \xrightarrow{1.0652\text{V}} \text{Br}^-$$
- Then the species undergoing disproportionation is
- a)  $\text{Br}_2$       b)  $\text{BrO}_4^-$       c)  $\text{BrO}_3^-$       d)  $\text{HBrO}$
22. For the cell reaction
- $$2\text{Fe}^{3+}(\text{aq}) + 2\text{I}^{-}(\text{aq}) \rightarrow 2\text{Fe}^{2+}(\text{aq}) + \text{I}_2(\text{aq})$$
- $E_{\text{cell}}^\circ = 0.24\text{V}$  at 298K. The standard Gibbs energy ( $\Delta_r G^\circ$ ) of the cell reactions is :
- a)  $-46.32 \text{ KJ mol}^{-1}$       b)  $-23.16 \text{ KJ mol}^{-1}$       c)  $46.32 \text{ KJ mol}^{-1}$       d)  $23.16 \text{ KJ mol}^{-1}$
23. A certain current liberated 0.504gm of hydrogen in 2 hours. How many grams of copper can be liberated by the same current flowing for the same time through copper sulphate solution
- a) 31.75      b) 15.8      c) 7.5      d) 63.5
24. A gas X at 1 atm is bubbled through a solution containing a mixture of  $1\text{M Y}^-$  and  $1\text{M Z}^-$  at  $25^\circ\text{C}$ . If the reduction potential of  $\text{Z} > \text{Y} > \text{X}$ , then
- a) Y will oxidize X and not Z      b) Y will oxidize Z and not X  
d) Y will oxidize both X and Z      d) Y will reduce both X and Z
25. Cell equation :  $\text{A} + 2\text{B}^+ \rightarrow \text{A}^{2+} + 2\text{B}$ ;  
 $\text{A}^{2+} + 2\text{e}^- \rightarrow \text{A}$     $E^\circ = +0.34\text{V}$  and  $\log_{10} K = 15.6$  at 300K for cell reactions find  $E^\circ$  for  
 $\text{B}^+ + \text{e}^- \rightarrow \text{B}$  (AIIMS – 2018)
- a) 0.80      b) 1.26      c) -0.54      d) -10.94



## Surface Chemistry



### EVALUATION

Choose the correct answer:

1. For Freundlich isotherm a graph of  $\log \frac{x}{m}$  is plotted against  $\log p$ . The slope of the line and its y – axis intercept respectively corresponds to  
 a)  $\frac{1}{n}$ , k                      b)  $\log \frac{1}{n}$ , k                      c)  $\frac{1}{n}$ ,  $\log k$                       d)  $\log \frac{1}{n}$ ,  $\log k$
2. Which of the following is incorrect for physisorption?  
 a) reversible                      b) increases with increase in temperature  
 c) low heat of adsorption                      d) increases with increase in surface area
3. Which one of the following characteristics are associated with adsorption? (NEET)  
 a)  $\Delta G$  and  $\Delta H$  are negative but  $\Delta S$  is positive  
 b)  $\Delta G$  and  $\Delta S$  are negative but  $\Delta H$  is positive  
 c)  $\Delta G$  is negative but  $\Delta H$  and  $\Delta S$  are positive      d)  $\Delta G$ ,  $\Delta H$  and  $\Delta S$  all are negative.
4. Fog is colloidal solution of  
 a) solid in gas                      b) gas in gas                      c) liquid in gas                      d) gas in liquid
5. Assertion : Coagulation power of  $Al^{3+}$  is more than  $Na^+$ .  
 Reason : greater the valency of the flocculating ion added, greater is its power to cause precipitation  
 a) if both assertion and reason are true and reason is the correct explanation of assertion.  
 b) if both assertion and reason are true but reason is not the correct explanation of assertion.  
 c) assertion is true but reason is false                      d) both assertion and reason are false.
6. Statement :  
 To stop bleeding from an injury, ferric chloride can be applied. Which comment about

the statement is justified?

- a) It is not true, ferric chloride is a poison.
- b) It is true,  $\text{Fe}^{3+}$  ions coagulate blood which is a negatively charged sol
- c) It is not true; ferric chloride is ionic and gets into the blood stream.
- d) It is true, coagulation takes place because of formation of negatively charged sol with  $\text{Cl}^-$ .

7. Hair cream is

- a) gel
- b) emulsion
- c) solid sol
- d) sol.

8. Which one of the following is correctly matched?

a) Emulsion	–	Smoke
b) Gel	–	butter
c) foam	–	Mist
d) whipped cream	–	sol

9. The most effective electrolyte for the coagulation of  $\text{As}_2\text{S}_3$  Sol is

- a)  $\text{NaCl}$
- b)  $\text{Ba}(\text{NO}_3)_2$
- c)  $\text{K}_3[\text{Fe}(\text{CN})_6]$
- d)  $\text{Al}_2(\text{SO}_4)_3$

10. Which one of the is not a surfactant?

- a)  $\text{CH}_3 - (\text{CH}_2)_{15} - \overset{+}{\text{N}} - (\text{CH}_3)_2 \text{CH}_2\text{Br}$
- b)  $\text{CH}_3 - (\text{CH}_2)_{15} - \text{NH}_2$
- c)  $\text{CH}_3 - (\text{CH}_2)_{16} - \text{CH}_2 - \text{OSO}_2^- \text{Na}^+$
- d)  $\text{OHC} - (\text{CH}_2)_{14} - \text{CH}_2 - \text{COO}^- \text{Na}^+$

11. The phenomenon observed when a beam of light is passed through a colloidal solution is

- a) Cataphoresis
- b) Electrophoresis
- c) Coagulation
- d) Tyndall effect

12. In an electrical field, the particles of a colloidal system move towards cathode. The coagulation of the same sol is studied using  $\text{K}_2\text{SO}_4$  (i),  $\text{Na}_3\text{PO}_4$  (ii),  $\text{K}_4[\text{Fe}(\text{CN})_6]$  (iii) and  $\text{NaCl}$  (iv) Their coagulating power should be

- a)  $\text{II} > \text{I} > \text{IV} > \text{III}$
- b)  $\text{III} > \text{II} > \text{I} > \text{IV}$
- c)  $\text{I} > \text{II} > \text{III} > \text{IV}$
- d) none of these

13. Collodion is a 4% solution of which one of the following compounds in alcohol – ether mixture?

- a) Nitroglycerine
- b) Cellulose acetate
- c) Glycoldinitrate
- d) Nitrocellulose

14. Which one of the following is an example for homogeneous catalysis?

- a) manufacture of ammonia by Haber's process
- b) manufacture of sulphuric acid by contact process
- c) hydrogenation of oil
- d) Hydrolysis of sucrose in presence of dil  $\text{HCl}$

15. Match the following

A) $V_2O_5$	i) High density polyethylene
B) Ziegler – Natta	ii) PAN
C) Peroxide	iii) $NH_3$
D) Finely divided Fe	iv) $H_2SO_4$

A	B	C	D
a) (iv)	(i)	(ii)	(iii)
b) (i)	(ii)	(iv)	(iii)
c) (ii)	(iii)	(iv)	(i)
d) (iii)	(iv)	(ii)	(i)

16. The coagulation values in millimoles per litre of the electrolytes used for the coagulation of  $As_2S_3$  are given below

(I) (NaCl)=52                      (II) ((BaCl<sub>2</sub>)=0.69                      (III) (MgSO<sub>4</sub>)=0.22

The correct order of their coagulating power is

a) III > II > I                      b) I > II > III                      c) I > III > II                      d) II > III > I

17. Adsorption of a gas on solid metal surface is spontaneous and exothermic, then

a)  $\Delta H$  increases                      b)  $\Delta S$  increases                      c)  $\Delta G$  increases                      d)  $\Delta S$  decreases

18. If x is the amount of adsorbate and m is the amount of adsorbent, which of the following relations is not related to adsorption process?

a)  $\frac{x}{m} = f(P)$  at constant T                      b)  $\frac{x}{m} = f(T)$  at constant P  
c)  $P = f(T)$  at constant  $\frac{x}{m}$                       d)  $\frac{x}{m} = PT$

19. On which of the following properties does the coagulating power of an ion depend ? (NEET – 2018)

a) Both magnitude and sign of the charge on the ion.  
b) Size of the ion alone  
c) the magnitude of the charge on the ion alone  
d) the sign of charge on the ion alone.

20. Match the following

A) Pure nitrogen	i) Chlorine
B) Haber process	ii) Sulphuric acid
C) Contact process	iii) Ammonia
D) Deacons Process	iv) sodium azide (or) Barium azide

Which of the following is the correct option?

	A	B	C	D
a)	(i)	(ii)	(iii)	(iv)
b)	(ii)	(iv)	(i)	(iii)
c)	(iii)	(iv)	(ii)	(i)
d)	(iv)	(iii)	(ii)	(i)


## Hydroxy Compounds and Ethers



### EVALUATION

**Choose the correct answer:**

1. An alcohol (x) gives blue colour in Victor Meyer's test and 3.7g of X when treated with metallic sodium liberates 560 mL of hydrogen at 273 K and 1 atm pressure what will be the possible structure of X?




2. Which of the following compounds on reaction with methyl magnesium bromide will give tertiary alcohol.

a) benzaldehyde




b) propanoic acid

c) methyl propanoate

d) acetaldehyde



The X is



d) None of these

4. In the reaction sequence, Ethene  $\xrightarrow{\text{HOCl}}$  A  $\xrightarrow{\text{X}}$  ethan -1, 2 - diol . A and X respectively are

a) Chloroethane and NaOH

b) ethanol and  $\text{H}_2\text{SO}_4$

c) 2 - chloroethan -1-ol and  $\text{NaHCO}_3$

d) ethanol and  $\text{H}_2\text{O}$

5. Which one of the following is the strongest acid


a) 2 - nitrophenol

b) 4 - chlorophenol

c) 4 - nitrophenol

d) 3 - nitrophenol

6.  on treatment with  $\text{Con H}_2\text{SO}_4$ , predominately gives



7. Carboxylic acid is


- a) Phenol                      b) Picric acid                      c) benzoic acid                      d) phenylacetic acid

8. Which one of the following will react with phenol to give salicylaldehyde after hydrolysis.

- a) Dichloro methane    b) trichloroethane    c) trichloro methane    d) CO<sub>2</sub>

9.  $(\text{CH}_3)_3\text{C}-\text{CH}(\text{OH})\text{CH}_3 \xrightarrow{\text{Con H}_2\text{SO}_4} \text{X}$  (major product)

- a)  $(\text{CH}_3)_3\text{CCH}=\text{CH}_2$                       b)  $(\text{CH}_3)_2\text{C}=\text{C}(\text{CH}_3)_2$   
 c)  $\text{CH}_2=\text{C}(\text{CH}_3)\text{CH}_2-\text{CH}_2-\text{CH}_3$                       d)  $\text{CH}_2=\text{C}(\text{CH}_3)-\text{CH}_2-\text{CH}_2-\text{CH}_3$

10. The correct IUPAC name of the compound, 

- a) 4-chloro-2,3-dimethylpentan-1-ol  
 b) 2,3-dimethyl-4-chloropentan-1-ol  
 c) 2,3,4-trimethyl-4-chlorobutan-1-ol  
 d) 4-chloro-2,3,4-trimethylpentan-1-ol

11. Assertion : Phenol is more acidic than ethanol


Reason: Phenoxide ion is resonance stabilized

- a) if both assertion and reason are true and reason is the correct explanation of assertion.  
 b) if both assertion and reason are true but reason is not the correct explanation of assertion.  
 c) assertion is true but reason is false  
 d) both assertion and reason are false.

12. In the reaction  $\text{Ethanol} \xrightarrow{\text{PCl}_5} \text{X} \xrightarrow{\text{alc.KOH}} \text{Y} \xrightarrow[298\text{K}]{\text{H}_2\text{SO}_4/\text{H}_2\text{O}} \text{Z}$ . The 'Z' is

- a) ethane                      b) ethoxyethane                      c) ethylbisulphite                      d) ethanol


13. The reaction



Can be classified as

- a) dehydration                      b) Williamson alcohol synthesis  
 c) Williamson ether synthesis                      d) dehydrogenation of alcohol

14. Isopropylbenzene on air oxidation in the presence of dilute acid gives  
 a)  $C_6H_5COOH$       b)  $C_6H_5COCH_3$       c)  $C_6H_5COC_6H_5$       d)  $C_6H_5 - OH$
15. Assertion : Phenol is more reactive than benzene towards electrophilic substitution reaction  
 Reason : In the case of phenol, the intermediate arenium ion is more stabilized by resonance.  
 a) if both assertion and reason are true and reason is the correct explanation of assertion.  
 b) if both assertion and reason are true but reason is not the correct explanation of assertion.  
 c) assertion is true but reason is false  
 d) both assertion and reason are false.
16.  $HOCH_2CH_2OH$  on heating with periodic acid gives  
 a) methanoic acid      b) Glyoxal      c) methanal      d)  $CO_2$
17. Which of the following compound can be used as antifreeze in automobile radiators?  
 a) methanol      b) ethanol      c) Neopentyl alcohol      d) ethan -1, 2-diol
18. The reactions



is an example of

- a) Wurtz reaction      b) cyclic reaction      c) Williamson reaction      d) Kolbe reactions
19. One mole of an organic compound (A) with the formula  $C_3H_8O$  reacts completely with two moles of HI to form X and Y. When Y is boiled with aqueous alkali it forms Z. Z answers the iodoform test. The compound (A) is  
 a) propan - 2-ol      b) propan -1-ol      c) ethoxy ethane      d) methoxy ethane
20. Among the following ethers which one will produce methyl alcohol on treatment with hot HI?  
 a)  $(H_3C)_3C-O-CH_3$       b)  $(CH_3)_2CH-CH_2-O-CH_3$   
 c)  $CH_3(CH_2)_3-O-CH_3$       d)  $CH_3-CH_2-\underset{\substack{| \\ CH_3}}{CH}-O-CH_3$
21. Williamson synthesis of preparing dimethyl ether is a / an /  
 a)  $SN^1$  reactions      b)  $SN^2$  reaction  
 c) electrophilic addition      d) electrophilic substitution
22. On reacting with neutral ferric chloride, phenol gives  
 a) red colour      b) violet colour      c) dark green colour      d) no colouration.



# Carbonyl Compounds & Carboxylic Acids



## EVALUATION

Choose the correct answer:

1. The correct structure of the product 'A' formed in the reaction (NEET)




2. The formation of cyanohydrin from acetone is an example of

- a) nucleophilic substitution                      b) electrophilic substitution  
 c) electrophilic addition                      d) Nucleophilic addition


3. Reaction of acetone with one of the following reagents involves nucleophilic addition followed by elimination of water. The reagent is

- a) Grignard reagent                      b) Sn / HCl  
 c) hydrazine in presence of slightly acidic solution                      d) hydrocyanic acid

4. In the following reaction,




- a) Tollen's test                      b) Victor meyer test  
 c) Iodoform test                      d) Fehling solution test



- a) Formaldehyde                      b) di acetone ammonia  
 c) hexamethylene tetraamine                      d) oxime







6. Predict the product Z in the following series of reactions



- a)  $(\text{CH}_3)_2\text{C}(\text{OH})\text{C}_6\text{H}_5$                       b)  $\text{CH}_3\text{CH}(\text{OH})\text{C}_6\text{H}_5$   
 c)  $\text{CH}_3\text{CH}(\text{OH})\text{CH}_2 - \text{CH}_3$                       d)

7. Assertion: 2,2 - dimethyl propanoic acid does not give HVZ reaction.

Reason: 2 - 2, dimethyl propanoic acid does not have  $\alpha$  - hydrogen atom

- a) if both assertion and reason are true and reason is the correct explanation of assertion.  
 b) if both assertion and reason are true but reason is not the correct explanation of assertion.  
 c) assertion is true but reason is false  
 d) both assertion and reason are false.
8. Which of the following represents the correct order of acidity in the given compounds  
 a)  $\text{FCH}_2\text{COOH} > \text{CH}_3\text{COOH} > \text{BrCH}_2\text{COOH} > \text{ClCH}_2\text{COOH}$   
 b)  $\text{FCH}_2\text{COOH} > \text{ClCH}_2\text{COOH} > \text{BrCH}_2\text{COOH} > \text{CH}_3\text{COOH}$   
 c)  $\text{CH}_3\text{COOH} > \text{ClCH}_2\text{COOH} > \text{FCH}_2\text{COOH} > \text{Br-CH}_2\text{COOH}$   
 d)  $\text{ClCH}_2\text{COOH} > \text{CH}_3\text{COOH} > \text{BrCH}_2\text{COOH} > \text{ICH}_2\text{COOH}$
9. Benzoic acid  $\xrightarrow[\text{ii) } \Delta]{\text{i) } \text{NH}_3}$  A  $\xrightarrow{\text{NaOBr}}$  B  $\xrightarrow{\text{NaNO}_2/\text{HCl}}$  C 'C' is  
 a) anilinium chloride  
 b) O – nitro aniline  
 c) benzene diazonium chloride  
 d) m – nitro benzoic acid
10. Ethanoic acid  $\xrightarrow{\text{P/Br}_2}$  2 – bromoethanoic acid. This reaction is called  
 a) Finkelstein reaction  
 b) Haloform reaction  
 c) Hell – Volhard – Zelinsky reaction  
 d) none of these
11.  $\text{CH}_3\text{Br} \xrightarrow{\text{KCN}} (\text{A}) \xrightarrow{\text{H}_2\text{O}^+} (\text{B}) \xrightarrow{\text{PCl}_5} (\text{C})$  product (c) is  
 a) acetylchloride  
 b) chloro acetic acid  
 c)  $\alpha$  - chlorocyano ethanoic acid  
 d) none of these
12. Which one of the following reduces tollens reagent  
 a) formic acid  
 b) acetic acid  
 c) benzophenone  
 d) none of these
13.   
 a)   
 b)   
 c)   
 d) 
14. The IUPAC name of 








a) but – 3- enoicacid

b) but – 1- ene-4-oicacid





c) but – 2- ene-1-oic acid

d) but -3-ene-1-oicacid

15. Identify the product formed in the reaction



16. In which case chiral carbon is not generated by reaction with HCN



17. Assertion : p – N, N – dimethyl aminobenzaldehyde undergoes benzoin condensation

Reason : The aldehydic (-CHO) group is meta directing

a) if both assertion and reason are true and reason is the correct explanation of assertion.

b) if both assertion and reason are true but reason is not the correct explanation of assertion.

c) assertion is true but reason is false

d) both assertion and reason are false.

18. Which one of the following reaction is an example of disproportionation reaction

a) Aldol condensation

b) cannizaro reaction



# Organic Nitrogen Compounds










## EVALUATION






Choose the correct answer:

- Which of the following reagent can be used to convert nitrobenzene to aniline
  - Sn / HCl
  - ZnHg / NaOH
  - LiAlH<sub>4</sub>
  - All of these
- The method by which aniline cannot be prepared is
  - degradation of benzamide with Br<sub>2</sub> / NaOH
  - potassium salt of phthalimide treated with chlorobenzene followed by hydrolysis with aqueous NaOH solution.
  - Hydrolysis of phenylcyanide with acidic solution
  - reduction of nitrobenzene by Sn / HCl.
- Which one of the following will not undergo Hofmann bromamide reaction
  - CH<sub>3</sub>CONHCH<sub>3</sub>
  - CH<sub>3</sub>CH<sub>2</sub>CONH<sub>2</sub>
  - CH<sub>3</sub>CONH<sub>2</sub>
  - C<sub>6</sub>H<sub>5</sub>CONH<sub>2</sub>
- Assertion : Acetamide on reaction with KOH and bromine gives acetic acid  
Reason : Bromine catalyses hydrolysis of acetamide.
  - if both assertion and reason are true and reason is the correct explanation of assertion.
  - if both assertion and reason are true but reason is not the correct explanation of assertion.
  - assertion is true but reason is false
  - both assertion and reason are false.
- CH<sub>3</sub>CH<sub>2</sub>Br  $\xrightarrow[\Delta]{\text{aq NaOH}}$  A  $\xrightarrow[\Delta]{\text{KMnO}_4/\text{H}^+}$  B  $\xrightarrow[\Delta]{\text{NH}_3}$  C  $\xrightarrow{\text{Br}_2/\text{NaOH}}$  D 'D' is
  - bromomethane
  - α - bromo sodium acetate
  - methanamine
  - acetamide
- Which one of the following nitro compounds does not react with nitrous acid
  - CH<sub>3</sub>-CH<sub>2</sub>-CH<sub>2</sub>-NO<sub>2</sub>
  - (CH<sub>3</sub>)<sub>2</sub>CH - CH<sub>2</sub>NO<sub>2</sub>
  - (CH<sub>3</sub>)<sub>3</sub>C NO<sub>2</sub>
  - $$\begin{array}{c} \text{CH}_3 - \text{C} - \text{CH} - \text{NO}_2 \\ \parallel \quad | \\ \text{O} \quad \text{CH}_3 \end{array}$$
- Aniline + benzoylchloride  $\xrightarrow{\text{NaOH}}$  C<sub>6</sub>H<sub>5</sub> - NH - COC<sub>6</sub>H<sub>5</sub> this reaction is known as
  - Friedel - crafts reaction
  - HVZ reaction
  - Schotten - Baumann reaction
  - none of these
- The product formed by the reaction an aldehyde with a primary amine (NEET)





- Page No:19*

- a) red solution                      b) blue solution                      c) green solution                      d) yellow solution
17. Which of the following amines does not undergo acetylation?
- a) t – butylamine                      b) ethylamine                      c) diethylamine                      d) triethylamine
18. Which one of the following is most basic?
- a) 2,4 – dichloroaniline                      b) 2,4 – dimethyl aniline
- c) 2,4 – dinitroaniline                      d) 2,4 – dibromoaniline
19. When  is reduced with Sn / HCl the pair of compounds formed are
- a) Ethanol, hydroxylamine hydrochloride                      b) Ethanol, ammonium hydroxide
- c) Ethanol, .NH<sub>2</sub>OH.                      d) C<sub>3</sub>H<sub>5</sub>NH<sub>2</sub>, H<sub>2</sub>O
20. IUPAC name for the amine
-  is
- a) 3 – Bimethylamino – 3 – methyl pentane
- b) 3 (N,N – Triethyl) – 3- amino pentane
- c) 3 – N,N – trimethyl pentanamine
- d) 3 – (N,N – Dimethyl amino) – 3- methyl pentane
21.  + CH<sub>3</sub>MgBr  $\xrightarrow{\text{H}_3\text{O}^+}$  P Product 'P' in the above reaction is
- a) 
- b) 
- c) 
- d) 
22. Ammonium salt of benzoic acid is heated strongly with P<sub>2</sub>O<sub>5</sub> and the product so formed is reduced and then treated with NaNO<sub>2</sub>/ HCl at low temperature. The final compound formed is
- a) Benzene diazonium chloride                      b) Benzyl alcohol
- c) Phenol                      d) Nitrosobenzene






23. Identify X in the sequence give below.



24. Among the following, the reaction that proceeds through an electrophilic substitution, is :



25. The major product of the following reaction









19. Which of the following statement is correct?
- a) Ovalbumin is a simple food reserve in egg-white
  - b) Blood proteins thrombin and fibrinogen are involved in blood clotting
  - c) Denaturation makes protein more active
  - d) Insulin maintains the sugar level of in the human body.
20. Glucose is an aldose. Which one of the following reactions is not expected with glucose?
- a) It does not form oxime
  - b) It does not react with Grignard reagent
  - c) It does not form osazones
  - d) It does not reduce tollens reagent
21. If one strand of the DNA has the sequence 'ATGCTTGA', then the sequence of complementary strand would be
- a) TACGAACT                      b) TCCGAACT                      c) TACGTACT                      d) TACGRAGT
22. Insulin, a hormone chemically is
- a) Fat                                      b) Steroid                                      c) Protein                                      d) Carbohydrates
23.  $\alpha$ -D (+) Glucose and  $\beta$ -D (+) glucose are
- a) Epimers                                      b) Anomers
  - c) Enantiomers                                      d) Conformational isomers
24. Which of the following are epimers
- a) D(+)-Glucose and D(+)-Galactose                      (b) D(+)-Glucose and D(+)-Mannose
  - c) Neither (a) nor (b)                      (d) Both (a) and (b)
25. Which of the following amino acids are achiral?
- a) Alanine                                      b) Leucine                                      c) Proline                                      d) Glycine

*By*

*M. Kesavan MSc.BEd.,*

*SVM Mat Hr Sec School...*

*Palacode...*




## Chemistry in Everyday Life




### EVALUATION

**Choose the correct answer:**

1. Which of the following is an analgesic?  
 a) Streptomycin                      b) Chloromycetin                      c) Asprin                      d) Penicillin
2. Dettol is the mixture of  
 a) Chloroxylenol and bithionol                      b) Chloroxylenol and  $\alpha$ -terpineol  
 c) phenol and iodine                      d) terpineol and bithionol
3. Antiseptics and disinfectants either kill or prevent growth of microorganisms. Identify which of the following statement is not true.  
 a) dilute solutions of boric acid and hydrogen peroxide are strong antiseptics.  
 b) Disinfectants harm the living tissues.  
 c) A 0.2% solution of phenol is an antiseptic while 1% solution acts as a disinfectant.  
 d) Chlorine and iodine are used as strong disinfectants.
4. Saccharin, an artificial sweetener is manufactured from  
 a) cellulose                      b) toluene                      c) cyclohexene                      d) starch
5. Drugs that bind to the receptor site and inhibit its natural function are called  
 a) antagonists                      b) agonists                      c) enzymes                      d) molecular targets
6. Aspirin is a/an  
 a) acetylsalicylic acid    b) benzoyl salicylic acid    c) chlorobenzoic acid    d) anthranilic acid
7. Which one of the following structures represents nylon 6,6 polymer?





8. Natural rubber has

- a) alternate cis- and trans-configuration      b) random cis- and trans-configuration  
c) all cis-configuration      d) all trans-configuration

9. Nylon is an example of

- a) polyamide      b) polythene      c) polyester      d) poly saccharide

10. Terylene is an example of

- a) polyamide      b) polythene      c) polyester      d) polysaccharide

11. Which is the monomer of neoprene in the following?

- a)  $\text{CH}_2 - \underset{\text{Cl}}{\text{C}} - \text{CH} = \text{CH}_2$       b)  $\text{CH}_2 = \text{CH} - \text{C} \equiv \text{CH}$   
c)  $\text{CH}_2 = \text{CH} - \text{CH} = \text{CH}_2$       d)  $\text{CH}_2 = \underset{\text{CH}_3}{\text{C}} - \text{CH} = \text{CH}_2$

12. Which one of the following is a bio-degradable polymer?

- a) HDPE      b) PVC      c) Nylon 6      d) PHBV

13. Non stick cook wares generally have a coating of a polymer, whose monomer is

- a) ethane      b) prop-2-enenitrile      c) chloroethene      d) 1,1,2,2-tetrafluoroethane

14. Assertion: 2-methyl-1,3-butadiene is the monomer of natural rubber

Reason: Natural rubber is formed through anionic addition polymerisation.

- a) If both assertion and reason are true and reason is the correct explanation of assertion.  
b) if both assertion and reason are true but reason is not the correct explanation of assertion.  
c) assertion is true but reason is false.      d) both assertion and reason are false.

15. An example of antifertility drug is

- a) novestrol      b) seldane      c) salvarsan      d) Chloramphenicol

16. The drug used to induce sleep is

- a) paracetamol      b) bithional      c) chloroquine      d) equanil

17. Which of the following is a co-polymer?

- a) Orlon      b) PVC      c) Teflon      d) PHBV

18. The polymer used in making blankets (artificial wool) is

- a) polystyrene      b) PAN      c) polyester      d) polythene

19. Regarding cross-linked or network polymers, which of the following statement is incorrect? (NEET)
- a) Examples are Bakelite and melamine
  - b) They are formed from bi and tri-functional monomers
  - c) They contain covalent bonds between various linear polymer chains
  - d) They contain strong covalent bonds in their polymer chain
20. A mixture of chloroxylenol and terpinecol acts as (NEET)
- a) antiseptic
  - b) antipyretic
  - c) antibiotic
  - d) analgesic

Padasalai

*I never learn anything talking. I only  
learn things when I ask questions...*

*- Lou Holtz*